


MARYLAND & RARE BOOK ROOM UNIVERSITY OF MARYLAND LIBRARY COLLEGE PARK, MD.

LIBRARY-COLLEGE PARK



Digitized by the Internet Archive in 2010 with funding from Lyrasis Members and Sloan Foundation



3

# ANNUAL REPORT

OF THE

# State Board of Health

OF MARYLAND

FOR THE

YEAR ENDING DECEMBER 31, 1915





Baltimore: KING BROTHERS State Printers 413° E. Lexington Street 1918 .

# State Department of Health of Maryland, 1915.

# STATE BOARD OF HEALTH.

Dr. WM. H. WELCH	. President
Dr. JOHN S. FULTON Secretary and Executive	tive Officer
Dr. WM. W. FORD.	
Dr. NATHAN R. GORTER.	
Dr. JOHN D. BLAKEComm. of Health of Baltimore (	ex officio)
Hon. EDGAR ALLAN POE. Attorney General of Maryland (	ex officio)

# EXECUTIVE OFFICE.

Walter N. Kirkman E. Robinson	
M. L. Wiedefeld	Chief Clerk's Stenographer
J. M. Conway and J. F. McLain	Multigraph Operators
A. E. LENZ	
N. WILKINSON, A. REICHERT and E. POFFE	LStock Clerks
B. MALEY	Telephone Operator
General Counsel.	Special Counsel,
WM. PINKNEY WHYTE, JR.	H. M. McCullough.

# BUREAU OF BACTERIOLOGY.

Dr. Wm. Royal Stokes	
Dr. Harry W. Stoner	Assistant Chief
Dr. Wm. Greenfeld	
R. L. SULLIVAN	
C. McCrackan, L. Egan, E. Cresap	

#### BUREAU OF CHEMISTRY.

Dr. W. B. D. Penniman	
DR. WYATT W. RANDALL	Assistant Chief Chemist
C. O. MILLER and L. H. ENSLOW	
E. HOPKINS and R. BREYER	
M. E. SEIDELL	

# BUREAU OF COMMUNICABLE DISEASES.

Dr. C. Hampson Jones	
Dr. C. W. G. Rohrer	
M. A. MILLER	
J. J. KING	
M. EURICH and O. PORTER	
A. Dushane	Tuberculosis Registration Clerk

# BUREAU OF SANITARY ENGINEERING.

Robert B. Morse	
H. R. HALL	Assistant Chief Engineer
F. W. CASPARI	Special Assistant Engineer
G. L. HALL	Office Division Engineer
R. T. Davis	
J. F. Diggs, Jr	
A. WOLMAN	
J. S. Gorsuch, Jr	
A. E. GOODRICH	Resident Engineer
F. P. GILBERT	
C. R. Cox	
C. E. Fox	
H. R. DEVILBISS	
H. L. WEAVER	Draftsman
C. M. FANKHANEL	
D. R. Makibbin	
H. A. WALKER	
F. E. HILDEBRANDT	
H. P. DELANEY	
M. E. Dodson	
E. C. Hughes	

# BUREAU OF VITAL STATISTICS.

Dr. Frederic V. Beitler	
T. NICKLAS, M. BAER, E. IJAMS	
E. Scott, R. Gees, H. Noel, M. Wilson, M. Gosline	
B. G. BetzTabulating Clerk	

# DIVISION OF FOOD AND DRUGS.


Dr. Chas. Caspari, JrState Food and Drug Commissioner
R. W. BatemanStenographer
J. W. ARNOLD, R. H. HOPKINS, C. E. TAYLOR. Food and Drug Inspectors
Dr. W. B. Billingsley
A. K. Beasley
C. N. MITTEN

# DEPUTY STATE HEALTH OFFICERS.

Dr. R. H. Riley
W. A. Gunther
E. BergmanLaboratory Assistant
Dr. R. B. Norment, Jr
Dr. W. C. Stone3rd Sanitary District
A. W. Stone
Dr. J. S. Bowen
Dr. Caldwell Woodruff
J. E. HAYESOffice Boy
V. E. Robertson
Dr. E. A. Jones
M. J. BryanStenographer
Dr. A. McC. Stevens

# **CONTENTS**

List of Officers and Employees	1-2
Report of the Chief of the Bureau of Vital Statistics	5-118
Report of the Chief of the Bureau of Communicable Diseases	119-163
Report of the Chief of the Bureau of Bacteriology	164-165
Report of the Chief of the Bureau of Chemistry	166-169
Report of the State Food and Drug Commissioner	170-199
Report of the Chief of the Bureau of Sanitary Engineering	200-230
Report of the Chief Clerk	231-234



# Report of the Bureau of Vital Statistics.

FREDERIC V. BEITLER, Chief

# POPULATION, BIRTHS AND DEATHS.

The efforts of Boards of Health and sanitarians are directed toward increasing the birth rate, diminishing the death rate and increasing the duration of life. As a basis upon which to direct their efforts in the State of Maryland, the tables contained within this report have been devised. All the earlier tables will be maintained and others added from time to time.

All Vital Statisticians use the Census figures of the population. The figures for census years, of course, are most valuable. In inter-census years, the population is computed by the arithmetic method.

The returned mortality in Maryland probably now constitutes 100 per cent. of the total deaths. Birth registration constitutes about 98 per cent. of the total births. Maryland statistics show in this respect the deficiency existing in the majority of the American States in the registration of births.

The deaths registered in Maryland during 1915 were, for \*Rural Maryland, 12,023; in Baltimore City, 9,327; making a total for the State of 21,350.

\* The births reported in 1915 were, for Rural Maryland, 18,659; for Baltimore City, 13,708; making a total for the State of 32,367, giving an apparent increase in the population of the State of 11,017.

The figures for Maryland during 1914 were: births 30,652, deaths 21,383, an increase of 9,269 births over deaths. The 1915 figures show an increase of 1,715 births over those registered during 1914.

# Population of Maryland—1915.

The figures for the population for the inter-census years are estimated by the arithmetic method as advised by the American Public Health Association, and are corrected as of July 1st of each year. The yearly increment is given along with the figures

<sup>\*</sup>The term Rural Maryland which is used in this report destinguishes that part of the State outside of the corporate limits of Baltimore City from the city proper. We are aware that the term is somewhat misleading, but has been used in this report for a number of years for that purpose.

for each division of the population, so that persons wishing the use of figures for this State for any inter-census year may do so without having to estimate them.

The population of Maryland is given in Table 1 for the male, female, white and colored population of Baltimore City, Rural Maryland and the State of Maryland.

The estimated population, white and colored, for the counties appears in Table 1-A. The estimated population by ages, without distinction of sex or color, appears in Table 19-A.

TABLE I.

ESTIMATED POPULATION IN MARYLAND FOR THE YEAR 1915.

Maryland.	Population.	Yearly In	crement.
Total population	. 1,351,940	+10,8	366.0
White		+11,	
Colored	004 4 = 4		295.0
Male	.=0.00=	+5.5	564.5
Female			301.5
Baltimore City.			
Total population	. 584,608	+5,0	015.5
White		+4,	472.8
Colored		+	542.7
Rural Maryland.			
Total population	. 767.332	<b>1</b> 5	850.5
White			688.2
Colored			837.7
Colored	. 145,240		001.1
White	Yearly C	olored	Yearly
Maryland. Population. In	crement. Pop	ulation. In	crement.
	-5,666.1 11	14.624	101.6
	-,	, -	_193.4
1 chares 602,100	0,202.0	.0,021	
Baltimore City.			
	-2,133.6	11.408	+389.5
			$\pm 153.2$
2 0000000 11111111111111111111111111111	,	,	•
Rural Maryland.			
	-3,532.5	73,216	_491.1
		70,030	_346.6
· ·			

TABLE I-A.

ESTIMATED POPULATION FOR THE YEAR 1915.

	W	White.		Colored.		Total.	
	Popula-	Incre-	Popula	- Incre-	Popula-	Incre-	
County.	tion.	ment.	tion.	ment.	tion.	ment.	
Allegany	65,573	+898.6	1,435	-15.9	67,008	+882.7	
Anne Arundel	26,008	+117.5	13,510	124.2	39,518	6.7	
Baltimore	125,877	+3.099.7	13,135	+99.6	139,012	+3,199.3	
Calvert	5,384	+20.1	4,995	_9.8	10,379	+10.3	
Caroline	15,702	+244.9	5,079	+55.7	20,781	+300.6	
Carroll	32,029	+20.7	1,945	_13.2	33,974	+7.5	
Cecil	20,201	_43.0	3,082	-48.4	23,283	-91.4	
Charles	7,707	-20.3	8,006	108.9	15,713	_129.2	
Dorchester	19,654	+78.1	9,388	-6.5	29,042	+71.6	
Frederick	47,993	+138.4	5,077	-62.2	53,070	+76.2	
Garrett	21,276	+245.4	97	<b>—1.9</b>	21,373	+243.5	
Harford	23,080	+44.4	4,725	75.1	27,805	30.7	
Howard	12,344	+2.3	3,441	64.0	15,785	<u>61.7</u>	
Kent	10,506	-55.5	$5,\!486$	-129.7	15,992	-185.2	
Montgomery	24,140	+248.4	8,813	-82.5	32,953	+165.9	
Prince George's	28,194	+681.8	11,249	_49.0	39,443	+632.8	
Queen Anne's	10,512	98.0	$5,\!522$	-56.4	16,034	154.4	
Somerset	17,290	+59.8	9,446	6.0	26,736	+53.8	
St. Mary's	10,148	+81.0	$6,\!802$	-96.4	16,950	-15.4	
Talbot	12,823	3.4	$6,\!416$	<b>—</b> 69.7	19,239	<b>—73.1</b>	
Washington	50,057	+491.6	1,924	-37.6	51,981	+454.0	
Wicomico	22,340	+352.5	6,565	+48.8	28,905	+401.3	
Worcester	15,248	+83.2	7,108	+15.6	22,356	+98.8	
Total Counties	624,086	+6,688.2	143,246	_837.7	767,332	+5,850.5	
Baltimore City	496,683	+4,472.8	87,925	+542.7	584,608	+5,015.5	
Total Maryland	1,120,769	+11,161.0	231,171	_295.0	1,351,940	+10,866.0	

# Births in Maryland.

The number of births recorded in the State of Maryland during 1915 was 32,367, of which 18,659 were reported from Rural Maryland, and 13,708 from Baltimore City.

The number of white births reported was 26,126. The number of colored births reported was 6,241.

The total number of male births, 16,541; of female births, 15,826.

Table II gives the birth rates per thousand of population for Baltimore City, Rural Maryland and the State of Maryland, for the years 1905 to 1915, inclusive. The birth rate for Baltimore City was 23.45 for the year 1915, for Rural Maryland 24.32, and for the entire State 23.94. The increase in the year 1912 for both Rural Maryland and Baltimore City was due to the new registration law which went into effect on July 1st of that year.

TABLE II.
BIRTH RATES, MARYLAND, 1905-1915.

i	Birth Rates, Rural Maryland	Birth Rates, Baltimore City	Birth Rates, Maryland
1905	13.15	16.75	14.70
1906	13.72	16.87	15.08
1907		16.09	14.31
1908	14.07	16.70	15.20
1909	14.80	15.86	15.26
1910	14.51	17.62	15.85
1911		16.44	15.17
1912		20.01	18.67
1913		21.83	21.34
1914		21.80	22.86
1915	0.1.00	23.45	23.94

In Table III and III-A, the births, deaths and increase are given by counties for the whole population, and separately for white and colored races; also the birth rate, death rate and rate of increase per 1,000 of the population (as estimated from the United States Census figures for 1910).

# TABLE III.

BIRTH RATES, DEATH RATES AND RATE OF INCREASE, 1915.

۰		White.		1	Colored			Total.	
	Births	Deaths	Inc. or	Births	Deaths	Inc. or	Births	Deaths	Inc. or
	per	per	Dec. per	per	per	Dec. per	per	per	Dec. per
Counties.	1,000.	1,000.	1,000.	1,000.	1,000.	1,000.	1,000.	1,000.	1,000.
	26.44	11.79	+14.65	34.15	34.15	00.0	26.61	12.27	+14.34
Anne Arundel.	29.26	14.03	+15.23	30.57	27.76	+2.81	29.71	18.73	+10.98
	21.03	17.24	+3.79	16.98	40.35	23.37	20.65	19.42	+1.23
	17.64	10.22	+7.42	30.23	19.62	+10.61	23.70	14.74	+8.96
	23.18	11.72	+11.46	27.17	19.69	+7.48	24.16	13.67	+10.49
	23.23	16.33	+6.90	24.68	22.11	+2.57	23.31	16.66	+6.65
	21.29	15.44	+5.85	16.55	19.79	-3.24	20.66	16.02	+4.64
	26.08	12.72	+13.36	35.72	21.11	+14.61	30.99	16.99	+14.00
	26.31	13.74	+12.57	37.18	24.71	+12.47	29.85	17.29	+12.53
	23.69	13.61	+10.08	29.35	25.01	+4.34	24.23	14.70	+9.53
	25.85	10.72	+15.13	0.00	0.00	0.00	25.73	10.67	+15.06
	18,15	12.44	+5.71	16.72	21.80	-5.08	17.91	14.03	+3.88
	19.04	10.21	+8.83	31.10	21.21	+9.89	21.67	12.61	+9.06
	19.32	12.28	+7.04	23.88	26.43	2.55	20.89	17.13	+3.76
	20.17	11.64	+8.53	25.53	17.59	+7.94	21.61	13.23	+8.38
ge's	21.92	10.53	+11.39	30.67	20.09	+10.58	24.41	13.26	+11.15
8.0	24.54	13.60	+10.94	26.26	23.54	+2.72	25.13	17.03	+8.10
	24.00	13.53	+10.47	30.28	17.57	+12.71	26.22	14.97	+11.25
	25.13	11.23	+13.90	31.90	22.93	+8.97	27.85	15.93	+11.92
	22.85	13.65	+9.20	23.38	21.66	+1.72	23.03	16.32	+6.71
	29.45	13.94	+15.51	19.23	27.55	8.35	29.02	14.45	+14.62
	21.89	12.76	+9.13	31.68	21.17	+10.51	24.11	14.67	+9.44
	22.23	14.30	+7.93	29.12	19.13	+9.99	24.42	15.83	+8.59
	23.50	13.81	+9.69	27.88	23.77	+4.11	24.32	15.67	+x.65
Baltimore City	23.08	14.39	+8.69	25.56	24.77	+0.79	23.45	15.95	+4.50
-	1 8		100	00	7 7 7 6	200	00 00	18.70	0 -
Maryland	23.31	14.07	+9.24	27.00	61.42	+2.39	79.94	19.13	

TABLE III-A.
BIRTHS, DEATHS AND INCREASE OF POPULATION, 1915.

		White.			Colored.			Total.	
			Inc. or			_			Inc. or
Counties	Births.		Dec.		Deaths.		Births.	Deaths.	Dec.
Allegany			+961		49		1,783	822	+961
Anne Arundel			+396		375		1,174	740	+434
Baltimore			+477		530		2,870	2,700	+170
Calvert			+40		86		246	153	+93
Caroline			+180		100		205	284	+218
Carroll			+221		43		792	566	+226
Cecil			+118		61		481	373	+108
Charles			+103		169		487	267	+220
Dorchester			+247		232		998	502	+364
Frederick			+484		127		1,286	282	+206
Garrett			+322		0		550	228	+322
Harford			+132		103		498	390	+108
Howard			+109		73		342	199	+143
Kent			+74		145		334	274	09+
Montgomery			+206		155		712	436	+276
Prince George's			+321		526		963	523	+440
Queen Anne's			+115		130		403	273	+130
Somerset			+181		166		701	400	+301
St. Mary's			+141		156		472	270	+202
Talbot			+118		139		443	314	+129
Washington			+776		53		1,511	751	+760
Wieomico			+204		139		697	424	+273
Worcester			+121		136		546	354	+192
			1000		107 6		10.000	10000	1000
Politimone City			+0,047		0,400		18,009	12,023	+0,030
			+4,312		2,178		13,708	9,327	+4,381
Maryland '	26,126		+10,359	6,241	5,583	+658	32,367	21,350	+11,017

By reference to the part of the table below-dealing with birth rates, death rates and rates of increase per thousand, a comparison may be made of the efficiency of registration in the various counties of Maryland. The birth rate per 1,000 for Maryland probably lies between the figures 24 and 27. It can be assumed, therefore, that returns from any of the counties of Maryland, which are below 20 per 1,000, indicate defective returns, and a birth rate of 15 or less means very defective returns.

The death rate exceeds 15 per 1,000 in eleven counties. The lowest death rate is in Garrett County, 10.67 per 1,000. This county has almost entirely a white population, and its returns to this office are good. The death rate here given we believe is correct.

Table IV gives a summary of the births, birth rates, deaths, death rates, and excess of births over deaths per thousand among male, female, white and colored for the total population of Rural Maryland and of Baltimore City. The birth rates and death rates per 1,000 are calculated both in relation to the total population and to the four divisions of population, male, female, white and colored.

\* TABLE IV. Births, Déaths and Rates—Maryland—1915.

SUMMARY.

#### Births. Male. Female. White. Colored. Total. Rural Maryland..... 9.513 9,146 14,665 3,994 18,659 2,247 Baltimore City..... 7,028 6,680 11,461 13,708 Maryland ...... 16,541 15,826 26,126 6,241 32,367 Deaths. Rural Maryland..... 6,386 5.637 8.618 3,405 12.023 4.835 Baltimore City..... 4,492 7,149 2,178 9,327 Maryland ..... 11,221 10,129 15,767 5,583 21,350 Birth Rate (Computed on Total Populations). Rural Maryland..... 12.40 11.92 19.11 5.21 24.32 Baltimore City..... 12.02 11.43 19.60 3.85 23.45 Maryland ..... 12.23 11.71 19.32 4.62 23.94 Death Rate (Computed on Total Populations). Rural Maryland..... 8.32 7.35 11.23 4.44 15.67 8.27 Baltimore City..... 7.68 12.23 3.72 15.95 Maryland ..... 8.30 7.4911.66 4.13 15.79

Difference Between Birth and Death Rates. Rural Maryland Baltimore City Maryland	$+4.08 \\ +3.75$	Female. +4.57 +3.75 +4.22	White. +7.88 +7.37 +7.66	Colored. +0.77 +0.13 +0.49	Total. +8.65 +7.50 +8.15
Birth Rate (Computed on Specific Groups of Population).					
Rural Maryland Baltimore City	$24.28 \\ 24.98$	24.36 $22.03$	$23.50 \\ 23.08$	$27.88 \\ 25.56$	24.32 $23.45$
Maryland	24.56 $24.57$	23.32	23.31	$\frac{25.56}{27.00}$	23.43
Death Rate (Computed on Specific Groups of Population).					
Rural Maryland	16.30	15.01	13.81	23.77	15.67
Baltimore City Maryland	17.19 $16.67$	14.81 $14.92$	$14.39 \\ 14.07$	$24.77 \\ 24.15$	15.95 15.79
Difference Between Birth and Death Rates.	20101	22,02	22101		20110
Rural MarylandBaltimore CityMaryland		+9.35  +7.22  +8.40	+9.69 $+8.69$ $+9.24$	+4.11  +0.79  +2.85	$+8.65 \\ +7.50 \\ +8.15$

The succeeding tables (Tables V and VI) give births and still-births for Rural Maryland and Baltimore City.

For the State, there were registered in 1915, 32,367 living births and 2,777 still-births, a proportion of 8.58 per cent. of all births stillborn.

In Table V the male and female births are given by months, with the corresponding months of conception.

The greatest number of living births in Rural Maryland was recorded in August (1,707), the corresponding period of conception being the month of November.

During 1914 the maximum number of births in Rural Maryland occurred during the month of September (1,670).

The minimum number of births during 1915 in Rural Maryland occurred in November (1,357). During 1914, February furnished the minimum number of births.

The greatest number of male births in Rural Maryland occurred in March (866). The greatest number during 1914 occurred in September (837). The greatest number of female births in 1915 occurred in August (846). The greatest number in 1914 occurred in September (833). The male births exceeded

the female births in every month except April, July, September and October.

Table VI (still-births) shows practically no seasonal variation, the fluctuation being proportionate only to the number of total births.




CHART 1-Births by months, Male, Female and Total Maryland-1915.

ABLE V.

BIRTHS, MALE AND FEMALE, BY MONTHS, MARYLAND, 1915.

Month		R	ural Maryl	and.	В	altimore Ci	ty.		Maryland.	_
Month of Birth	Conception.	Male.	Female.	Total.	Male.	Female.	Total.	Male.	Female.	Total.
January		827	782	1,609	601	522	1,123	1.428	1,304	2,732
February		777	740	1,517	587	545	1,129	1,364	1,282	2,646
March		998	778	1,644	009	621	1,221	1,466	1,399	2,865
April		790	798	1,588	208	510	1,018	1,298	1,308	2,606
May		763	730	1,493	, 549	505	1,054	1,312	1,235	2,547
June		296	748	1,544	620	523	1,143	1,416	1,271	2,687
July		784	608	1,593	641	628	1,269	1,425	1,437	2,862
August		861	846	1,707	589	592	1,181	1,450	- 1,438	2,888
September		815	816	1,631	583	584	1,167	1,398	1,400	2,798
October		763	773	1,536	616	522	1,138	1,379	1,295	2,674
November		731	626	1,357	275	561	1,103	1,273	1,187	2,460
December		740	. 700	1,440	592	570	1,162	1,332	1,270	2,602
Total		9,513	9,146	9,513 9,146 18,659	7,028	7,028 6,680 13,708	13,708	16,541	16,541 15,826 32,367	32,367

TABLE VI.

# STILL-BIRTHS-MARYLAND, 1915.

	3. Total. 494 315 342 		2. Total	2. Total. 1,639
	Dec. 30 40 60 80 80 80 80 80 80 80 80 80 80 80 80 80	Dec. 22 13 14 42 42	. : : : : : : : : : : : : : : : : : : :	Dec. 141
	Nov. 42 16 22 28 80	Nov. 18 15 4 4 37	Nov. :::   :	Nov. 117
	0ct. 42 28 33 103	00ct.	oet: ::   :	Oct. 141
	Sept. 30 34 32 32 36 96	Sept.  17  14  5  36	Sept.	Sept. 132
	Aug. 30 18 34 82	Aug. 18 16 3 3 37	Aug	Aug. 119
	July. 50 27 32 109	July. 17 19 5	July	July. 151
ND.	June. 34 26 21 81	June. 16 12 5 33		June. 114
MARYLAND.	May. 45 26 23 23 94	May. 23 13 3 39	May. :: ::	May. 133
RURAL	$\frac{Apr.}{49}$	Apr. 20 20 118 10 10 48	Apr.	<i>Apr.</i> 141
	Mar. 41 21 33 33 95	Mar. 27 22 7 7 7 56	Mar.	Mar. 151
	Feb. 39 28 28 28 95	$Feb.$ 16 19 3 $\times$ 38	Feb	Feb. 133
	Jan. 52 37 35 35 124	Jan. 19	Jan	Jan 166
				:
	Male Female Unknown	Male Female Unknown .	Male	Maryland
	White.	Colored.	Color Unknown.	Total Rural

TABLE VI-Continued.

					BALTIM	BALTIMORE CITY.	Ý.							
		Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Totai.
	( Male	26	28	38	47	32	36	37	23	20	34	63	30	389
	Female	6.	22	16	19	55	27	15	25	19	19	83	55	248
White.	Unknown	14	∞ ∞	12	12	12	œ	11	17	14	2	œ	œ	131
			1	1	1			İ	1	1	8	3	8	000
	[ Total	29	28	99	28	99	99	63	65	3	99	69	3	2
		,				3.6	Trees	Tortor	4310	Sont	Oct	Wow.		Total
		Jan.	F'cb.	Mar.	Apr.	May.	June.	July.	Aug.	sept.	000			anao r
	f Male	18	15	18	13	16	16	18	15	14	გ ე	14	-	196
	Female	12	17	19	11	ಸಾ	15	10	11	10	12	13	16	150
Colored.	Unknown	:	:	:	-	-	ಣ	<b>c</b> 1	ಣ	c1	4	က		20
			ļ			1				1	1			
	Total	30	32	37	25	22	34	30	20	56	36	29	36	366
		Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Scpt.	Oct.	Nov.	Dcc.	Total.
	( Male	:	:	:	:	:	:	:	:	:	:	:	:	:'
7	Female	:	:	:	:	:	:	:	<del>, -</del>	:	:	:	:	<b>⊣</b> •
Cotor	Unknown	:	61	:	:		:	:	:	:	:	:	:	3
O reknotore.		1	1	1	1	1	1		١,	ĺ	1	1		
	[ Total	:	2	:	:	H	:	:	-	:	:	:	:	4
		Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Total.
Total Baltimore	imore City	83	92	103	103	68	100	93	35	88	96	94	96	1,138

TABLE VI-Continued.

	Total.	883	563	473		,919	stal.	432	342	79	1	853	Total.	:	٦,	4		က	Total.	,777
	Dec.	2	22	37		159	Dec	4	83	œ		28	Dec.	:	:	:		:	Dec.	237
	Nov.	11	4	ဓ္က	1	145	Nov.	35	22	2		99	Nov.	:	:	:		:	Nov.	211
	Oct.	92	47	40		163	Oct.	43	23	∞		74	Oct.	:	:	:		:	Oct.	237
	Sept.	29	53	46	1	158	Sept.	31	24	2		62	Sept.	:	:	:	1	:	Sept.	550
	Aug.	53	43	21	}	147	Aug.	33	22	9	1	99	Aug.	:	_	:		<del>,</del>	Aug.	214
	July.	87	45	43	1	172	July.	35	53	2		11	July.	:	:	-		-	July.	244
	June.	70	48	53	1	147	June.	35	27	00		29	June.	:	:	:			June.	214
MARYLAND.		2.2					May.	33	18	4		19	May.	:		1	1	-	May.	222
MAR		96				171				11			Apr.	:	:	:	1	:	Apr.	244
	Mar.	62	37	45	1	161	Mar.	45	41	2	1	93	Mar.	:	:	:		:	Mar.	254
	Feb.	67	20	36	1	153	Feb.	31	36	က		20	Feb.	:	•	01		61	Feb.	225
	Jan.	48	56	49	1	183	Jan.	37	33	က	1	7.5	Jan.	:	:	:		:	Jan.	255
			Female			Total		C Male				[ Total		( Male	Female	Unknown		[ Total		
_				White.						Colored.					Color	Coton	Ournoun.			Total Maryland

327 illegitimate stillbirths; 39 cases of twins; 1 case of triplets.

The succeeding table (Table VII) gives illegitimate births by counties. These figures include probably all of the returns of white illegitimates and a large number of the colored illegitimates; but, owing to the peculiar marital relations of the colored race, it is difficult to establish a standard of legitimacy for children born of colored parents. Ecclesiastical marriages are not performed for a large proportion of colored persons who are living as man and wife. In 1914, among the whites in Rural Maryland, there were 18.71 illegitimates per 1,000 white births and in 1915, 18.14. In the colored population, Rural Maryland in 1914, there were 182.15 illegitimates per 1,000 colored births and in 1915, 184.53 per 1,000 births.

TABLE VII. .

ILLEG	ITIMATE	Births—Ma	RYLAND-	1915.  Total	Illegiti- mate Births per 1,000
Counties	White.	Colored.	Total.	Births.	of Births.
Allegany	42	9	51	1,783	28.60
Anne Arundel	6	64	70	1,174	59.63
Baltimore	26	25	51	2,870	17.77
Calvert	<b>2</b>	30	32	246	130.08
Caroline	9	38	47	502	93.63
Carroll	16	12	28	792	35.35
Cecil	14	16	30	481	62.37
Charles	4	45	49	487	100.62
Dorchester	19	74	93	866	107.39
Frederick	20	41	61	1,286	47.43
Garrett	18		18	550	32.73
Harford	6	17	23	498	46.18
Howard	1	19	20	342	58.48
Kent	<b>2</b>	38	40	334	119.76
Montgomery	<b>2</b>	39	41	712	57.58
Prince George's	5	39	44	963	45.69
Queen Anne's	3	30	33	403	81.89
Somerset	8	46	54	701	77.03
St. Mary's	$^2$	33	35	472	74.15
Talbot	. 3	39	42	443	94.81
Washington	44	8	52	1,511	34.41
Wicomico	. 8	· 35	43	697	61.69
Worcester	6	40	46	546	84.25
				<del></del>	
Total Counties	266	737	1,003	18,659	53.75
Baltimore City	356	558	914	13,708	66.68
Total Maryland	622	1,295	1,917	32,367	59.23

PERCENTAGE OF NATIVE AND FOREIGN BORN PARENTS OF CHILDREN BORN IN MARYLAND DURING 1915.

TABLE VIII.

# Parent Nativity.

Both   One   Neither   Total   Both     Allegany   Auryland, Maryland, Maryland, Natives, Foreign     Allegany   44:59   28.44   17.11   90.14   5.55     Anne Arundel   65.09   10.14   4.51   80.83   14.57     Baltimore   66.18   10.14   4.51   80.83   14.57     Caroline   62.55   24.4   0.41   100.00   0.00     Caroline   62.55   24.4   0.41   98.23   1.01     Caroline   62.55   25.10   7.97   95.62   2.59     Caroline   62.55   22.45   6.24   98.23   1.01     Caroline   64.45   22.45   6.24   97.94   1.44     Dorchester   87.99   7.85   2.08   97.92   0.69     Frederick   78.77   14.93   5.52   99.22   0.31     Garrett   75.30   15.86   5.62   99.73   1.75     Harrord   78.95   13.45   97.37   1.75     Howard   78.95   13.45   97.90   0.60     Howard   78.95   13.45   99.23   0.00     Went   66.57   15.73   14.61   90.91   1.40     Prince George's   66.57   15.75   2.85   98.85   0.00     Washington   67.11   25.28   96.89   0.43     Worcester   68.86   20.70   7.83   96.89   2.38     Total Counties   69.11   16.72   7.20   98.73     Total Counties   69.11   16.72   7.20   98.75     Total Marriand   61.40   7.55   7.50   7.50     Total Marriand   61.40   7.55   7.50   7.50     Total Marriand   61.40   7.51   7.50   7.50   7.50     Total Marriand   61.40   7.51   7.50   7.50   7.50     Total Marriand   61.40   7.51   7.50	- `		Nati	ve.			Fore	ign.		
Maryland, Maryland, Maryland, Natives.       1el.     66.18       66.18     10.14     4.51       66.18     10.14     4.51       66.18     10.14     4.51       66.18     10.14     4.51       66.19     15.02     5.37       85.10     2.44     0.41       62.55     25.10     7.97       85.10     9.72     3.41       98.23     6.24     97.94       87.99     7.82     2.08       87.99     7.82     2.08       78.77     14.93     5.62       78.77     14.93     5.62       78.77     14.93     5.62       78.77     14.93     5.62       78.74     14.93     5.62       78.75     13.45     97.37       78.31     14.61     96.91       78.32     17.65     20.04       88.74     11.55     2.85       88.62     6.36     0.85       88.62     6.36     0.85       98.85     98.85       98.86     98.65       98.87     98.85       98.89     68.86     2.87       98.89     68.86     2.87     98.99       98.89     68.86		Both	One	Neither	Total	Both	Father	Mother	Total	Both
10.14   44.59   28.44   17.11   90.14     66.18   10.14   4.451   80.83     65.09   15.02   5.37   85.48     67.15   2.44   0.41   100.00     62.55   25.10   7.37   95.62     85.10   9.72   3.41   90.23     85.10   9.72   3.41   98.23     87.99   8.21   1.64   97.94     88.09   8.21   1.64   97.94     88.09   8.21   1.64   97.95     78.77   14.93   5.52   99.22     78.77   14.93   5.52   99.23     78.85   13.45   20.04   94.28     88.45   11.55   20.04   94.28     88.45   11.74   4.06   97.06     88.26   6.36   0.35   96.83     98.27   11.74   4.06   97.06     98.28   11.74   4.06   97.06     98.29   11.74   16.72   7.20   93.73     98.20   11.74   7.22   7.216     98.20   14.04   7	County.	Haryland.	Maryland.	Maryland.	Natives.	Foreign.	Foreign.	Foreign.	Foreign.	Unknown.
10.14		. 44.59	28.44	17.11	90.14	5.55	2.86	1.40	9.81	0.02
65.09 15.02 5.37 85.48  97.15 2.44 0.41 100.00  62.55 25.10 7.97 95.62  85.10 9.72 3.41 98.23  64.45 22.45 6.24 93.14  88.09 7.87 14.93 5.52  97.92 7.84 97.94  87.99 7.87 14.93 5.52  97.92 7.84 97.94  78.77 14.93 5.52 99.22  78.95 13.45 4.97 97.90  78.95 11.74 4.06 97.06  88.45 11.74 4.06 97.06  88.20 11.74 4.06 97.06  88.20 11.74 4.06 97.06  98.20 11.74 4.06 97.06  98.20 11.74 4.06 97.06  98.20 11.74 4.06 97.06  98.20 11.74 4.06 97.06  98.20 11.74 4.06 97.06  98.20 11.74 4.06 97.06  98.20 11.74 4.06 97.55  98.20 11.74 4.06 97.55  98.20 11.74 4.06 97.55  98.20 11.74 4.06 97.55  98.20 11.74 4.06 97.55  98.20 11.74 16.72 7.90 98.73  98.20 11.74 7.22 7.20  98.72 14.60	lel	. 66.18	10.14	4.51	80.83	14.57	2.81	1.62	19.00	0.17
97.15 2.44 0.41 100.00 (2.55 25.10 7.97 95.62 85.10 9.72 3.41 98.23 85.10 9.72 3.41 98.23 85.10 9.72 3.41 98.23 85.09 8.21 1.64 97.94 97.99 7.87 14.93 18.45 99.22 75.30 15.85 5.52 96.78 75.30 15.85 5.65 96.78 75.30 15.73 14.61 96.91 76.85 96.57 15.73 14.61 96.91 97.90 97.90 97.90 97.10 10.42 1.99 99.51 98.85 98.65 98.65 97.00 98.73 77.76 18.36 20.70 7.38 96.89 97.55 97.70 18.36 20.70 7.38 96.89 97.55 97.70 18.36 20.70 7.38 96.89 97.55 97.70 98.70 98.70 98.70 98.70 98.70 98.70 98.70 98.70 98.70 98.70 98.70 97.70		. 65.09	15.02	5.37	85.48	8.19	4.29	1.99	14.47	0.07
85.10 7.97 95.62 85.10 9.72 3.41 98.23 85.10 9.72 3.41 98.23 88.09 8.21 1.64 97.94 87.99 7.85 2.08 97.92 78.77 14.93 5.52 99.22 78.77 14.93 5.62 99.22 78.70 15.86 5.62 96.78 78.95 13.45 4.97 97.90 78.50 10.42 1.99 99.51 88.45 11.74 4.06 97.06 88.65 20.70 7.85 96.89 mties 69.11 16.72 7.90 98.73 77.76 18.86 2.87 98.99 mties 69.11 16.72 7.90 98.73 78.10 1.672 7.91 98.73 77.76 18.86 2.87 98.99 mties 69.11 16.72 7.90 98.73 78.10 1.04 7.22 7.16	•	. 97.15	2.44	0.41	100.00	0.00	0.00	0.00	0.00	0.00
85.10 9.72 3.41 98.23 (6.45 22.45 6.24 93.14 98.23 (6.45 8.09 8.245 6.24 93.14 98.23 (6.24 93.14 98.29 7.82 1.64 97.94 97.94 97.95 78.77 14.93 5.52 99.22 78.77 14.83 6.57 15.73 15.		62.55	25.10	76.7	95.62	2.59	1.79	0.00	4.38	0.00
88.09 8.21 88.09 8.21 1.64 98.14 88.09 8.21 1.64 97.94 97.94 97.95		. 85.10	9.72	3.41	98.23	1.01	0.38	0.38	1.77	0.00
88.09 8.21 1.64 97.94 87.39 7.85 7.85 7.85 7.85 7.85 7.85 7.85 7.85		64.45	22.45	6.24	93.14	4.99	1.04	0.62	6.65	0.21
87.99 7.85 2.08 97.92 78.77 14.93 5.52 99.22 78.95 13.45 96.22 78.95 13.45 97.92 78.95 13.45 4.97 97.37 79.34 14.97 3.59 97.30 79.85 16.59 17.65 20.04 94.28 68.50 17.65 2.85 98.85 87.10 10.42 1.99 99.51 87.11 25.28 15.16 97.55 77.76 18.36 2.87 98.99 mties 69.11 16.72 7.90 93.73 78.90 14.04 7.22 7.16		60.88	8.21	1.64	97.94	1.44	0.21	0.41	2.06	0.00
7.8.77 14.93 5.52 99.22 7 15.30 15.86 5.62 99.22 7 13.45 99.22 7 15.30 15.86 5.62 96.78 75.30 15.86 5.62 96.78 75.30 15.87 14.97 97.37 79.34 14.97 3.59 97.90 17.65 20.04 94.28 efs. 56.59 17.65 20.04 94.28 efs. 56.59 17.65 20.04 99.51 88.45 11.75 2.85 98.85 98.85 11.74 4.06 97.06 87.06 87.01 17.75 18.36 2.87 98.99 17.776 18.36 2.87 98.99 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.30 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 7.33 96.89 17.776 18.36 20.70 98.73 17.70 98.73 17.70 17.70 17.70 17.70 17.70 17.70 17.70 17.70		87.99	7.85	2.08	97.92	0.69	1.04	0.35	2.08	0.00
F0.55 29.27 13.45 93.27 75.30 15.86 5.62 96.78 78.95 18.45 4.97 97.37 78.34 14.97 97.37 78.34 14.97 97.37 78.34 14.97 97.37 78.34 14.97 97.37 78.34 14.97 97.37 14.61 96.91 96.91 17.65 20.04 94.28 87.10 10.42 11.99 99.51 87.45 11.74 4.06 97.06 87.25 87.11 25.28 15.16 97.55 17.77 18.36 2.87 98.89 98.85 98.86 97.06 97.06 97.06 97.06 97.06 97.06 97.07 17.76 18.36 2.87 98.89 98.99 98.89 98.89 98.89 98.89 98.89 98.89 98.89 98.89 98.99 98.89 98.89 98.99 98.89 98.99 98.99 98.99 98.99 98.99 98.99 98.99 98.99		. 78.77	14.93	5.52	99.22	0.31	0.23	0.23	0.77	0.00
75.30 15.86 562 96.78 78.95 13.45 4.97 97.37 78.95 13.45 4.97 97.37 78.85 13.45 4.97 97.37 78.85 15.73 14.61 96.91 rge's 56.59 17.65 20.04 94.28 e's 87.10 10.42 1.99 99.51 88.45 11.55 20.04 94.28 88.45 11.74 4.06 97.66 87.26 11.74 4.06 97.55 88.86 20.70 7.33 96.89 unties 69.11 16.72 7.90 93.73 rahand 61.40 15.50 7.51		. 50.55	29.27	13.45	93.27	5.27	1.09	0.36	6.72	0.00
78.95 13.45 4.97 97.37 78.95 13.45 6.97 97.37 79.34 14.97 3.59 97.90 97.90 97.90 15.73 14.61 96.91 15.73 14.61 96.91 15.73 14.61 94.28 e <sup>5</sup> S. 87.10 10.42 1.99 99.51 84.45 11.55 20.04 94.28 89.62 6.36 0.85 98.85 89.62 17.76 18.36 2.87 98.99 17.77 18.36 2.87 98.99 18.36 2.87 98.99 18.36 2.97 18.36 2.87 98.99 18.36 2.97 18.36 2.87 98.99 18.36 2.97 18.36 2.97 18.36 2.37 18.37 18.36 2.37 18.36 2.37 18.36 2.37 18.37		. 75.30	15.86	5.62	96.78	1.00	1.61	0.60	3.21	0.00
rege's. 79.34 14.97 3.59 97.90 17.65 66.57 15.73 14.61 96.91 97.90 17.65 20.04 94.28 17.65 20.04 94.28 17.65 20.04 94.28 17.65 20.04 94.28 17.65 20.04 94.28 17.65 20.04 99.51 17.65 20.04 99.51 17.6 18.86 2.85 96.85 96.85 11.74 4.06 97.06 17.75 18.36 2.87 98.99 17.75 18.36 2.87 98.99 17.75 18.36 20.70 17.35 96.89 17.75 18.36 20.70 17.35 96.89 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.35 96.89 17.75 17.75 18.36 20.70 17.75 18.3		. 78.95	13.45	4.97	97.37	1.75	0.88	0.00	2.63	0.00
rge's 56.59 17.65 20.04 94.28 e's 56.59 17.65 20.04 94.28 e's 17.10 10.42 1.99 99.51 e's 84.45 11.55 2.85 98.85 89.62 6.36 0.85 96.83 81.26 11.74 4.06 97.06 87.11 25.28 15.16 97.55 88.62 20.70 7.33 96.89 unties 69.11 16.72 7.90 93.73 fryland 61.40 15.50 7.61		. 79.34	14.97	3.59	97.90	0.60	06.0	0.30	1.80	0.30
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1	. 66.57	15.73	14.61	96.91	1.40	0.42	0.98	2.80	0.28
e's. 87.10 10.42 1.99 99.51 88.45 11.55 2.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.85 98.86 97.06 97.06 97.06 97.06 97.06 97.06 98.86 20.70 7.33 96.89 98.80 98.8	rge's	. 56.59	17.65	50.04	94.28	2.60	1.77	1.35	5.72	0.00
89.45 11.55 2.85 98.85 88.62 6.36 0.85 96.83 88.62 6.36 0.85 96.83 96.83 87.06 11.74 4.06 97.06 97.06 97.05 97.06 97.06 97.06 97.06 98.86 20.70 7.83 96.89 96.89 97.05 98.15 96.89 97.05 98.15 96.89 97.05 98.15 96.89 97.05 98.15 97.05 9	e's	87.10	10.42	1.99	99.51	0.25	0.25	0.00	0.50	0.00
89.62 6.36 0.85 96.83 88.83 11.74 4.06 97.06 97.06 97.06 97.07 11.74 15.16 97.55 98.99 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 96.89 97.55 97.		84.45	11.55	2.85	98.85	0.00	0.43	0.29	0.72	0.43
81.26 11.74 4.06 97.06   77.11 25.28 15.16 97.55   77.76 18.36 20.70 7.33 96.89   68.86 20.70 7.33 96.89   68.81 16.72 7.90 93.73   69.91 16.72 7.90 93.73   7.91 61.40 15.60 7.61 61.40   7.61 61.40 15.60 7.61 61.60		. 89.62	6.36	0.85	96.83	2.12	0.42	0.42	2.96	0.21
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	albot	. 81.26	11.74	4.06	97.06	06.0	1.58	0.45	2.93	0.00
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	ashington	. 57.11	25.28	15.16	97.55	1.32	0.99	0.13	2.44	0.00
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	/icomico	. 77.76	18.36	2.87	66.86	0.43	0.57	0.00	1.00	0.00
ies	orcester	. 68.86	20.70	7.33	96.89	2.38	0.55	0.18	3,11	0.00
ies $69.11$ $16.72$ $7.90$ $93.73$ $50.90$ $14.04$ $7.22$ $72.16$ $14.04$ $7.22$ $72.16$ $14.04$ $1.80$ $14.04$ $1.80$ $18.00$			1	1					1	1
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	ies	. 69.11	16.72	7.90	93.73	3.72	1.67	0.80	6.19	0.02
6140 1850 7.61		. 50.90	14.04	7.22	72.16	20.86	4.36	2.59	27.81	0.03
61 AO 18 RO 78 RO				]		1		1		]
00.40 10.1 66.61 04.10	Total Maryland	. 61.40	15.59	7.61	84.60	10.98	2.81	1.56	15.35	0.05

In the preceding table (Table VIII) the nationality of parents of children born in Maryland during 1915 is returned in six columns. In the sub-columns the native and foreign-born parents are considered separately. In the column headed "Neither Maryland," the American-born parents are included, both born in the United States, but neither a native of Maryland. The foreign column is in three sub-divisions, in the first of which are included parents, both of whom are of foreign birth; and second, father foreign; and third, mother foreign. By reference to this column it will be seen that the largest proportion of native parents occurred in Calvert County, 100 per cent. The greatest proportion of foreign parents was in Anne Arundel County, 19 per cent. Baltimore County was next in order with 14.47 per cent. In two counties the proportion of native-born parents exceeded 99 per cent., Frederick and Queen Anne's. In Calvert County 97.15 per cent. of the parents of children born in 1915 were both natives of Maryland, and over 80 per cent. of the parents of children born in Carroll, Charles, Dorchester, Queen Anne's, Somerset, St. Mary's and Talbot counties were both natives of this State. In Rural Maryland at large there were 93.73 per cent. of the parents native of the United States, and 6.19 per cent. foreign-born. The highest percentage of both parents foreign was in Anne Arundel County. 14.57 per cent., and the highest percentage of mothers foreign was in Baltimore County, 1.99 per cent. In Baltimore County 4.29 per cent. of the fathers were foreign-born.

# FECUNDITY.

Under the title of "Fecundity" is included a review of the fecundity rates and the maximum fecundities by counties in the white and colored population. Along with these we place the tables showing maximum and minimum ages of parents, by counties and color, and the average age of parents.

# TABLE IX.

Table of Fecundity for White and Colored Mothers by Age Periods—Rural Maryland—1915.

			Wh	ite.				
Ayes of Mothers.	No. of Mothers.	Total No. of Children Born.	Total No. of Children Living.	Total Children per Mother.	Living Children per Mother.	Total Children per 10,000 of Fe- male Population.	Living Children per 10,000 of Fe- male Population.	Estimated Female Population.
10-15 years	5	5	5	1.00	1.00	1.55	1.55	32,208
15-20 years	1,437	1,749	1,656	1.22	1.15	585.87	554.72	
20-25 years	4,035	8,006	7,278	1.98	1.80	2,998.73	2,726.05	26,698
25-30 years	3,631	11,279	10,072	3.11	2.77	4,701.74	4,198.59	23,989
30-35 years	2,727	12,127	10,509	4.45	3.85	5,491.55	4,758.86	22,083
35-40 years	1,965	12,083	10,290	6.15	5.24	5,800.77	4,939.99	20,830
40-45 years	719	5,592	4,701	7.78	6.54	3,158.79	2,655.48	17,703
45-50 years	103	897	755	8.71	7.33	590.44	496.97	15,192
Age Unknown	43	149	122	3.47	2.84	• • • • • •	•••••	• • • • • • ′
Total	14,665	51,887	45,388	3.54	3.09	2,751.81	2,407.14	188,556
			Colo	red.				
10-15 years	23	25	25	1.09	1.09	28.29	28.29	8,836
15-20 years	796	1.080	971	1.36	1.22	1,433,50	1,288.82	7,534
20-25 years	1,069	2,960	2,411	2.77	2.26	4.832.65	3,936.33	6.125
25-30 years	774	3,612	2,943	4.67	3.80	6,959.54	5,670.52	5,190
30-35 years	603	3,830	3,115	6.35	5.17	9,200.10	7,482.58	4,163
35-40 years	481	4,062	3,155	8.44	6.56	10,012.32	7.776.68	4,057
40-45 years	177	1,732	1,346	9.79	7.60	5,077.69	3,946.06	3,411
45-50 years	22	214	167	9.73	7.59	724.20	565.14	2,955
Age Unknown	49	163	124	3.33	2.53		• • • • • •	
Total	3,994	17,678	12,911	4.18	3.23	4,182.06	3,054.34	42,271
A form mothone	include	and my boar	DONOG TE	TOPO O	70r 50	TOOPE		

A few mothers included whose ages were over 50 years.

TABLE IX-A.

Table of Fecundity for White and Colored Mothers by Age Periods—Baltimore City—1915.

			Wh	ite.			<i>'</i>	
Ages of Mothers.	No. of Mothers.	Total No. of Children Born.	Total No. of Children Living.	Total Children & per Mother.	or Living Children of per Mother.	Total Children per 10,000 of Fe-	Living Children per 10,000 of Fe-	Estimated Female Population.
10-15 years 15-20 years	3 1,042	$\frac{3}{1,261}$	3 1,180	1.00 $1.21$	1.13	1.33 488.19	456.83	22,620 25,830
15-20 years 20-25 years	3,610	6,478	5,818	1.79	1.61	2,421.32	2,174.63	26,754
25-30 years	3,150	8,701	7,645	2.76	2.43	3,687.49	3,239.96	23,596
30-35 years	2,031	8,392	7,134	4.13	3.51	4,137.25	3,517.06	20,284
35-40 years	1,183	6,712	5,538	5.67	4.68	3,537.47	2,918.73	18,974
40-45 years	405	3,032	2,417	7.49	5.97	1,808.42	1,441.61	16,766
45-50 years	30	230	182	7.67	6.07	160.25	126.80	14,353
Age Unknown	7	16	16	2.29	2.29			
	'							
Total	11,461	34,825	29,933	3.04	2.61	2,058.49	1,769.33	169,177
			Colo	red.				
10-15 years	12	12	12	1.00	1.00	34.72	34.72	3,456
15-20 years	525	702	623	1.34	1.19	1,524.43	1,352.88	4,605
20-25 years	713	1,644	1,318	2.31	$^{-}1.85$	2,757.00	2,210.30	5,963
25-30 years	435	1,697	1,283	3.90	2.95	2,918.31	2,206.36	5,815
30-35 years	298	1,560	1,065	5.23	3.57	3,751.80	2,561.33	4,158
35-40 years	193	1,232	869	6.38	4.50	2,972.26	2,096.50	4,145
40-45 years	60	483	314	8.05	5.23	1,511.26	982.48	3,196
45-50 years	6	48	28	8.00	4.67	181.68	105.98	2,642
Age Unknown	5	17	9	3.40	1.80		•••••	
Total	2,247	7,395	5,521	3.29	2.46	2,176.28	1,624.78	33,980
1 white mother	r at the	age of	58 years	<b>5.</b>				

In order to ascertain the fecundity rates of the child-bearing portion of the population, Tables IX and 9-A have been devised. These tables are valuable for comparative purposes demonstrating particularly the difference in rates in a wholly urban population and the population of Maryland outside of Baltimore City, which is to a large extent a rural population. Of equal interest is a comparison of the difference of rates in the white and colored populations.

Chart II shows the fecundity rates for Rural Maryland and Baltimore City for total children born and total children living per 10,000 of female population, white and colored, in 1915.




CHART 2.

The succeeding table (Table X) gives figures of maximum fecundity in Rural Maryland during 1915.

Only children in excess of the tenth during 1915 are considered. Of these, there were 349 white and 262 colored, a total of 611. The percentage of all births being: white, 1.87 percent.; colored, 1.40 per cent.; total, 3.27 per cent.

Twin births occurred in the white population 149 times and in the colored population 50 times. There were also two triple births in the colored population.

# TABLE X.

Table of Maximum Fecundity—Number of Child in Excess of Tenth Born in Maryland, Exclusive of Baltimore City, 1915.

				-			_							
. County.		11	12	13	14	15	16	17	18	19	20	Total.	Twins.	Trip- lets.
Allegany	W. C.	18	7	5 2	$\begin{vmatrix} 2 \\ \cdot \\ 1 \end{vmatrix}$	3						36  16	22	
Anne Arundel }	W. C.	11	5	6	4	3			1		1	31	2	
Baltimore	W.   C.	23	$\begin{vmatrix} 14 \\ 5 \end{vmatrix}$	15   1	$\begin{vmatrix} 6 \\ 2 \end{vmatrix}$	3	$\begin{bmatrix} 2 \\ \cdot \cdot \end{bmatrix}$	i	1	::		64 13	33 3	
Calvert	W. C.	1 5	1	$\frac{\cdot \cdot}{2}$	2							1 10	$\frac{2}{3}$	
Caroline	W. C.	5 2	3 3	1 5	1 1	1 1		• •			• •	11 12	5 3	
Carroll	W.	10 2	4	$\frac{2}{1}$	• •	1	•			 		17 4	4	
Cecil	W. C.	$\begin{vmatrix} 4\\2 \end{vmatrix}$	1 1	1 1		• •	• •	• •			•	$\frac{1}{6}$	3 0	
Charles	W. C.	3 5	3 6	3								9 15	$\frac{2}{1}$	
Dorchester }	W. C.	3 14		4	1	1		2			 1	7 29	4	
Frederick	W. C.	10	16 1	6 3	$\frac{1}{2}$	2						35 10	5 0	
Garrett	W. C.	3	4	3	1	1		1			••	13	5	
Harford	W.		4	1 2	3	1	1					9	8	• •)• •
Howard	W.	$\frac{1}{2}$	$\frac{2}{7}$	$\frac{1}{2}$		1						8	$\frac{1}{2}$	
Kent	W.	4	$\begin{vmatrix} 3 \\ 4 \end{vmatrix}$	2 4	$egin{array}{c} 1 \ 2 \end{array}$	1						10 11	$\frac{1}{2}$	
Montgomery }	W. C.	5 3	4	$\frac{1}{2}$	3		1		1			11 11	2 5	
Prince George's	W. C.	9	3 5	4 6	1	1	$\frac{1}{2}$	2			1	18 27	9	
Queen Anne's	W. C.	$\frac{2}{4}$	$\frac{1}{2}$	1	1	••	••	•-		$\cdots$		4 8	3	1
Somerset	W. C.	3	$\frac{1}{2}$	$\frac{1}{2}$	1 2	1	1					7 9	2	
St. Mary's	W. C.	4	1 3	$egin{array}{c} 2 \ 1 \ \end{array}$	$\frac{1}{2}$	1				1		9	3	*1
Talbot	w. C.		$\cdot \cdot \cdot \cdot 2$	2	••	••	••	$\cdot \cdot  $	••	•••		7	1	
Washington	W. C.	17	10	7	4	3		2		$\cdot \cdot  $		43	15 1	
Wicomico	W. C.	4 5	1 5	3	1	1	1					11 12	10	• • • •
Worcester	- W. C.	4	5				1	••	$\cdot \cdot  $	• •	$ \cdot $	4 16	3 2	
White	· · · · ·	$\begin{vmatrix} 140 \\ 90 \end{vmatrix}$	88 65			$egin{array}{c c} 20 \ 12 \ \end{array}$	5	3	$\begin{vmatrix} \cdot \cdot \\ 2 \\ 2 \end{vmatrix}$	· ·		- 1	$\begin{array}{c} 2\\149\\50\end{array}$	2
Total		230	[	—-j	-/-		— - 11	9	4	1	-1	611		$\frac{2}{2}$
*0										!		- '		

<sup>\*</sup>One child stillborn, two living.

In Table XI the maximum age of fathers and mothers of children born in Rural Maryland during 1915 are shown for the white and colored races by counties. This table considers A, maximum age of father; B, maximum age of mother; C, minimum age of father; D, minimum age of mother.

Maximum age of father, 81 years (mother 40, black). Maximum age of mother, 58 years (father 59, white). Minimum age of father, 15 years (mother 18, black). Minimum age of mother, 12 years (father 20, black).

TABLE XI.

MAXIMUM AND MINIMUM AGES OF FATHERS AND MOTHERS OF CHILDREN BORN IN MARYLAND, 1915.

ALLEGAN	COUNT	Y.		
Age of father		W. 49 47 B	W. 18 17 C	W. Unk. 14 D
Anne Arun	DEL COU	NTY.		
Age of fatherAge of mother		B. 24 52 B	W. 17 16 C	B. Unk. 14 D
BALTIMOR	E COUNT	Υ.		
Age of fatherAge of mother	W. 64 30 A	W. 54 48 B	B. 18 15 C	B. Unk. 13 D
CALVERT	COUNTY	•		
Age of father		W. 43 43 B	B. 17 13 C D	
Caroline	COUNTY	τ.		
Age of father		W. 50 45 B	B. 17 15 U D	
CARROLL	COUNTY			
Age of fatherAge of mother		W. 36 49 B	W. 16 16 C	W. 26 14 D

STATE BOARD OF HEALTH.									
CECIL CO	OUNTY.								
	W.	В.	w.	B.					
Age of father	61	45	17	35					
Age of mother	41	50	17	15					
	A	В	C	D					
CHARLES	County	ī <b>.</b>							
	В.	w.	В.	В.					
Age of father	70	51	18	24					
Age of mother	39	46	15	14					
	A	В	Ö	D					
Dorchester County.									
•	W.	В.	В.	В.					
Age of father	72	$\tilde{32}$	16	$\frac{26}{26}$					
Age of mother	33	48	16	13					
6	A	В	C	D					
Frederick County.									
	w.	w.	W.	W.					
Age of father	75	53	16	21					
Age of mother	35	48	18	14					
· ·	A	В	C .	D					
Garrett (	County								
Gilling I			***						
	W.	W.	W.	W.					
Age of father	75	$\frac{52}{12}$	17	39					
Age of mother	26 A	46 B	17 C	14 D					
	11.	ъ	C,	D					
Harford	COUNTY	7.							
•	w.	w.	W.	В.					
Age of father	55	48	17	20					
Age of mother	39	$\overset{10}{45}$	20	15					
	A	B	C	$\widetilde{\mathbf{D}}$					
Howard County.									
	В.	В.	В.	В.					
Age of father	71	63	17	$\frac{2}{45}$					
Age of mother	43	46	15	15					
	A	В	O	D					
KENT C	OUNTY.								
	В.	W.	В.						
Age of father	72	56	17						
Age of mother	14	49	16						
3	A	В	C						
	D								
Montgomer	Y Cour	NTY.							
	В.	В.	В.	В.					
Age of father	70	$\overline{52}$	15	17					
Age of mother	42	47	18	13					
	A	В	C	D					

50 REPORT (	)F TH	E		
Prince Georg	e's Co	UNTY.		
	В.	W.	В.	В.
Age of father	81	60	16	$\frac{2}{21}$
Age of mother	40	50	$\frac{10}{27}$	14
Age of mother	A	B	Ċ	D
	Δ.,	ь		D
QUEEN ANNE	's Cou	NTY.		
	В.	W.	W.	В.
Age of father	63	48	18	23
Age of mother	32	42	18	13
	A	′ B	$\mathbf{C}$	D
Somerset	Count	Y.		
	w.	В.	В.	· B.
Ago of father	61	50	18	Unk.
Age of father		48	14	12
Age of mother	29			D 12
	A	В	С	D
St. Mary's	Coun	TY.		
	W.	W.	В.	В.
Age of father	65	49	15	25
Age of mother	33	46	18	15
,	A	В	$\tilde{\mathbf{c}}$	D
Talbot (	County	ī.		
	w.	W.	В.	
Age of father	69	46	15	
Age of mother	42	46	15	
age of mother	A	B	Č	
	Α	Б	$\mathbf{\tilde{D}}$	
Washington	n Cou	NTY.		
	w.	w.	- w.	w.
Age of father	72	34	17	22
Age of mother	32	48	18	15
arge of mother	Ā	В	Ö	D
,	21	Б	O	D
Wicomico	Count	ry.		
	В.	В.	В.	В.
Age of father	74	45	17	19
Age of mother	30	53	16	14
	A	В	G	D
Worcester	Coun	TY.	•	
	W.	В.	В	. В.
Age of father	65	50	17	23
Age of mother	42	46	15	13
	A	В	- O	D
Baltimor	E CITY	r <b>.</b>		
	W.		В.	D
Age of father		W.		B.
	73 45	59 50	15 15	20
Age of mother	45	58 P	15	12
	A	В	Q	D

### AVERAGE AGES OF PARENTS.

The succeeding table (Table XII) gives the ages of fathers and mothers of children born in Rural Maryland and Baltimore City for the year 1915. The table demonstrates the superior fecundity of females in early life and the superior fecundity of males at the higher age periods. This table shows an earlier average age for parents in Baltimore City than in Rural Maryland.

TABLE XII.

Ages of Parents of Children Born in Maryland During 1915.

		Rural	Marylan	d.		Baltlimore City.			
Age Period.	Number of Fathers.	Per Cent.	Number of Mothers.	Per Cent.	Number of Fathers.	Per Cent. Number of	Mothers.	Per Cent.	
10-15			28	0.15			15	0.11	
15-20	309 [	1.66	2,233	11.97	197	1.44	1,567	11.43	
20-25	3,451	18.50	5,104	27.35	2,824	20.60	4,323	31.54	
25-30	4,445	23.82	4,405	23.61	3,994	29.14	3,585	26.15	
30-35	3.691	19.78	3,330	17.85	2,805	20.46	2,329	16.99	
35-40	3.060	16.40	2,446	13.11	1.972	14.39	1.376	10.04	
40-45	1.976	10.59	896	4.80	1,080	7.88	465	3.39	
45-50	919	4.93			442	3.22			
50 and over	543	2.91			173	1.26			
45 and over			125	0.67			36	0.26	
Age unknown.	265	1.42	92	0.49	221	1.61	12	0.09	
Total	18,659		18,659		13,708		13,708		

Two mothers at the age of 12; eight mothers at the age of 13; thirty-three mothers at the age of 14.

Table XIII gives the average ages of parents of children born in Rural Maryland for the years 1906 to 1915, inclusive. Except for slight variations in any single year, the average age of both fathers and mothers remains the same. The average difference between the ages of parents shows no marked variation.

TABLE XIII.

AVERAGE AGE OF PARENTS OF CHILDREN BORN IN RUBAL MARYLAND, 1906-1915.

	Average Age of	•	Average Age of	Average Difference Between Father's
Year.	Father.		Mother.	and Mother's Ages.
1906	32.44		27.73	4.71
1907	33.06		28.42	- 4.64
1908	~~ ~~		28.31	4.62
1909	32.97		28.36	4.61
1910	32.83		28.32	4.51
1911			28.35	4.50
1912			28.23	4.53
1913	. 32.82		28.23	4.59
1914			28.00	4.53
1915	00.10		27.99	4.49

# Birth Registration.

For the purpose of ascertaining the extent of the practice of midwives as shown by registration, two tables were made. Table XIV gives the actual number of births attended by physicians, midwives or others, i. e., no physician or midwife in attendance. Of a total of 26,126 white births, 20,179 were attended by physicians, 5,802 by midwives and 145 did not have the attendance of either a physician or midwife. Of a total of 6,241 colored births, 3,243 received the attention of a physician, 2,927 were attended by midwives and 71 had neither the attention of a physician or midwife. The total number of births attended by physicians was 23,422, the total number attended by midwives was 8,729, and the total number without the attention of midwife or physician was 216.

TABLE XIV.

Proportion of Births Attended by Physicians and Midwives, White and Colored, Maryland, 1915.

		White.				Colored.			
Counties.	Physician.	Midwife.	Other Person.	Total.	Physician.	Midwife.	Other Person.	Total.	( \
Allegany	1,612	95	27	1,734	36	11	$\frac{2}{3}$	49	1,783
Anne Arundel	525	236	0	761	131-	279	3	413	1,174
Baltimore	1,938	700	9	2,647	152	64	7	223	2,870
Calvert	63	32	0	95	13	137	1	151	246
Caroline	266	93	5	364	20	114	4	138	502
Carroll	727	11	6	744	38	10	0	48	792
Cecil	425	3	2	430	36	14	1	51	481
Charles	106	89	6	201	22	256	8	286	487
Dorchester	392	115	10	517	108	238	3	349	866
Frederick	1,082	41	14	1,137	116	29	4	149	1,286
Garrett	489	28	33	550	0	0 8	0	79	$\frac{550}{498}$
Harford	409	8	2	419	68	_	3 1	107	342
Howard	214	20	1	235	59 48	47 80	3	131	$\frac{342}{334}$
Kent	176	$\frac{27}{14}$	0	$\frac{203}{487}$	128	94	3	$\frac{151}{225}$	712
Montgomery	470	66"	$\frac{3}{2}$	618	108	233	4	345	963
Prince George's.	$\frac{550}{227}$	31	ő	$\frac{018}{258}$	36	108	1	145	403
Queen Anne's Somerset	387	$\frac{51}{28}$	0	415	118	165	3	286	701
Somerset St. Mary's	172	83	ő	255	50	165	$\frac{3}{2}$	217	472
Talbot	257	35	1	293	52	97	ĩ	150	443
Washington	1,392	69	13	1,474	29	6	2	37	1.511
Wicomico	415	71	3	489	87	115	$\bar{6}$	208	697
Worcester	292	42	5	339	49	153	5	207	546
	12,586	1,937	142	14,665	1,504	2,423	67	3,994	18,659
Baltimore City	7,593	3.865	3	11,461	1,739	504	4	2,247	13,708
Maryland	20,179	-5,802	145	26,126	3,243	2,927	71	6,241	32,367

Table XV gives the percentage of births, white and colored, attended by physicians and midwives, respectively. In all counties the majority of white mothers had the attendance of a physician. In Charles County, 44.28 per cent. of white births were attended by midwives, and in a number of the other counties, namely, Anne Arundel, Baltimore, Calvert, Caroline and St. Mary's, over 25 per cent. of white mothers were attended by midwives. Counties in which the largest proportion of white mothers had neither the services of physicians or midwives were Garrett, 6 per cent. of births reported, and Charles, 2.99

per cent. of births reported. The large practice of midwives is seen among the colored women. In thirteen counties over 50 per cent. of colored births were attended by midwives. The county in which the largest percentage of colored mothers were attended by midwives is Calvert County, 90.73 per cent.; Charles county second, with 89.51 per cent. The counties in which the largest percentage of colored mothers had neither the attention of physician or midwife are Washington County, 5.41 per cent., and Allegany County, 4.08 per cent.

TABLE XV.

Percentage of Births Attended by Physicians and Midwives, White and Colored, Maryland, 1915.

	Percentage of White Births.			Percen	tage of Colored Births.		
Countics.	Physicians.	Midwives.	Others	Physicians.	Midwives.	Others.	
Allegany	92.96	5.48	1.56	73.47	22.45	4.08	
Anne Arundel	68.99	31.01	0.00	31.72	67.55	0.73	
Baltimore	73.21	26.45	0.34	68.16	28.70	. 3.14	
Calvert	66.32	33.68	0.00	8.61	90.73	0.66	
Caroline	73.08	25.55	1.37	14.49	82.61	2.90	
Carroll	97.72	1.48	0.81	79.17	20.84	0.00	
Cecil	98.84	0.70	0.47	70.59	27.45	- 1.96	
Charles	52.74	44.28	2.99	7.69	89.51	2.80	
Dorchester	75.82	22.24	1.93	30.95	68.19	0.86	
Frederick	95.16	3.61	1.23	77.85	19.46	2.68	
Garrett	88.91	5.09	6.00	0.00	0.00	0.00	
Harford	97.61	1.91	0.48	86.08	10.13	3.80	
Howard	91.06	8.51	0.43	55.14	43.93	0.93	
Kent	86.70	13.30	0.00	36.64	61.07	2.29	
Montgomery	96.51	2.87	0.62	56.89	41.78	1.33	
Prince George's	89.00	10.68	0.32	31.30	67.54	1.16	
Queen Anne's	87.98	12.02	0.00	24.83	74.48	0.69	
Somerset	93.25	6.75	0.00	41.26	57.69	1.05	
St. Mary's	67.45	32.55	0.00	23.04	76.04	0.92	
Talbot	87.71	11.95	0.34	34.67	64.67	0.67	
Washington	94.44	4.68	0.88	78.38	16.22	-5.41	
Wicomico	84.87	14.52	0.61	41.83	55.29	2.88	
Worcester	86.14	12.39	1.47	23.67	73.91	2.42	
	·						
Total Counties		13.21	0.97	37.66	60.67	1.68	
Baltimore City	66.25	33.72	0.03	77.39	22.43	0.18	
Maryland	77.24	22.21 ·	0.56	51.96	46.90	1.14	

Table XVI gives the births, estimated population and births per 1,000 in cities of Maryland with population of 2,000 and over. Upon investigation we found it was the practice of physicians and midwives to enter the nearest town as the address instead of giving the exact location, and in this manner a number of births which did not occur within their limits were credited to some cities. A system has been devised which will correct this error in tables beginning with the year 1917.

#### TABLE XVI.

BIRTHS BY SEX AND COLOR, ESTIMATED POPULATION AND BIRTH RATES PER 1,000 OF CITIES AND TOWNS IN MARYLAND WITH POPULATION OF 2,000 AND OVER, 1915.

						ed tion.	Per
Towns.	le.	Female.	White.	Colored.	Total.	Estimated Population.	Births 1,000.
10000.	Male.	Fe	<b>A</b>	Co -	To	$E_{P}$	Bi
Baltimore City7	,028	6,680	11,461	2,247	13,708	584,608	23.45
Cumberland	384	344	689	39	. 728	25,564	28.48
Hagerstown	322	325	631	16	647	25,233	25.64
Frederick	115	130	203	42	245	10,999	22.27
Annapolis	129	101	152	78	230	8,751	26.28
Salisbury	105	106	169	42	211	<b>7,963</b>	26.50
Cambridge	135	110	149	96	245	6,755	36.27
Frostburg	82	113	187	8	195	6,426	30.35
Havre de Grace	42	38	66	14	80	4,628	17.29
Brunswick	58	53	107	4	111	4,380	25.34
Crisfield	61	57	85	33	118	. 3,628	32.52
Westminster	` 35	47	80	2	82	3,346	24.51
Easton	54	42	66	30	96	3,088	31.09
Westernport	53	47	98	2	100	3,073	32.54
Laurel	29	20	43	6	49	2,592	18.90
Chestertown	12	26	22	16	38	2,591	14.67
Pocomoke City	43	42	54	31	85	2,498	34.03
Elkton	33	28	57	4	61	2,283	$24.8\hat{2}$
Hyattsville	28	25	45	8	53	2,283	23.22
		<u></u>					
TotalS	3,748	8,334	14,364	2,718	17,082	710,864	24.03

One case sex unknown.

# Deaths in Maryland.

The total number of deaths recorded in Maryland during 1915 was 21,350; of these, 12,023 occurred in Rural Maryland and 9,327 in Baltimore City.

The total number of white deaths was 15,767; of colored deaths, 5,583.

The total number of male deaths was 11,221; of female deaths, 10,129.

Tables III and 3-A give the births, deaths and increase, and the birth rates, death rates and rate of increase (per 1,000) for the white and colored population of Maryland.

Table IV gives the births, birth rates, deaths, death rates and excess of births over deaths per 1,000 of population for the male, female, white and colored, and total population of Rural Maryland, Baltimore City and the State of Maryland.

#### TABLE XVII.

DEATHS BY SEX AND COLOR, ESTIMATED POPULATION AND DEATH RATES PER 1,000 OF CITIES AND TOWNS IN MARYLAND WITH POPULATION OF 2,000 AND OVER, 1915.

Towns.	Male.	Female.	White.	Colored.	Total.	Estimated Population.	Deaths Per 1,000.
Baltimore City	4,835	4,492	7,149	2,178	9,327	584,608	15.95
Cumberland	202	193	364	31	395	25,564	15.45
Hagerstown	184	187	333	38	371	25,233	14.70
Frederick	84	. 96	145	35	180	10,999	16.37
Annapolis	82	76	77	81	158	8,751	18.06
Salisbury	89	73	107	55	162	7,963	20.34
Cambridge	- 93	79	87	85	172	6,755	25.46
Frostburg	58	· 42	93	7	100	6,426	15.56
Havre de Grace	41	$^{26}$	51	16	67	4,628	14.48
Brunswick	19	20	34	5	39	4,380	8.90
Crisfield	32	45	53	$^{24}$	77	3,628	21.22
Westminster	- 36	24	50	10	60	3,346	17.93
Easton	61	34	52	43	.95	3,088	30.76
Westernport	21	19	-36	4	40	3,073	13.02
Laurel	20	29	45	4	49	2,592	18.90
Chestertown	30	$^{27}$	23	34	57	2,591	22.00
Pocomoke City	32	34	40	26	66	2,498	26.42
Elkton	44	38	65	17	82	2,458	33.36
Hyattsville	11	11	19	3	22	2,283	9.64
Total	5,974	5,5 <del>4</del> 5	8,823	2,696	11,519	710,864	16.20

Table XVII gives deaths male, female, white, colored, estimated population and deaths per 1,000 for cities of Rural Maryland with a population of 2,000 or over. The rates for a number of the cities as given in the table are high. There are two reasons for this; first, that a number of the cities support hospitals and all the deaths which occur in the hospitals add to the death rate for the cities; second, the improper addresses on certificates causes a number of deaths which did not occur within the corporate limits of the cities to be included in the total number of deaths. We have taken means to eliminate this error in all tables beginning with 1917.

## TABLE XVIII.

Deaths for the Year 1915 by Ages, Showing Percentages of Total Mortality in the Several Age Periods of Life in Rural Maryland, Baltimore City and the State of Maryland.

	Rural Maryland.		Baltimo	re City.	Maryland.	
Age Periods.	Deaths.	Per Ct.	Deaths.	Per Ct.	Deaths.	Per Ct.
0- 1 year	2,238	18.61	1,624	17.41	3,862	18.09
1- 5 years		5.70	537	5.76	1,222	5.72
5-10 years	213	1.77	158	1.69	371	1.74
10-15 years	199	1.66	109	1.17	308	1.44
15-20 years	323	2.69	221	2.37	544	2.55
20-25 years	428	3.56	309	3.31	737	3.45
25-30 years	425	3.53	370	3.97	795	3.72
30-35 years	446	3.71	371	3.98	817	3.83
35-40 years		3.93	458	4.91	930	4.36
40-45 years	456	3.79	486	5.21	942	4.41
45-50 years	481	4.00	551	5.91	1,032	4.83
50-55 years		4.72	643	6.89	1,210	5.67
55-60 years	654	5.44	663	7.11	1,317	6.17
60-65 years	763	6.35	672	7.20	1,435	6.72
65-70 years	855	7.11	607	6.51	1,462	6.85
70–75 years	933	7.76	564	6.05	. 1,497	7.01
75-80 years	821	6.83	482	5.17	1,303	6.10
80 years and over.		8.18	502	5.38	1,485	6.96
Age unknown	. 81	0.67	0	0.00	81	0.38
Total	.12,023		9,327		21,350	




Chart 3—State of Maryland—Deaths by Ages, 1915.




CHART 4—Deaths by Ages.

The preceding table (Table XVIII) gives the distribution of the deaths in Rural Maryland, Baltimore City and the State of Maryland; according to age, giving the number of deaths at each age period and the percentages of total mortality. Chart 3 and Chart 4 show graphically the percentage of deaths at the several age periods; the charts indicate the excessive percentage of deaths in infancy (under 2 and under 5 years).

The table shows that the infantile mortality is slightly higher in Rural Maryland than in Baltimore City, as 18.61 per cent. of the deaths in Rural Maryland occurred under the age of 1 year and 17.41 per cent. in Baltimore City. The figures under the age of 5 for Baltimore City are 23.17 and for Rural Maryland 24.31 per cent. This difference is shown graphically in the accompanying charts.

Tables 19-A, 19-B, 19-C, give the estimated population, death rates per 1,000 at age periods and the mortality per 1,000 for age periods for the white, colored and total populations of Maryland. The mortality per 1,000 at age periods in the total population is lowest in the third quinquennium, after which it slowly rises until the period between 60 and 65 years, when it exceeds the mortality at the period of 0-5/years.

The uniformly higher death rate by ages in the colored population is well demonstrated up to the age period of 55-60, when the colored mortality almost doubles the white.

# TABLE XIX-A.

Population and Deaths Per Thousand at the Several Age Periods—Maryland, 1915.

• = "	Per	Esti- mated Popu-		of Those Living at	Mortality Per 1,000 for Age
Age.	Cent.	lation.	Deaths.	$the\ Age.$	Periods.
Under 5 years	10.61	143,457	5.084	35.44	177.20
5 to 10 years		139,148	371	2.67	13.35
10 to 15 years		135,032	308	2.28	11.40
15 to 20 years		133.517	544	4.07	20.35
20 to 25 years		128,457	737	5.74	28.70
25 to 30 years		114,752	795	6.93	34.65
30 to 35 years		100,049	817	8.17	40.85
35 to 40 years		96,262	930	9.66	48.30
40 to 45 years		82,091	942	11.47	57.35
45 to 50 years	5,24	70,867	1,032	14.56	72.80
50 to 55 years		61,725	1.210	19.60	98.00
55 to 60 years		45,106	1,317	29.20	146.00
60 to 65 years		36,514	1,435	39.30	196.50
65 to 70 years		27,752	1,462	52.68	263.40
70 to 75 years		18,435	1,497	81.20	406.00
75 to 80 years		10,280	1,303	126.75	633.75
80 years and over.		7,115	1,485	208.71	
Unknown		1,381	81	58.65	
Total		1,351,940	21,350	15.79	

# TABLE XIX-B.

ESTIMATED WHITE POPULATION AND DEATHS PER THOUSAND AT THE AGE PERIODS FOR THE YEAR 1915—MARYLAND'S ESTIMATED WHITE POPULATION, 1,120,769.

	Per	Esti- mated Popu-		Mortality Per 1,000 of Those Living at	Mortality Per 1,000 for Age
Age.	Cent.	lation,	Deaths,	the Age.	Periods.
Under 5 years		117.681	3,490	29.66	148.30
5 to 10 years		113.534	256	2.25	11.25
10 to 15 years	9.87	110,620	186	1.68	8.40
15 to 20 years	9.84	110,284	324	2.94	14.70
20 to 25 years	9.37	105,016	446	4.25	.21.25
25 to 30 years	8.37	93,808	515	5.49	27.45
30 to 35 years	7.45	83,497	564	6.75	33.75
35 to 40 years	7.11	79,687	632	7.93	39.65
40 to 45 years	6.12	68,591	647	9.43	47.15
45 to 50 years	5.30	59,401	729	12.27	61.35
50 to 55 years	4.67	52,340	852	16.28	81.40
55 to 60 years	3.48	39,003	1,043	26.74	133.70
60 to 65 years	2.80	31,382	1,181	37.63	188.15
65 to 70 years	2.14	23,984	1,224	51.03	255.15
70 to 75 years	1.42	15,915	1,269	79.74	398.70
75 to 80 years	0.81	9,078	1,149	126.57	632.85
80 years and over	0.54	6,052	1,224	202.25	
Unknown	0.08	896	36	40.18	• • • •
Total		1,120,769	15,767	14.07	

## TABLE XIX-C.

ESTIMATED COLORED POPULATION AND DEATHS PER THOUSAND AT THE AGE PERIODS FOR THE YEAR 1915—MARYLAND'S ESTIMATED COLORED POPULATION, 231,171.

Per Cent. Under 5 years. 11.15 5 to 10 years. 11.08 10 to 15 years. 10.56 15 to 20 years. 10.05 20 to 25 years. 10.14 25 to 30 years. 9.06 30 to 35 years. 7.16 35 to 40 years. 7.17	Esti- mated Popu- lation. 25,776 25,614 24,412 23,233 23,441 20,944 16,552 16,575	Deaths. 1,594 115 122 220 291 280 253 298	Mortality Per 1,000 of Those Living at the Age. 61.84 4.49 5.00 9.47 12.41 13.37 15.29 17.98	Mortality Per 1,000 for Age
40 to 45 years       5.84         45 to 50 years       4.96         50 to 55 years       4.06         55 to 60 years       2.64         60 to 65 years       2.22         65 to 70 years       1.63         70 to 75 years       1.09         75 to 80 years       0.52	13,500 11,466 9,385 6,103 5,132 3,768 2,520 1,202	295 303 358 274 254 238 228 <sub>4</sub> 154	21.85 26.43 38.15 44.90 49.49 63.16 90.48 128.12	109.25 132.15 190.75 224.50 247.45 315.80 452.40 640.60
80 years and over. 0.46 Unknown 0.21	1,063 485 231,171	$ \begin{array}{r} 261 \\ 45 \\ \\ 5,583 \end{array} $	$ \begin{array}{r} 245.53 \\ 92.78 \\ \hline 24.15 \end{array} $	••••




Chart 5—Distribution of the Population and Deaths by Ages, Maryland, 1915.

The figures for the total population (Table 19-A) are graphically shown in Chart 5. The profile of the black area, including the portion below, indicates the population; the number of deaths is shown in the black area described below.

From the chart it is seen that the curve plotted for population and the curve plotted for deaths bear no relation to one another except in the first period 0-5 years. The deaths are fewer in the larger population, and as the population diminishes, the deaths increase.

The diseases producing the general mortality analyzed in the preceding tables are shown in Tables A and B, at the end of the report, and in the special tables of this section.

It is evident from the sanitary point of view that important differences exist in the nature of these diseases in their relations to the public health, both as regards their infectiousness and the possibility of their prevention; accordingly, the diseases have been classified in the table from a strictly sanitary standpoint.

The following main divisions have been recognized—parasitic diseases, constitutional diseases, congenital diseases and malformations, poisonings and intoxications, malignant neoplasms, degenerations, pregnancy and violence; other obscure affections not properly falling in one of the previous classes.

The most important of these classes is that including parasitic diseases, which are mostly communicable, and are to a large extent subject to sanitary control. All diseases of parasitic origin may be assumed to be communicable to a certain degree, although important differences exist as to the amount and extent of their contagiousness. Accordingly, these diseases are considered in three classes: (a) infectious and contagious diseases; (b) communicable diseases; (c) other infections of parasitic origin.

The diseases classified under each heading are shown in Table XX. Parasitic diseases form a class largely under the control of sanitary authorities. Poisonings and intoxications are to some extent controlled by law. Constitutional dyscrasiae are not, as a rule, subject to administrative control, while congenital diseases and malformations are wholly beyond administrative influence. Deaths from violence are also usually outside the control of sanitary authorities. The diseases classified under degenerations and malignant neoplasms form a class of

maladies which are imperfectly understood, and are accordingly not capable of control by our present methods. The comparative importance of these classes as causes of death is graphically shown in Chart 6.

### TABLE XX.

A CLASSIFICATION OF CAUSES OF DEATH, WITH THE NUMBER OCCURRING AND THEIR RATIOS TO THE MORTALITY (MARYLAND, 1915).

7.1		70		n a of
Diseases. PARASITIC DISEASES. (Infectious and Contagious Diseases.)	Balto. City.	Rural Mary- land.	Total.	P. C. of Total Mort.
Typhoid fever, scarlatina, whooping cough, diphtheria, influenza, smallpox, measles, glanders, anthrax, actinomycosis	346	592	938	4.39
(*Communicable Diseases.) Malaria, dysentery, tuberculosis, syphilis, tetanus, pneumonia, gonorrhœa, rabies, erysipelas	1,748	2,345	4,093	19.17
(†Other Infections.)  Septicæmia, pyemia, rheumatism (febril), meningitis, bronchitis, broncho-pneumonia, gastro-intestinal inflammations (summer diarrhea of infants), cholera nostras, tonsilitis, pharyngitis, cholecystitis (and other inflammations of the liver and gall bladder), pericarditis, cystitis, peritonitis, acute nephritis, gangrene, abscess, furuncle, pleurisy, appendicitis, laryngitis, metritis, endometritis, endocarditis			9	
(acute), salpingitis, pellagra	1,718	2,065	3,783	17.72
Total Parasitic Diseases	3,812	5,002	8,814	41.28
Constitutional Dyscrasias.  Diabetes, exophthalmic goitre, gout, anæmia, chlorosis, leukemia, Addison's disease, diseases of the thyroid body	<b>1</b> 63	159	322	1.51
Congenital Diseases and Malformations.  Morbus ceruleus, icterus neonatorum, marasmus, sclerema	541	726	1,267	5.93
Poisonings and Intoxications. Alcoholism, saturnism and occupational intoxications, scorbutus	30	60	, 90	0.42
Malignant Neoplasms. Epithelioma, carcinoma, sarcoma	600	608	1,208	5.66
†Degenerations.  Cerebral congestion and hemorrhage, paralysis (without specified cause), meningo-encephalitis, cerebral softening (?), epilepsy, organic diseases of the heart, angina pectoris, arteriosclerosis, aneurism (and allied arterial degenerations), asthma (in all forms), Bright's disease, senile debility and dementia, locomotor ataxia, myelitis, insanity.		3,555	6,344	29.71

n	T A	DI	10	XX-	a	20 4 60	Low
- 1	14		1 1		- ( ) ( )	HLLUI	unea.

THE THE CONTINUE	ve a.			
	Balto.	Rural Mary-		P. C. of Total
Pregnancy.	City.	land.	Total.	Mort.
Puerperal hemorrhage, puerperal septicæmia, puerperal albuminuria and convulsions, pleg-				
masia alba dolens	<b>7</b> 9	112	191	0.89
~				
VIOLENCE.				
Suicide, homicide, murder, dueling, accidental				
violence, poisoning, gas inhalation, drowning,				
strangulation and legal execution, death by				
insulation, lightning, freezing, burns and scalds	573	698	1,271	5.95
All other causes	740	1,103	1,843	8.63
-				
Total	9,327	12,023	21,350	

\*All communicable diseases have been assumed to be due to a living organism, and included in this list whether the specific cause has been discovered or not. †The distinction between these three classes is one of kind, rather than degree, as all parasitic diseases may at some time be communicable. ‡Includes mainly the disorders dependent on advanced years and prolonged strain.

strain.




CHART 6—Maryland Classification of Causes of Death, 1915

# Principal Causes of Death.

The principal causes of death are separately considered, as together they cause about 75 per cent. of the total mortality, and they are in the main preventable diseases.

In Table XXI the twenty principal causes of death are tabulated for Rural Maryland, Baltimore City and total Maryland. The causes come in their proper numerical order under the column marked "Total Maryland." The percentages and figures are entered in the columns marked "Rural Maryland" and "Baltimore City," but do not follow in numerical order.

Deaths from tuberculosis always occupies first place in this table, and deaths from organic heart disease and chronic Bright's disease occupy, respectively, the second and third places. Deaths from cerebral congestion and hemorrhage will always be found in the upper part of this table, and causes like diarrhoea and enteritis, broncho-pneumonia, typhoid fever and influenza or other infectious diseases should gradually take lower positions. Diphtheria for a number of years has been eliminated from the table altogether, while such causes as accidental violence and malignant neoplasms will gradually assume more important positions.

TABLE XXI.

TWENTY PRINCIPAL CAUSES OF DEATH, MARYLAND, 1915.

(	.000,0	.tali		16.52	13.72	19.75 25.55	8.94	8.70	<b>x</b> .54	8.13 8.13 1.13 1.13 1.13 1.13 1.13 1.13	7.47	6.58	20 9 Si o	2:40	1.37	?; ;	2 S	1.66	1.01	0.90	0.75	67.7	
MARYLAND.	t. al lity.	01.10 Tot		10.46	9 9 9 9	04.8 9.40 9.40	2.66	5.51	5.41	5.54	4.73	4.17	1.76	1.52	1.50	1.40	1.17	1.05	0.0₹	0.57	0.47	<del>0.10</del>	
		·841	рәп	2,233	1,855	1,793	1,208	1,176	1,155	1,119	1,010	890	376	324	320	598 -	520	150	137	122	101	97	
Crry.	.000,	ilut 01 r	olf Pe	14.04	15.00	14.76	10.26	10.35	8.19	8.26	8.62	6.65	1.37	2.09	2.48	1.61	1.42	1.66	1.18	0.51	0.36	0.68	
BALTIMORE CITY.		Lot		8.80	9.40	9.25 6.65	6.43	6.40	5.14	5.18	5.40	4.17	0.86	1.31	1.55	1.01	0.80	1.04	0.74	0.32	0.23	0.43	
BAI		.81[1	noo Q	821	877	863	88	605	479	483	504	383	8	122	145	<del>1</del> 6	es S	97	69	30	21	40	
AND.	*000°		170]( 194	18.40	12.75	12.12	0 0 0 0 0 0 0 0 0 0 0	7.4	8.81	8.29	6.59	6.53	3.86	2.63	2.28	5.66	2.18	1.67	0.89	1.20	1.04	0.74	
RUBAL MARYLAND.		101		11.74	8.13	7.74	5.92 5.06	4.75 5.75	5.62	5.29	4.21	4.17	2.46	1.68	1.46	1.70	1.39	1.06	0.57	0.77	0.67	0.47	
RURA		·sų	<sub>1</sub> və (	*1.412	978	930	200	. 575	929	636		202	296	205	13	20 <del>1</del>	167	128	89	36	8	57	
				The majoris Pulmonary and Laryngeal	Organic Heart Disease	Chronic Bright's Disease	Cerebral Congestion and Hemorrhage	Mangnant Neoplasms	Diemboo and Enteritie (under 2 vears)	Concenited Dobility	Unigenital Debitely	Academtal Violence	Davalucia	Tunboid Down	A fractions of the Arteries	Souile Debility	Diambea and enteritis (2 years and over)	Influence	Castrio Diseases	Commissions (Infantila)	Whoming Cough	Meningitis	

\*Including 357 deaths of residents of Baltimore City who died at tuberculosis sanatoria.




Chart 7—Twenty Principal Causes of Death, 1915.


For the purpose of making a special study of tuberculosis Table XXII is inserted which gives the deaths from tuberculosis by age periods, percentage of deaths in age periods and percentage per 10,000 of the population living at the various age periods, white and colored. A much higher death rate from tuberculosis in the colored race per 10,000 of the living population is seen throughout. The excess of deaths in the colored population is extremely disproportionate up to the eightieth year.

On account of the number of deaths occurring from tuberculosis at early ages, we have always been inclined to feel that tuberculosis is a disease of early adult life but it is evident from this table and the accompanying chart that in the white population at least, deaths from tuberculosis are equally prevalent throughout life after the twentieth year. The chart for tuberculosis for the colored population is rather irregular but I attribute this to the difficulty in receiving proper returns of ages from this part of our population. Chart 9, which gives the percentage of deaths of total deaths from tuberculosis which occur at various age periods I think, is the ordinary conception which we have had of the distribution of deaths from this cause.

Howard County shows the smallest number of deaths per 10,000 among the white population while Baltimore County shows the highest. This is due of course to the large number of deaths from tuberculosis at Bay View Asylum. Garrett County gives the lowest death rate among the colored population. However, the colored population in this county is so small that the statistics are not of much value when considered separately. The greatest number of deaths among the colored population occurred in Baltimore County.


PUL

	AT AGE		8.	ц1рэ До	дигу Тар	ьы. Тот	3.57	2.05	4.40	12.02	15.95	13.69	11.55	9.64	6.90	6.79	5.12	3.81	2.14	. 1.31	0.00	0.12	0.24	0.12		:
	OPULATION	Colored.		ıə	.000 A 84	Deat 10,01	11.64	6.64	15.16	43.47	57.16	54.91	58.60	48.87	42.96	49.71	45.82	52.43	35.07	29.19	19.84	8.32	18.81	20.62		36.34
	Per 10,000 of Population at and Colored.	Co			ւթլուն Əւթս		25,776	25,614	24,412	23,233	23,441	20,944	16,552	16.575	13,500	11,466	9,385	6,103	5,132	3,768	2,520	1,202	1,063	485		231,171
	-4	ĺ			*8¥.	1v3 A	. Og	17	50	101	134	115	97	8.	58	57		c1 :	18	11	ರ		21	1	1	0 <del>1</del> 8
XXII.	Ages, Period,		8.	յօ Ծարթ	Cent. Talbi	0.T 0.T	0.93	0.57	0.79	6.03	11.41	12.13	11.63	10.70	11.20	7.25	6.25	6.60	5.17	4.38	2.51	1.65	0.36	0.43		:
TABLE	DEATHS BY	₩нте. √		.19,	.000 A 84	Deat ,01	1.10	0.70	0.99	7.62	15.14	18.02	$\frac{19.40}{10.20}$	18.70	22.74	17.00	16.62	23.59	22.94	25.43	21.99	25.34	8.26	96.99		12.43
	TABLE AMII.  TUBERCULOSIS—1915—GIVING DEATHS BY AGES, DEATHS PERIODS, PER CENT, IN EACH AGE PERIOD, WHITE.		b noit	ıvınd Əşvu	ii18A .oA	117,681	113,534	110,620	110,284	105,016	93,808	83,497	79,687	68,591	59,401	52,340	39,003	31,382	23,984	15,915	9,078	6,052	896		1,120,769	
	osis—19 Periods, 1	ł			*8 <i>ų</i>	реар	13	œ	11	# 5	159	169	162	. 149	156	$\frac{101}{\tilde{0}1}$	87 87	33 (	21	19	65	60 61	ည	9		1,393
	PULMONARY TUBERCUI					AGES.	Under 5 years	5 to 10 years	to 15	to 20	to 25	to 30.	 	to 40	to #5	to 50	to 55	to 60	10 10 10	5 10	to	to 80	-80 years and over	Unknown		Total



TUBERCULOSIS, 1915.

Chart 8—Comparative Chart Giving Deaths per 10,000 at Various Age Periods, Maryland, White and Colored.



TUBERCULOSIS, 1915.

Chart 9—Comparative Chart Demonstrating the Percentage of Deaths at Various Age Periods from Tuberculosis, White and Colored, Maryland.

## TABLE XXIII.

Deaths Per 10,000 of Population from Pulmonary Tuberculosis— White and Colored—Maryland—1915.

County.	White.	Colored.	Total.
Allegany	6.10	27.87	6.57
Anne Arundel	8.46	28.87	15.44
*Baltimore	27.88	113.44 .	35.97
Calvert	9.29	20.02	14.45
Caroline	8.92	21.66	12.03
Carroll	12.18	10.28	12.07
Cecil	8.91	32.45	12.03
Charles	10.38	28.73	19.73
Dorchester	11.19	46.87	22.72
†Frederick	18.96	29.55	19.97
Garrett	5.64	0.00	5.61
Harford	6.07	35.98	11.15
Howard	4.86	29.06	10.14
Kent	9.52	41.92	20.64
Montgomery	9.94	24.96	13.96
Prince George's	9.93	25.78	14.45
Queen Anne's	13.32	34.41	20.58
Somerset	17.35	26.47	20.57
St. Mary's	14.78	39.69	24.78
Talbot	12.48	24.94	16,63
Washington	9.19	20.79	9.62
Wicomico	13.43	19.80	14.88
Worcester	13.77	33.76	20.13
Total Counties	14.04	*37.42	18.40
Baltimore City	10.41	34.57	14.04
Maryland	12.43	36.34	16.52

<sup>\*</sup>Including 323 non-residents of Baltimore County who died at T. B. Sanatoria.

<sup>†</sup>Including 65 non-residents of Frederick County who died at Sabillasville.

In Table XXIV the principal diseases are divided according to the percentages occurring in the three periods of life already used, with the exception of senile debility, congenital debility, infantile convulsions, unspecified and ill-defined causes, the three former of which fall by reason of their classification in only one period of life.

Deaths in the middle period of life have an importance from the economic standpoint far greater than those occurring in either extreme. In Chart 10, the principal causes of death are arranged according to their importance as causes of death between the ages of 15 and 45 years, for the State of Maryland.

Tuberculosis of the lungs still easily retains first place on this chart. Accidental violence, which ranks tenth in Table 21, comes in second with this arrangement; typhoid fever is third, and Bright's disease fourth. A number of diseases which bear a high ratio to the total number of deaths do not occupy important positions as causes of death in the age periods of 15-45.




Chart 10—Principal Causes of Death, Classified According to their Importance in the Middle Period of Life.

TABLE XXIV.

DEATHS FROM FIFTEEN PRINCIPAL CAUSES IN MARYLAND, 1915, SHOWING NUMBER AND PERCENTAGE OF DEATHS IN EACH OF THREE AGE PERIODS.

	RURAI	L MARYLAND.	BALT	IMORE CITY.	MA	MARYLAND,			
TYPHOID FEVER.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.			
Under 15		22.28	17	13.93	62	19.14			
15 to 45		59.90	83	68.03	204	62.96			
45 and over		$17.33 \\ 0.50$	$\frac{22}{0}$	$\frac{18.03}{0.00}$	$\begin{array}{c} 57 \\ 1 \end{array}$	$17.59 \\ 0.31$			
Age duknown		0.90		0.00		0.51			
Total	202		122		324	•			
PARALYSIS.									
Under 15	1	0.48	0	0.00	1	0.37			
15 to 45 45 and over	$\begin{array}{c} 7 \\ 198 \end{array}$	3.38	3	4.76	10	3.70			
Age unknown	198	95.65 $0.48$	60	$95.24 \\ 0.00$	$258 \\ 1$	$95.56 \\ 0.37$			
rige dinkhown		. 0.10		0.00		0.51			
Total	207		63	-	270				
Bright's Disease.									
Under 15		3.98	17	1.73	60	2.91			
15 to 45	145	13.43	193	19.67	338	16.40			
45 and over	883 9	$81.76 \\ 0.83$	$\begin{array}{c} 771 \\ 0 \end{array}$	$78.59 \\ 0.00$	1,654	$80.25 \\ 0.44$			
Age unknown		0.88		0.00	9	0.44			
Total	1,080		981		2,061				
Whooping Cough.									
Under 15	79	98.75	21	100.00	100	99.01			
15 to 45	0	0.00	0	0.00	0	0.00			
45 and over	$\frac{1}{0}$	$\frac{1.25}{0.00}$	0	0.00	1	0.99			
Age unknown		0.00		0.00	0	0.00			
Total	80		21		101				
HEART DISEASE.									
Under 15	20	2.04	23	2.62	43	2.32			
15 to 45	96	9.82	158	18.02	254	13.69			
45 and over	851	87.01	696	79.36	1,547	83.40			
Age unknown	11	1.12	0	0.00	11	0.59			
Total	978		877		1,855				
Malignant Neoplasms.									
Under 15	7	1.15	3	0.50	10	0.83			
15 to 45		12.66	105	$\frac{0.50}{17.50}$	182	15.07			
45 and over	523	86.02	492	82.00	1.015	84.02			
Age unknown	1	0.16	0	0.00	1	0.08			
Total	608		600		1,208				
Broncho-					_, <b>_</b> 00				
PNEUMONIA.									
Under 15	304	60.08	335	66.47	639	63.27			
15 to 45		$\frac{00.55}{2.57}$	39	7.74	$\frac{050}{52}$	5.15			
45 and over	189	37.35	130	25.79	319	31.58			
Age unknown	0	0.00	0	0.00	0	0.00			
Total	506		504		1,010				

## TABLE XXIV-Continued.

DEATHS FROM FIFTEEN PRINCIPAL CAUSES IN MARYLAND, 1915, SHOWING NUMBER AND PERCENTAGE OF DEATHS IN EACH OF THREE AGE PERIODS.

	RURAI	L MARYLAND.	BALT	IMORE CITY.	MARYLAND.			
CEREBRAL HEMORRHAGE. Under 15	$\begin{array}{c} 25 \\ 682 \end{array}$	Per Cent. 0.00 3.51 95.79 0.70	No. 3 47 570 0	Per Cent. 0.48 7.58 91.94 0.00	No. 3 72 1,252 5	Per Cent. 0.23 5.41 93.99 0.38		
Total  Diarrhæa and Enteritis.	712		620		1,332			
Under 15	13 99 1 —	86.60 1.54 11.74 0.12	$ \begin{array}{r} 502 \\ 17 \\ 43 \\ 0 \\ \hline 562 \end{array} $	89.32 3.02 7.65 0.00	$   \begin{array}{c}     1,232 \\     30 \\     142 \\     \hline     1,405   \end{array} $	87.69 2.14 10.11 0.07		
Total  Influenza. Under 15 15 to 45 45 and over Age unknown	15 10 100	11.72 $7.81$ $78.13$ $2.34$	7 29 61 0	7,22 29,90 62,89 0.00	1,403 22 39 161 3	9.78 17.33 71.56 1.33		
Total  Meningitis. Under 15	128 45 7 5	78.95 12.28 8.77 0.00	97 30 8 2 0	75.00 20.00 5.00 0.00	75 15 7 0	77.32 15.46 7.22 0.00		
Total  ACCIDENTAL VIOLENCE. Under 15	123 205 159 14	24,55 40,92 31,74 2,79	$ \begin{array}{r} 105 \\ 132 \\ 152 \\ 0 \\ \hline 389 \end{array} $	26.99 33.93 39.07 0.00	97  228 337 311 14 890	25.62 37.87 34.94 1.57		
Tuberculosis (Lungs and Larynx). Under 15	914 422 7	4.89 64.73 29.89 0.50	$ \begin{array}{r} 47 \\ 551 \\ 223 \\ 0 \\ \hline 821 \end{array} $	5.72 67.11 27.16 0.00	116 1,465 645 7 	5.19 65.61 28.88 0.31		

### TABLE XXIV-Continued.

DEATHS FROM FIFTEEN PRINCIPAL CAUSES IN MARYLAND, 1915, SHOWING NUMBER AND PERCENTAGE OF DEATHS IN EACH OF THREE AGE PERIODS.

	RURAI	L MARYLAND.	BALT	IMORE CITY.	MA	RYLAND.
PNEUMONIA.	No.	Per Cent.	No.	Per Cent.	No.	Per Cent.
Under 15		23.82	109	18.02	245	20.83
15 to 45		22.94	159	26.28	290	24.66
45 and over		52.54	337	55.70	637	54.17
Age unknown	4	0.70	0	0.00	4	0.34
Total	571		605		1,176	
OTHER CAUSES.						
Congenital Debility						
(under 1 year)			483		1,119	
Senile Debility (over						
50 years)			94		298	
Convulsions of Chil-						
dren (less than 10 years)			30		122	
Unspecified or ill-			50		122	
defined causes			28		383	

The age distribution of the principal causes of death is shown in Charts Nos. 12, 13, 14 and 15. The age distribution of the general mortality is shown in Chart No. 11. These charts illustrate the age distribution of the principal causes of death, with the exception of senile debility, congenital debility and infantile convulsions (whose age distribution is fixed by their classification) and the unclassified diseases. By referring to Chart No. 11 it will be seen that as has been shown in previous charts, a great portion, 24.31 per cent. of the general mortality occurs under the age of 5. The percentage rapidly declines to the ages between 10 and 15 years, 1.66 per cent., when the lowest actual mortality is reached. The percentage of deaths thereafter remains fairly constant until after the age period 45-50, when it increases. The general percentage ranges below 5 per cent. for all periods after the first quinquennium until the age period of 55-60 years. The percentage of deaths from lobar pneumonia has a large distribution corresponding closely to that of the general mortality. If the pneumonia and broncho-pneumonia curves were combined, the analogy would appear even more striking. This fact shows that the term pneumonia as generally employed, is composite, and relates to a number of diseases, mostly infectious, of which pneumonia is the terminal or secondary symptom. Both pneumonia and

broncho-pneumonia are important causes of death only at the extremes of life. After the age of 10 years, broncho-pneumonia becomes a small factor in the mortality, except for persons over 65, although it will be seen from the chart the largest proportion of deaths from this cause is in the first five years of life. Broncho-pneumonia and whooping cough have an almost wholly infantile distribution. 87.50 per cent. of the deaths from whooping cough occur in the first five years of life; 7.50 per cent. in the age period of 5-10. 55.08 per cent. of all deaths from diphtheria appears in the first quinquennium and 27.96 per cent. in the second.

Organic diseases of the heart, malignant neoplasms, cerebral hemorrhage, paralysis and Bright's disease occur almost entirely in the late periods of life. The mortality from cerebral hemorrhage rises above 5 per cent. at the fiftieth year, and increases rapidly thereafter and reaches its maximum between 65 and 70. Malignant neoplasms rise above 5 per cent. at from 40 to 45 and reaches its greatest height at 60-65. Bright's disease rises above 5 per cent. between 45 and 50; paralysis between 60 and 65 and organic heart disease between 50 and 55.

Tuberculosis—The greatest number of deaths from pulmonary tuberculosis occurs in early adult life. It will be noted in the chart that the percentage remains low to the 15th year, when there is a rapid rise, the curve reaching its maximum between 20 and 25 years, declining slowly until it reaches the 75th year.

Typhoid Fever—The greater number of deaths from typhoid fever occur in early adult life. The maximum is reached at 15-20 years.

The causes of accidental violence in infancy are due to the inability of the children to protect themselves and to the carelessness of parents. Burns and scalds account for a large portion of this mortality, and accidental drowning for a further considerable portion. A large number of these deaths are among the children of colored persons and foreigners, and are due, on the one hand, to the carelessness in handling fire, and on the other, to the ready access to water afforded by the Maryland coast line and Chesapeake Bay and its tributaries. The percentages are shown in Table XXV.

TABLE XXV.

AGE DISTRIBUTION OF MORTALITY FROM TWELVE DISEASES BY PERCENTAGES, RURAL MARYLAND, 1915.

амоплаП	0.50	0.00	0.50	0.70	0.00	1.12	0.48	0.83	2.79	0.16	0.00	0.70	0.67
80 years and over.													
.08-37													
.87-07													
.07-58													
.59-09													
•09 <u>-</u> 66									-				
50-55.													
.06-64	4.95	0.00	6.73	2.63	1.38	4.81	5.90	5.28	5.99	7.24	0.00	4.64	4.00
.64-04	5.94	0.00	9.21	3,33	0.50	3.48	1.93	4.35	3.59	6.25	1.69	1.40	3.79
32-40.	7.92	0.00	9.70	4.73	0.79	2.35	0.48	3.33	6.33	2.63	2.54	1.26	3.93
.56-95.	5.45	0.00	11.54	4.20	0.40	1.43	0.00	2.55	6.79	2.14	0.85	0.42	3.71
.55-30.	10.40	0.00	12.25	5.08	0.59	1.02	0.48	1.39	7.78	1.15	0.85	0.42	3.53
20-25.	13.37	0.00	14.09	2.80 80	0.50	0.51	0.00	1.30	8:38	0.16	0.00	8.0	3.56
.02-20	16.83	0.00	7.93	2.80	0.40	1.02	0.48	0.83	7.98	0.33	1.69	0.00	2.69
.61–01	8.91	3.75	2.55	1.23	0.40	0.92	0.48	0.74	6.59	0.33	6.78	0.00	1.66
.01-G											•		
.ā-0	. 5.94	87.50	다. 다.	19.96	58.30	0.95	0.00		12.18	0.49			
f Death.	P	Whooping Cough	Fulmonary.	Lobar	monia	t Disease		ase	olence	Malignant Neoplasms		Cerebral Hemorrhage	uses
Causes of Death.	Typhoid Fever	noohing Co	perculosis,	Fuenmonia, Lobar.	oncho-Pneu	Organic Heart Disease.	Faralysis Beeter Bee	bright s Disease	cluental v	Highant No	Diplitheria	rebrai Hen	Deaths, all causes.
		= [	7 .	Z 6	i d	5	ਮ ਮ ਜ	DI -	AC	i i	1 2	בי בי	De





CHART 11—Age Distribution of General Mortality, 1915.



**CHART 12.** 




CHART 13.




CHART 14.




CHART 15.




Chart 16-Mortality from Five Respiratory Diseases, 1907-1915.

# TABLE XXVI.

DEATHS FROM FIVE PRINCIPAL RESPIRATORY DISEASES IN RURAL MARYLAND, 1907-1915, INCLUSIVE.

FAT	E I	3U.	AK	עו	O.	B,	HE	ALTI	н.
	1915	1.06	0.95	4.21	4.75		11.74		
	1914	0.99	0.61	4.37	4.13		11.21		
tality.	1913	1.33	0.61	4.26	3.36		10.10		
u Mor	1912	0.90	0.91	3.53	3.87		9.74		
of Tote	1911	1.31	0.81	3.87	4.25		10.70		
Percentage of Total Mortality.	1910	1.60	0.89	2.75	5.81		11,19		
Percei	1909	1.91	1.18	1.99	5.61		11.16		
	1908	2.55	1.01	1.85	4.76		10.65		
	1907	2.93	1.05	1.62	5.45		10.35		
	1915	128	111	506	571		*1,412	12,023	
	1914	107	99	473	447		1,214 *	-	274 41 68
			99				1,099 1	10,880 10	
	1912	91	35	358	393		989 1	10,156 10	micipal Tuberculosis Hospital
Deaths.			<b>%</b>				1,058	(C)	Hosp
De									osis
	-1910	156	87	569	568		1,094	9,775	nbercul
	1909	186	115	194	548		1,089	9,762	pal Tu
	1908	251	66	185	469		1,049	9,849	Munici Sanate
	1907	287	103	159	Pneumonia 534		culosis1,013 1,049	Total Deaths9,792 9,849	Bay View Municipal Tuberculosis Hospital Includes   Eudowood Sanatorium
		:		опіа	:	Pulmonary Tuber-		70	Bay Eudo Sabil
		:	nchi	eum	:	Τn	:	ath	S
		za .	Bro	-Pu	nia	ary	<u>x</u>	De	Jude
		Influenza	ite	nche	ume	mon	ulosi	'otal	*Inc
		Infl	Acu	Bro	Рпе	Pul	G	1	

Table XXVI gives the deaths from five principal respiratory diseases in Rural Maryland for the years 1907 to 1915, inclusive. In this table all of the figures show considerable variation except for broncho-pneumonia which has shown an increase. The figures given for pulmonary tuberculosis in Rural Maryland are always too high. While the general rate in the State as a whole has fallen, that for Rural Maryland has been increasing more so since the establishment of tuberculosis sanatoria.

Table XXVII gives the deaths from pulmonary and laryngeal tuberculosis by sex, percentage of the total mortality and the mortality per 10,000 for the years 1908 to 1915 inclusive. In general, the tuberculosis rates have shown a gratifying decrease. This decrease was very steady from the year 1908 to the year 1913. In 1914, the mortality rates per 10,000 exceeded those for the three previous years and the rate per 10,000 for the year 1915 still remains higher than the rate for 1912 and 1913. At the foot of Table XXVII, average figures for the eight years are given. It is evident that in Maryland, tuberculosis is more prevalent in the male population. The average rate per 10,000 for males is 18.05 and for females 16.21.

TABLE XXVII.


PULMONARY AND LARYNGEAL TUBERCULOSIS, 1908-1915.

		Mortalit	y	Per Cent.	Mor- tality
T/one	7/.7.		m. 1. 7	Mor-	Per
Year.	Male.	Female.	Total.	tality.	10,000.
1908	1,187	1,112	2,299	11.33.	18.02
1909	1,216	1,146	2,362	11.73	18.36
1910	1,192	1,136	2,328	11.10	17.94
1911	1,170	1,053	2,223	10.96	16.99
1912	1,175	1,003	2,178	10.57	16.51
1913	1,037	979	2,016	9.58	15.16
1914	1,274	1,085	2,359	11.03	17.59
1915	1,192	1,041	2,233	10.46	16.52
Total	9,443	8,555	17,998		
Average for 8 years	1,180	1,070	2,250		
Average Population6	53,731	660,178	1,313,909		
Death rate per 10,000	18.05	16.21	17.12		

# Seasonal Incidence of the Mortality.

The seasonal incidence of the general mortality is shown separately for Baltimore City and Rural Maryland in Chart No. 17.

It will be seen that the monthly mortality, as shown in the black portion of the charts, is of somewhat oval shape, with the poles lying in the winter and summer months, which are the seasons of greatest mortality. Generally speaking, the mortality may be divided into two classes, diseases of the gastrointestinal tract, prevalent in the summer months, and diseases of the respiratory tract, prevalent in the winter. The gastrointestinal diseases reach their altitude in the height of the March is the month of greatest mortality in Rural Maryland and in Baltimore City. June was the month of least mortality in both Rural Maryland and in Baltimore City. The expansion of the black portion of the chart during July and August is due almost entirely to the acute gastro-intestinal dis-The mortality during the winter months eases of children. is due mainly to pneumonia and influenza. While the summer mortality affects mainly children, the winter mortality affects both extremes of life (infancy and old age). The sudden rise from the months of low mortality, May and June, to the months of high mortality, July and August, produces a very notable eccentricity in the figures. The circles on the charts indicate the actual number of deaths in the two specified divisions of Maryland. A comparison of this chart with that of the preceding year shows a marked correspondence. The figure of the chart is apparently not subject to many changes, although it may rotate somewhat, either to a later or an earlier season. In the winter months, March furnished the highest mortality. In 1914 March furnished the highest mortality for Rural Maryland and for Baltimore City.



In the following table the deaths by months for six years, ending 1915, are given for Baltimore City and Rural Maryland (Table 28). From this table it appears that March is the month of greatest mortality in Rural Maryland as well as for Baltimore City. June is the month of least mortality for Rural Maryland and for Baltimore City.

The seasonal prevalence of eight prominent causes of death are given in the succeeding chart, No. 18.

In considering seasonal prevalence, two classes of diseases must be recognized:


- (1) Acute fatal diseases, where there is a short interval between commencing sickness and the date of death.
- (2) Chronic fatal diseases, where the date of death is separated by a long interval from the date of commencing illness.

The influence of pneumonia and influenza upon the tuberculosis mortality will be seen in a glance at the chart. The chronic degenerative diseases, cerebral appoplexy, heart disease and Bright's disease show little seasonal variation, and are apparently not much dependent upon acute infection. The pneumonia curve and influenza curve always correspond very closely. The gastro-intestinal diseases show a marked antithesis to pneumonia in their seasonal distribution. The mortality from acute gastro-intestinal infections shows the greatest seasonal variation of any of these diseases, as already observed. To illustrate the striking influence of pneumonia upon tuberculosis, the curves of these diseases, together with those of acute bronchitis and broncho-pneumonia, have all been introduced in Chart No. 18.

TABLE XXVIII.

DEATHS BY MONTHS FOR 1915, 1914, 1913, 1912, 1911 AND 1910.

			RURAL	. MARYLAND	LAND.		-			BAL	SALTIMORE	CITY.		
							Total							Total
	1915.		1913.	1912.	1911.		6 Years.	1915.	1914.	1913.	1912.	1911.	1910.	6 Years.
January	974		888	901	905		5,511	648	686	935	966	993	1,067	5,829
February	976	923	876	885	843	956	5,399	739	943	914	925	871	896	5,300
March	1,192		991	899	953		5,918	952	1,129	955	1,056	686	940	6,021
April	1,172		846	06 <u>7</u>	803		5.284	808	937	837	698	935	875	5,349
May	952		828	111	069		4.805	55	865	305	082	305	775	4,988
June	803		086	678	662		4,601	633	757	831	701	7.58	262	4,448
July	1,023		1,144	862	1,004		5,915	<u>#9</u> 2	208	9+6	922	65 6	981	5,439
August	1,067		96 <del>1</del>	<del>1</del> 96	1,017		5,928	0 <del>1</del> 2	845	792	830	855	950	4,982
September	965		818	950	848		5,272	0 <u>8</u> 9	771	703	747	752	805	4,458
October	981		006	800	713		5,185	721	736	748	836	798	834	4,673
November	800		807	817	730		4,968	755	805	732	850	801	860	4,800
December	1,076		801	806	721		5,209	831	880	873	929	854	930	5,297
Total	12,023	10,832	10,880	10,156	9,886	10,218	63,995	9,327	10,551	10,168	10,441	10,404	10,753	61,644



**CHART 18.** 

# EIGHT PROMINENT CAUSES OF DEATH BY MONTHS—RURAL - MARYLAND—1915.

### TYPHOID FEVER.

Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
7	12	8	6	. 11	10	15	36	29	28	19	21

### LOBAR PNEUMONIA.

Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
77	66	110	102	32	17	12	7	11	13	41	83

### HEART DISEASE.

Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
86	87	92	83	102	74	87	· 88	69	66	65	79

### BRIGHT'S DISEASE.

Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	$Sear{p}t.$	Oct.	Nov.	Dec.
93	79	113	121	86	73	87	69	86	86	86	101

### INFLUENZA.

Jan.	Fcb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
10	9	35	44	10	1	0	0	1	0	1	17

### TUBERCULOSIS (PULMONARY).

Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
130	124	138	150	130	108	134	116	85	103	84	110

### CEREBRAL HEMORRHAGE.

 Jan.
 Feb.
 Mar.
 Apr.
 May.
 June.
 July.
 Aug.
 Sept.
 Oct.
 Nov.
 Dec.

 56
 62
 64
 64
 64
 60
 57
 44
 50
 60
 50
 81

### DIARRHŒA.

 Jan.
 Feb.
 Mar.
 Apr.
 May.
 June.
 July.
 Aug.
 Sept.
 Oct.
 Nov.
 Dec.

 12
 19
 15
 26
 16
 33
 152
 192
 165
 141
 49
 23

# Eight Prominent Causes of Death.

The succeeding table and chart (Table 29, Chart 19), show the comparative susceptibility of the several organs to infection. In general, the table and charts have been arranged so as to show infections of the special organs. The total number of deaths considered in this table is 8,875, or about 41.57 per cent. of the total mortality. Certain other important classes of diseases, such as degenerations, congenital diseases and accidents, are not considered in this tabulation.

It will be seen by reference to the table and chart that most of the inflammatory diseases affect the lungs and alimentary canal, infections of the lungs ranking very much higher than those of the other organs.

The special predilection of the tubercle baccillus for the lung has been previously recognized, but the fact that a majority of pathogenic organisms find in the lungs a favorable habitat is not so well recognized. The organisms of the enteric group are, of course, one exception to this rule.

### TABLE XXIX.

# PARASITIC DISEASES, SHOWING THE ORGAN AFFECTED.

Pagningtony Infactions W	Rural aryland.	Baltimore City.	Mary- land.
Influenza, laryngeal and pulmonary tubercu-		ouy.	iana.
losis, bronchitis, broncho-pneumonia, pneumonia, pleurisy, pulmonary gangrene		2,179	4,993
Systemic Infections, Principally Respiratory.  Measles, scarlet fever, whooping cough, diph-	261	127	388
theria  Infections of the Alimentary Tract.	201	121	900
Typhoid fever, cholera, dysentery, abdominal tuberculosis, gastritis, diarrhœa and enteritis, peritonitis, appendicitis	1,289	889	2,178
Infections of the Urinary Tract.  Perinephritis, pyonephrosis, pyelitis, cystitis, nephrolithiasis	34	37	71
Systemic Infections, Principally of the			
Kidneys. Acute nephritis	150	118	268
Infections of the Nervous System. Rabies, meningeal tuberculosis, encephalitis, meningitis, tetanus, chorea	144	113	257
Systemic Infections, Not Localized in Any			
Organ or Tissue.  Pyemia and septicemia, general tuberculosis, syphilis	114	139	253
Infections of the Skin. Erysipelas, gangrene, abscess and furuncle, dermatitis	68	47	115
Systemic Infections, Principally of			
Smallpox	0	0	0
Arthritic Infections. Acute rheumatism	27	40	67
Systemic Infections, Principally Arthritic. Chronic rheumatism, gout	26	17	43
Infections of the Female Organs of Reproduction. Endometritis, metritis, salpingitis, puerperal			
septicemia	51	51	102
Infections of the Liver. Cholangitis, hepatitis, cholelithiasis	31	39	- 70
Infections of the Bones. Tuberculosis, osteomyletis	23	. 32	55
Infections of the Blood.	- 4		,
Malaria	14	1	
Total	5,046	3,829	8,875




Chart 19—Comparative Susceptibility of the Several Organs to Parasitic Infections—Maryland Mortality, 1915.

# Special Causes of Death.

Tables A. B and C, at the end of the report, give the causes of death by race, sex, age, county, and seasonal distribution for the 189 principal causes of death recognized in the international classification.

Many of the diseases recognized in this classification are only important causes of mortality during epidemics, and are not commonly prevalent in this State. Others are limited to Europe or other geographic or climatic divisions.

Among the diseases of the International Classification, from which there were no deaths in Maryland during 1915, were typhus fever, relapsing fever, miliary fever, Asiatic cholera, bubonic plague, yellow fever and leprosy.

The following table, which is for comparative purposes, gives the number of deaths from certain special causes not considered separately in this report, for a period of ten years.

### TABLE XXX.

### SPECIAL CAUSES OF DEATH.

S	M	A	L	L	P	o	X.	

1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland 0	0	0	0	0	0	0	2	5	0
Baltimore City 0	0	0	0	0	0	0	0	3	0
Maryland 0	0	0	0	0	0	0	2	8	0
		RAB	IES.			-			
1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland 1	0	0	0	0	1	0	1	0	0
Baltimore City 1	0	6	1	1	2	0	0	0	0
Maryland 2	0	6	1	1	3	0	1	0	0
	-A	LCOH	OLISM.						
1906	1907	1908	1909	1910	1911	1919	1913	1914.	1915
Rural Maryland 34	36	37	44		39	49	56	36	46
Baltimore City 59	53	46	40	49	46	49	57	55	25
Maryland 93	89	83	84	81	85	98	113	91	71
							*		
			Pois						
					1911.	1912.		1914.	1915.
Rural Maryland 1	1	0	0	0	0	0	0	0	1
Baltimore City 1 Maryland 2	0	1 1	1 1	$\frac{2}{2}$	3 3	0	$\frac{2}{2}$	1	3 4
Maryiand 2	1	т	1	_	9	U	_	1	4
DISEA	SES OF	THE	Nervo	us Sy	STEM.				
	· <i>B</i>	Enceph	alitis.						
1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland 21	11	19	7	14		18	10	5	14
Baltimore City 1	3	6	1	8	2	2	9	6	3
Maryland 22	14	25	8	22	12	20	19	11	17
		Menin	aitis.						
1006				1910	1911	1919	1913	1914.	1915.
Rural Maryland 130	124	117	107	103	88	101	105	73	57
Baltimore City 163	161	140	105		- 101	110	88	61	40
Maryland 293		257	212			211	193	134	97
		~							
		Epile							1011
								1914.	
Rural Maryland 39	35	48	30		41	44	56	$\frac{50}{22}$	53 14
Baltimore City 18 Maryland 57	19 54	18 66	15 45	_		15 59	$\frac{12}{68}$	72 72	67
maryianu	94	- 00	40	01	- 50	90	30		

General Paralysis of the Insane and Other Forms of Mental Alienation.

Rural Maryland..... 85

Baltimore City...... 37 Maryland ...... 122

1906. 1907. 1908. 1909. 1910. 1911. 1912. 1913. 1914. 1915.

### TETANUS.

		TETA	NUS.							
1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.	
Rural Maryland 32		19	29	24	22	26	26	21	23	
Baltimore City 27	21	15	19	16	23	32	16	16	11	
Maryland 59	52	34	48	40	45	58	42	37	34	
Intestinal Parasites.										
1906.	1907.	1908.	1909.							
Rural Maryland 2 Baltimore City 0		$\frac{1}{0}$	$\frac{3}{0}$	. 5	6	2	7	- 1	1	
Maryland 2		1	3	$\frac{1}{6}$	0 6	$\frac{0}{2}$	0	$\frac{1}{2}$	0	
		Pregn		Ū	· ·	_	•	_	1	
1000				1010	1011	1010	1019	1014	1015	
Rural Maryland 105	1907. 81	1908.	102	1910.	1911, 76	1912.				
Baltimore City 103	97	92	102		87	$\frac{100}{105}$	$\frac{113}{116}$	95 84	$\frac{112}{79}$	
Maryland 207	178	190	205	192	163	211	229	179	191	
		Suic	IDE.							
By Poison.										
1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.	
Rural Maryland 6	12	6	14	2	5	8	20	15	14	
Baltimore City 26	20	28	31	28	37	30	30	63	29	
Maryland 32	32	34	45	30	42	38	50	78	43	
	B	y Asp	hyxia.							
1906.	1907.	1908.	1909.	1910.	Ī911.	1912.	1913.	1914.	1915.	
Rural Maryland 0	2	2	2	1	0	4	3	0	4	
Baltimore City 7	10	17	12	11	18	22	35	33	24	
Maryland 7	12	19	14	12	18	26	38	33	28	
	Hangi	_								
	1907.	1908.	1909.	1910.		1912.	1913.	1914.	1915.	
Rural Maryland 9	13	10	18	14	12	13	15	15	14	
Baltimore City	$\begin{array}{c} 13 \\ 26 \end{array}$	$\frac{12}{22}$	11 29	5 19	$\frac{7}{19}$	$\begin{array}{c} 6 \\ 19 \end{array}$	$\frac{14}{29}$	10 25	$\frac{6}{20}$	
1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1			2.5 wning		1.0	10	20	29	20	
1906	1907.		_		1911	1912	1913	1914 -	1915	
Rural Maryland 3		4	6	6	5	10	4	3	9	
Baltimore City 2	4	4	8	. 4	$\frac{3}{2}$	13	$\hat{\bar{5}}$	3	3	
Maryland 5	10	8	14	10	7	23	9	6	12	
By Firearms.										
1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.	
Rural Maryland 12	22	22	28	20	16	23	39	30	34	
Baltimore City 20 Maryland 32	$\frac{27}{49}$	23 45	30 58	$\frac{22}{42}$	23 39	$\frac{21}{44}$	22 61	$\frac{24}{54}$	$\begin{array}{c} 31 \\ 65 \end{array}$	
By Cu.						- 11	01	01	00	
	1907.					1912	1913	1914	1915	
Rural Maryland 2	1.001.	3	3	3	2	6	5	4	2	
Baltimore City 6	3	7	3	5	5	2	8	4	$\frac{1}{5}$	
Maryland 8	4	10	6	8	7	8	13	8	7	

TABLE XXX—Continued.										
By Jumping from High Places.										
1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1	915.									
Rural Maryland 0 0 0 0 0 0 0 1 3	3									
Baltimore City 3 5 4 6 1 7 2 4 1	3									
Maryland	6									
,										
By Crushing or Other Means.										
1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1	915									
	2									
Rural Maryland 6 2 3 2 1 3 2 1 1 Baltimore City 2 1 3 0 1 1 2 0 0	$\bar{0}$									
Maryland	$\overset{\circ}{2}$									
	_									
ACCIDENTAL VIOLENCE.										
Poisoning by Food and Other Acute Poisonings.										
1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1	915.									
Rural Maryland 17 18 22 26 29 20 15 21 21	26									
Baltimore City 16 14 23 12 \( 9 \) 6 13 20 29	13									
Maryland	39									
Burns and Scalds and Burns by Corrosive Substances (Conflagration included from 1910 through 1915).										
	1015									
1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1										
Rural Maryland 70 54 68 64 56 50 58 59 69 Baltimore City 70 81 65 76 86 68 67 69 69	-88									
	155									
Maryland	100									
Absorption of Deleterious Gases (Conflagration included from 1906 through 19	909).									
1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1	1915.									
Rural Maryland 4 5 3 6 13 18 16 19 9	6									
Baltimore City 12 17 15 12 16 17 14 25 30	10									
Maryland 16 22 18 18 29 35 30 44 39	16									
Accidental Drowning.										
1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1	1915.									
Rural Maryland 90 101 107 98 79 95 115 110 105	141									
Baltimore City 61 54 55 47 49 45 52 58 46	58									
Maryland	199									
*Traumatism by Firearms.										
1910. 1911. 1912. 1913. 1914.										
Rural Maryland	20									
Baltimore City 7 8 1 3 5	7									
Maryland 35 29 22 31 35	27									
*Traumatism by Cutting or Piercing Instruments.										
1910. 1911. 1912. 1913. 1914. 1	1915.									
	2									
Baltimore City 3 1 0 0 2	$^{2}$									
Maryland	4									
•										

# \*Traumatism by Fall.

Traumation by	L wee.											
	1910.	1911.	1912.	1913.	1914.	1915.						
Rural Maryland		44	53	66	52	76						
		91	94	86	100	91						
Baltimore City												
Maryland	. 102	199	147	152	152	167						
•												
*Traumatism in Mines and Quarries.												
1910. 1911. 1912. 1913. 1914. 1915.												
				1913.	1914.	1919.						
Rural Maryland:	. 8	10	12	11	25	7						
Baltimore City	. 1	0	1	. 0	0	1						
Maryland	. 9	10	13	11	25	8						
*Traumation by Machines												
*Traumatism by Machines.												
	1910.	1911.	1912.	1913.	1914.	1915.						
Rural Maryland			7	7	11	10						
Baltimore City			15	14	6	10						
Maryland		_	22	$\frac{11}{21}$	17	$\frac{10}{20}$						
Maryland	. 10	10	ت ت	21	1.	20						
•												
*Traumatism by Crushing (Vehicles, Railroads, Landslides).												
	1010	1911.	1019	1012	1014	1015						
Rural Maryland			155	150	172	146						
Baltimore City			64	96	108	109						
Maryland	. 206	206	219	246	280	255						
*Electrical Shock (Lightni	na Ela	ccenter	7.)									
· · · · · · · · · · · · · · · · · · ·		_										
		1911.	1912.	1913.	1914.	1915.						
Rural Maryland	. 3	4	6	11	4	4						
Baltimore City	. 9	1	6	3	1	1						
Maryland	. 12	$\frac{1}{5}$	12	14	5	5						
Lightning.												
Lightheny.												
		1911.	1912.	1913	1914.	1915.						
Rural Maryland	. 2	4	7	9	6	9						
Baltimore City		3	Ó	0	0	0						
Maryland			7	9	6	9						
	_											
Drowning.												
DROWNING.												
Accidental and Suicidal.												
						1015						

	1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland	. 93	107	111	104	85	100	125	114	108	150
Baltimore City	. 63	58	59	55	53	47	55	63	49	61
Maryland	. 156	165	170	159	138	147	180	177	157	211

# UNSPECIFIED OR ILL-DEFINED CAUSES. .

# Ill-Defined Organic Causes.

1900	3. 1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland 4	0 27	30	36	22	5	4	1	1	5
Baltimore City	7 4	9	7	5	6	4	$^{2}$	0	1
Maryland	7 21	39	43	27	11	8	3	1	6

### Sudden Death.

	1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland	26	25	27	44	19	26	31	24	25	25
Baltimore City										
Maryland	27	25	27	44	19	26	32	24	25	27

# Unspecified Causes.

	1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.
Rural Maryland	. 425	503	421	347	437	367	220	208	283	325
Baltimore City	. 280	295	308	-326	313	-226	85	45	10	25
Maryland	. 705	798	729	673	750	593	305	253	293	350

<sup>\*</sup>Classified under one title, "Other Accidental Traumatisms," during the years 1906, 1907, 1908 and 1909.

Table XXX permits of rapid comparative observation for a period of ten years both for Rural Maryland and the City of Baltimore.

During the year 1915 there were no deaths from smallpox in Rural Maryland and Baltimore City. Except for the year 1913, in which there were two deaths from this cause, and 1914, in which there were eight deaths, not a single death from smallpox has been recorded for a period of seven years.

There were 71 deaths from alcoholism in 1915, a decrease of 20 over the previous year.

There were 34 deaths from tetanus, a decrease of three over the preceding year. This is the lowest number of deaths from this cause in ten years, except in the year 1908, when there were also 34 deaths. The majority of deaths from tetanus are in infants, and are due to infection of the umbilicus, the larger proportion occurring in the practice of midwives.

There were 183 deaths by suicide in the State in the last year. Of these, 43 were by poison, 28 by asphyxia, 20 by hanging and 65 by firearms. The greater number of suicides by poison and asphyxia occurred in Baltimore City, while the majority by firearms occurred in Rural Maryland. The total suicides in Rural Maryland were 82 and for Baltimore City 101.

The deaths from poisoning by food and other acute poisoning (not homicidal or suicidal) were 39; 26 in rural Maryland and 13 in Baltimore City.

Burns, scalds, etc., caused 155 deaths in 1915. Most of these deaths occurred in children and were due to clothes catching fire from stoves, open fireplaces or bonfires.

Accidental Drowning—199 deaths were reported from this cause during 1915. As usual, a majority of these deaths, 141 were reported from Rural Maryland, while approximately one-third, 58, were reported from Baltimore City.

Traumatism by Crushing (Vehicles, Railroads, Landslides, Etc.)—The number of deaths from this cause was 255 and has been on the increase for a number of years. The majority of these accidents occur on railroads. The proportion of deaths from automobile accidents is surprisingly small.

Lightning—There were nine deaths from this cause during 1915, all in Rural Maryland.

# Infant Mortality.

With a total number of 32,367 living births reported and a birth rate of 23.94 per thousand for Maryland, we feel as if we have a proper basis to compute a correct infant mortality rate.

Table 31 gives the infant mortality rate, white and colored for Rural Maryland, Baltimore City, and total Maryland for the year 1915. Birth registration in Maryland is now practically complete and these rates for ordinary purposes are considered correct. The mortality rate per 1,000 of living births for Baltimore City in the white population is 105.58 and in Rural Maryland is 97.58, while the rate in the colored population in Baltimore City is 184.25 and in Rural Maryland 202.05. The total rate for Rural Maryland is slightly higher than that for Baltimore City.

### TABLE XXXI.

INFANT MORTALITY RATE-1915-WHITE AND COLORED, MARYLAND.

	RURAL N	IARYLAND.	BALTIM	ORE CITY.	MARYLAND.		
•	Total	Mortality	Total	Mortality	Total	Mortality	
1	)eaths	Per 1,000		Per 1,000			
U	Inder	Living	Under	Living	Under	Living	
1	Year.	Births.	1 Year.	Births.	1 Year.	Births.	
White	1,431	97.58	1,210	105.58	2,641	101.09	
Colored	807	202.05	414	184.25	1,221	195.64	
						. ———	
Total	2,238	119.94	1.624	118.47	3.862	119.32	

### Duration of Life.

The main purpose of sanitary organization is to increase the duration of life. The duration of life is usually shown by tables of the mean duration and expectancy of life at given ages. Such tables were considered by Dr. Farr to represent the health and prosperity of the country, and to furnish a fair estimate of its living capital. Dr. Farr considered a table constructed in this way to be a measure of the life of the community, and hence it was called by him a biometer.

Other tables, popularly used by insurance companies, are tables of survivorship and tables showing the probability of living for a stated period beyond a given age—the probability of living one year being generally employed in the construction of this class of tables. Tables of duration of life are open to some objections for sanitary purposes. The extremes of either great or brief duration of life are largely lost, and certain components of the curve, which are of great sanitary importance, are not apparent.

Tables of duration of life are prognostic, the conditions of life in any given year being assumed to remain constant during a future period of years, i. e., 100 or 50 years. The particular question which interests the sanitarian is, at what period of life does the avoidable mortality occur. His problem is directed in general to increasing the duration of life, and the results of the work will be shown by fewer deaths in the early periods of life. The survivors, dying later, must, in proportion to their number, increase the number of deaths at an advanced age. Such tables and charts should illustrate, in the population of the State, under sanitary conditions present in the year of calculation, the number of the population living and the number dead at succeeding age periods.

The "Average Age of Death" may be considered in connection with the duration of life. This measurement is determined by adding the ages of the decedents and dividing them by the number of decedents. The fallacy that an increase in the average age at death indicates a corresponding increase in

the duration of life must be avoided. It is evident that in communities, made up of young adults and children, the average age at death will be low, while later, as the older portion of the population becomes numerous, the average age at death will be high, though there may have been no change in the death rate.

In 1915 there were 21,350 deaths in Maryland. Of these 21,269 occurred at known ages. Taking the sum of these ages, the 21,269 persons were found to have lived 868,616.932 years, giving an average age at death of 40.840 years. This number should be identical with the duration of life in a fixed population, therefore, the term "presumptive duration of life" may be applied to this factor in the succeeding table. The table of presumptive duration and presumptive expectancy is constructed as follows:

If we tabulate the decedents dying over the age of one year during 1915 we find 17,446 persons to have died, after living a total of 867,651.885 years, giving an average age at death of 49.734 years. This figure is entered in the table under the heading "Presumptive Duration — Years." Since this figure includes one year which has already been lived, the expectation of those decedent over one year will be, at the age of one year, 48.734 years. This factor appears in the table under the heading "Presumptive Expectancy at Age."

The factors of this calculation are shown in the first two columns of Table XXXII-A, while in the last two columns headed respectively "Presumptive Duration—Years" and "Presumptive Expectancy at Age," are given the presumptive expectation and duration of life by one-year periods.

To determine the total years lived by any number of decedents, it is necessary to add the ages of all the decedents and divide the sum of years by the number of decedents. This method involves considerable labor, and vital statisticians generally use the median age between the two age periods, multiplied by the number of decedents during the period.

By the aid of the Hollerith tabulating machine it has been possible for us to use the exact age of each individual of the 21,269 decedents of known ages in Maryland and secure a very accurate result. The ages being returned by years and months, the months are entered as decimal fractions of a year.

All registrations in which the number of days of the age of

the decedent was given were returned with an additional month, if over 15 days, and the additional days dropped, if under 15 days.

Thus the maximum error of any individual entry was only 0.0416 years. In any large series of entries the error should not exceed 0.001.

The following decimals were used for months, each being carried to the third place; one month, .083; two months, .166; three months, .249; four months, .333; five months, .416; six months, .499; seven months, .583; eight months, .686; nine months, .749; ten months, .833; eleven months, .916; twelve months, 1.000.

By reference to Table XXXII-A it will be seen that the presumptive duration of life steadily increases from the second year upwards, and the presumptive expectancy steadily diminishes.

The presumptive expectancy of life is greatest at two years, being greater than at birth (49.774, as compared with 48.734). At birth the expectancy and duration of life are identical, but after birth the expectancy increases to the second year, and thereafter steadily decreases, although the expectancy remains above that at birth up to the fifth year.

While the table of presumptive expectation and duration of life is not to be relied upon as an accurate measure of real expectation of life, such as is furnished by actuarial tables, its simplicity of construction and value for purposes of comparison give it a place in all our reports on vital statistics.

The figures over the age of 80 years are only included in the table for the interest they may possess, as they have no value for comparison, owing to the small number of entries and the great annual fluctuation in deaths over 80.

Table XXXII-B and XXXII-C are introduced in this report. Table XXXII-B gives the presumptive expectancy and duration of life worked on the total white deaths in Maryland during 1915, and Table, XXXII-C gives the presumptive expectancy and duration of life worked on the colored deaths in Maryland for the same year.

The presumptive expectation of life in the white population exceeds that of the colored population at the second year by 9.564 years; at the 20th year, it was 8.081 years greater; at the

30th year 6.106 years greater; at the 40th year it was 4.664 years greater; at the 50th year it was 2.746 years greater; at the 60th year it was only 0.401 years greater and at the 70th year the expectation of life in the colored population exceeds that of the white by 0.639 years.

# TABLE XXXII-A.

PRESUMPTIVE EXPECTANCY AND DURATION OF LIFE—MARYLAND—1915.

			Presumptive	Presumptive
Deaths Over		Total Years	Duration,	Expectancy
the Age of	Number.	Lived.	Years.	$at\ Age.$
1 year	17,446	867,651.885	49.734	48.734
2 years	16,740	866,692.196	51.774	49.774
3 years	16,472	866,052.391	52,577	49.577
4 years	16,312	865,517.568	53.060	49.060
5 years	16,191	864,990.322	53.424	48.424
6 years	16,097	864,485.899	53,705	47,705
7 years	16,034	864,084.703	53.891	46.891
8 years	15,958	863,526.265	54.112	46.112
9 years	15,883	862,896,780	54.328	45.328
10 years	15,816	862,265.379	54.519	44.519
11 years	15,754	861,623,578	54.692	43.692
12 years	15,698	860,984,217	54.847	42.847
13 years	15,647	860,356,782	54.985	41.985
14 years	15,594	859.645.680	55.127	41.127
15 years	15,509	858,425,785	55.350	40.350
16 years	15,420	857,051.126	55.580	39.580
	15,339	855,723.458	55.787	38.787
17 years	15,214	853,553.060	56.103	38.103
	15,098	851,419,106	56.393	37.393
19 years		848,862,444		36.719
20 years	14,966 14,832		56.719 57.048	36.048
21 years	,	846,141.873 843,239,184	57.0 <del>4</del> 8 57.379	35.379
22 years	14,696			
23 years	14,537	839,688.975	57.762	34.762
24 years	14,393	836,334.077	58.107	34.107
25 years	14,230	832,378.619	58.495	33.495
26 years	14,071	828,358.349	58.870	32.870
27 years	13,910	824,121.483	59.247	32.247
28 years	13,758	819,976.320	59.600	31.600
29 years	13,588	815,165.737	59.992	, 30.992
30 years	13,434	810,653,497	60.343	30.343
31 years	13,266	805,568.382	60.724	29.724
32 years	13,099	800,335.753	61.099	29.099
33 years	12,934	795,010.151	61.467	28.467
34 years	12,793	790,313.151	61.777	27.777
35 years	12,618	784,309.763	62.158	27.158
36 years	12,407	776,875.831	62.616	26.616
37 years	12,257	771,434.667	62.938	25.938
38 years	12,076	764,676,995	63.322	25.322
39 years	11,875	756,982,304	63,746	24.746
40 years	11,688	749,627.011	64.136	24.136
41 years	11,461	740,496.980	. 64.610	23.610
42 years	11,301	733,889.425	64.940	22.940
43 years	11,094	725,136.571	65.363	22.363
44 years	10,930	718,041.464	65.695	21.695
45 years	10,744	709,798.862	66.065	21.065
46 years	10,476	697,678.285	66.598	20.598
47 years	10,300	689,532,006	66.945	19.945
48 years	10,117	680,876,710	67.300	19.300
49 years	9,918	671,274.450	67.682	18.682
50 years	9,714	661,221.010	68.069	18.069
51 years	9,410	645,958,964	68.646	17.646
52 years	9,222	636,321.630	69.000	17.000
53 years	8,953	622,268.394	69.504	16.504

			Presumptive	Presumptive
Deaths Over		Total Years	Duration,	Expectancy
the Age of N	umber.	Lived.	Years.	$at\ Age.$
54 years	8,740	610,919,169	69,899	15.899
55 years	8,505	598,152.604	70.330	15.330
56 years	8.227	582,794.054	70.839	14.839
57 years	7,960	567,758.695	71.326	14.326
58 years	7,706	553,194.679	71.788	13.788
59 years	7,432	537,213.701	72.284	13.284
60 years	7,188	522,727.425	72.722	12,722
61 years	6,823	500,739.026	73,390	12.390
62 years	6,604	487,303.975	73.789	11.789
63 years	6,317	469,429,572	74.312	11.312
64 years	6.041	451,959,636	74.815	10.815
65 years	5,761	433,944.482	75.325	10.325
66 years	5,391	409,805.850	76.017	10.017
67 years	5,134	392,759.033	76.502	9.502
68 years	4.861	374,381.133	77.017	9.017
69 years	4,565	354,159.641	77.582	8.582
70 years	4,302	335,926.558	78.086	8.086
71 years	3,924	309,372.424	78.841	7.841
72 years	3,658	290,402.219	79.388	7.388
73 years	3,362	268,989.563	80.009	7.009
74 years	3,047	245,899.442	80.702	6.702
75 years	2,799	$227,\!454.808$	81.263	6.263
76 years	2,445	200.809.638	82.131	6.131
77 years	2,166	179,512.629	82.877	5.877
78 years	1,943	162,256.890	83.508	5.508
79 years	1,679	141,580.159	84.324	5.324
80 years	1,491	126,660.910	84.950	4.950
81 years	1,224	105,229.829	85.972	4.972
82 years	1,059	91,810.106	86.695	4.695
83 years	886	77,558.576	87.538	4.538
S4 years	758	66,893.947	88.251	4.251
85 years	619	55,169.431	89.127	4.127
86 years	498	44,852,922	90.066	4.066
87 years	408	37,082.781	90.889	3.889
88 years	324	29,746.779	91.811	3.811
89 years	269	24,880.251	92.492	3.492
90 years	217	20,235.107	93.249	$\frac{3.249}{3.429}$
91 years	156	14,730.982	94.429	3.444 3.444
92 years	118 91	11,262,376	$95.444 \\ 96.364$	3.364
93 years	61	$8,769.117$ $5,97\overline{1}.337$	90.304	3.891
94 years 95 years	49	3,911.551 4,840.488	98.785	3.785
95 years 96 years	38	3,794,407	99,853	3.853
97 years	30	3,025,159	100,839	3.839
98 years	$\frac{30}{24}$	2,440.996	101.708	3.708
99 years	14	1,458.916	104.208	5.208
100 years	13	1,359,916	104.609	4.609
100 years	10	1,059.833	105.983	4.983
102 years	9	958.000	106,444	4.444
105 years	8	856,000	107.000	2.000
106 years	$\ddot{6}$	646.000	107.667	1.667
110 years	$\check{2}$	222,000	111.000	1.000
112 years	1	112.000	112.000	0.000
Total	21,269	868,616.932	40.840	

# TABLE XXXII-B.

# PRESUMPTIVE EXPECTANCY AND DURATION OF LIFE-WHITE-1915.

			Presumptive	Presumptive
Deaths Over		Total Years	Duration,	Expectancy
the Age of Num	ber.	Lived.	Years.	$at\ Age.$
1 year 13,1	15	684,450.350	52.188	51.188
2 years 12,6		683,795.634	54,119	52.119
3 years 12,4		683,330.895	54.926	51.926
4 years 12,5		682,938.823	55.411	51.411
5 years 12,2	245	682,585.082	55.744	50.744
6 years 12,1		682,238.478	56.008	50.008
7 years 12,1		681,957.024	56.188	49.188
-8 years 12,0		681,564.931	56.402	48.402
9 years 12,0		681,158.768	56.593	47.593
10 years 11,9		680,697.148	56.786	46,786
11 years 11,9		680,293.377	56.938	45.938
12 years 11,9		679,894.708	57.072	45.072
13 years 11,8		679,463.101	57.203	44,203
14 years 11,8		679,006.243	57.329	43,329
15 years 11,7		678,362,098	57.493	42,493
16 years 11,7		677,632.897	57.661	41.661
17 years 11,7		677,042.541	57.788	40,788
18 years 11,6		675,750.668	58.044	40.044
19 years 11,5		674,330.800	58.308	39.308
20 years 11,4		672,643.157	58.603	38.603
21 years 11,5		670,953,676	58.881	37.881
		669,222.573	59.150	37.150
,		667,250.897	59.438	36.438
		665,128.275	59.733	35.733
		662,623,990	60.064	35.064
26 years 10,9 27 years 10,8		$659,940.017 \\ 657,090.114$	$60.401 \\ 60.740$	$34.401 \\ 33.740$
		654,247.168	61.065	33.065
		651,331.514	61.383	32.383
			61.669	31.669
30 years 10,5 31 years 10,4		648,577.309 $645,273.840$	_ 61.998	30.998
		641,636.836	62.343	30.343
		638,144.910	62.662	29.662
		634,842.335	62.949	28,949
	)54	630,342.643	63.326	28.326
		625,787.342	63.693	27.693
	325 717	621,866.354	63.998	26.998
	585	616,930.773	64.364	26.364
	156	611,982,544	64.719	25.719
	323	606,743.925	65.080	25.080
	182	601,064.938	65.461	24.461
	)69	596,391,205	65.762	23.762
	)35 	590,718.661	66.113	23.113
	,55 309	585,260.821	66.439	22,439
	573	579,228.036	66.785	21,785
	507	571,708.690	67.205	21.205
	876	565,639.564	67.531	20.531
	236	559,009.615	67.874	19.874
	.30 )96	552.246.699	68.212	19.212
	)46	544,846.332	68.569	18.569
	48	534,893.709	69.036	18.036
	619	528,272.571	69.336	17.336
	132	518,490.595	69.765	16.765
	275	510,119.730	70.120	16.120
or years 1,		910,110.100	10.120	10.120

			Presumptive	Presumptive
Deaths Over		Total Years	Duration,	Expectancy
the Age of	Number.	$. \hspace{1.5cm} Lived.$	Years.	at Age.
55 years	7.096	500,383,905	70.516	15.516
56 years	6,889	488,934,619	70.973	14.973
57 years	6,682	477,269.455	71.426	14.426
58 years	6,478	465,560.117	71.868	13.868
59 years	6,256	452,604.566	72.347	13.347
	6,051	440,426.157	72.786	12.786
60 years	5.782	424,205.354	73,367	12.367
61 years	5,182 $5,592$	412,544.880	73.774	11.774
· ·	5,358	397,961.490	74.274	11.274
63 years	,	382,955.733	74.781	10.781
64 years	$5,121 \\ 4.880$	367,442.609	75.296	10.131
65 years	. ,		75.923	9.923
66 years	4,592	348,640.047		9.426
67 years	4,363	333,414.580	76.426	
68 years	4,125	317,416.508	76.949	8.949
69 years	3,878	300,530.698	77.496	8.496
70 years	3,656	285,132,148	77.990	7.990
71 years	3,363	264,532.360	78.660	7.660
72 years	3,124	247,483.149	79.220	7.220
73 years	2,869	229,024.905	79.827	6.827
74 years	2,599	209,224.796	80.502	6.502
75 years	2,383	193,156.232	81.056	6.056
76 years	2,092	$171,\!238.141$	81.854	5.854
77 years	1,842	152,151.351	82.601	5.601
78 years	1,637	136,285.525	83.253	5.253
79 years	1,407	118,265.807	84.055	5.055
80 years	1,230	104,218.305	84.730	4.730
81 years	1,023	87,590.822	85.622	4.622
82 years	876	75,631.512	86.337	4.337
83 years	721	62,859.895	87.184	4.184
84 years	604	53,110.764	87.932	3.932
85 years	488	43,321.744	88.774	3.774
86 years	396	35,472,982	89.578	3.578
87 years	316	28,564.276	90.393	3.393
88 years	245	22,360.938	91.269	3.269
89. years	192	17,670.493	92.034	3.034
90 years	146	13,560.514	92.880	2.880
91 years	109	10,218.469	93,747	2.747
92 years	78	7.388.110	94.719	2.719
93 years	58	5,539.870	95.515	2.515
94 years	36	3,486,839	96.857	2.857
95 years	24	2,355.990	98.166	3.166
96 years	18	1,785,158	99.175	3.175
97 years	15	1,496.159	99.744	2.744
98 years	10	1,008,996	100.900	2.900
100 years	4	418.916	104.729	4.729
101 years	3	318.S33	106.278	5.278
105 years	$\frac{3}{2}$	217.000	108.500	3.500
112 years	ĩ	112.000	112.000	0.000
ira years		112,000		0.000
Total	15,731	685,096.801	43.551	

# TABLE XXXII-C.

# PRESUMPTIVE EXPECTANCY AND DURATION OF LIFE—COLORED—1915.

			Presumptive	Presumptive
Deaths Over		Total Years	Duration,	Expectancy
the Age of	Number.	Lived.	Years.	at Age.
1 year	4,331	183,201.535	42.300	41.300
2 years	4,105	182,896,562	44.555	42.555
3 years	4,031	182,721.496	45.329	42.329
4 years	3,987	182,578.745	45.794	41.794
5 years	3,946	182,405,240		
6 years	3,916	182,247.421	46.225	$\frac{41.225}{40.539}$
7 years	3,897	182,127.679	46.539	
8 years	3,874	181,961.334	46.735	39.735
9 years	3,847	181,738.012	$46.970 \\ 47.241$	$\frac{38.970}{38.241}$
10, years	3,829	181,568.231	47.419	
11 years	3,806			37.419
.12 years	3,785	181,330.201	47.643	36.643
13 years	3,769	181,089.509 180,893.681	47.844	35.844
14 years	3,750	180,639.437	/ 47.995 48.171	34.995
15 years	3,710		48.535	34.171
	3,668	180,063.687 179,418.229		33.535
16 years	3,623	178,680.917	48.914 $49.318$	32.914
18 years	3,572	177,802,392	49.777	32.318
19 years	3,533	177,088.306	50.124	31.777 $31.124$
20 years	3,488	176,219.287	50.522	30.522
21 years	3,437	175,188,197	50.971	29.971
22 years	3,382	174,016.611	51.454	29.454
23 years	3,311	172,438.078	52.080	29.080
24 years	3,258	171,205.802	52.549	28.549
25 years	3,198	169,754.629	53.081	28.081
26 years	3,145	168,418.332	53.551	27.551
27 years	3,092	167,031.369	54.020	27.020
28 years	3.044	165,729.152	54.445	26.445
29 years	2,977	163,834.223	55.033	26.033
30 years	2,917	162,076.188	55.563	25.563
31 years	2,858	160,294.542	56.086	25.086
32 years	2,807	158,698.917	56.537	24.537
33 years	2,750	156,865.241	57.042	24.042
34 years	2,708	155,470.816	57.412	23.412
35 years	2,664	153,967,120	57.795	22.795
36 years	2,582	151,088.489	58.516	22.516
37 years	2,540	149,568.313	58.885	21.885
38 years	2,491	147,746.222	59.312	21.312
39 years	2,419	144,999.760	59.942	20.942
40 years	2,365	142,883.086	60.416	20.416
41 years	2,279	139,432.042	61.181	20.181
42 years	2,232	137,498.220	61.603	19.603
43 years	2,159	134,417.910	62.259	19.259
44 years	2,121	132,780.643	62.603	18.603
45 years	2,071	130,570.826	63.047	18.047
46 years	1,969	125,969,595	63.976	17.976
47 years	1,924	123,892.442	64.393	17.393
48 years	1,881	121,867.095	64.788	16.788
49 years	1,822	119,027.751	65.328	16.328
50 years	1,768	116,374.678	65.823	15.823
51 years	1,662	111,065.255	66.826	15.826
52 years	1,603	108,049.059	67.404	15,404
53 years	1,521	103,777.799	68.230	15.230

### TABLE XXXII-C-Continued.

					Presumptive	Presumptive
Dear	ths Over			Total Years	Duration,	Expectancy
the	Age of	Number.		Lived.	Years.	at Age.
54	years	1,465		100,799.439	68,805	14.805
	years			97,768.699	69,389	14.389
	years			93,859.435	70.149	14.149
	years			90,489,240	70.805	13,805
	years	,		87,634.562	71.364	13.364
	years			84,609.135	71.947	12.947
	years			82,301,268	72.385	12.385
	years			76,533,672	73.519	12.519
	years			74,759.095	73.873	11.873
				,	74.524	11.524
	years			71,468.082		11.004
	years			69,003.903	75.004	10.485
	years			66,501.873	75.485	
	years			61,165.803	76.553	10.553
01	years	771		59,314.453	76.932	9.932
	years			56,964.625	77.398	9.398
	years			53,628.943	78.063	9.063
	years			50,794.410	78.629	8.629
	years	561		44,840.064	79.929	8.929
	years			42,919.070	80.373	8.373
	years			39,964.658	81.064	8.064
	years			36,674.646	81.863	7.863
	years	416		$34,\!298.576$	82.449	7.449
76	years			$29,\!571.497$	83.772	7.772
	years	324		27,361.278	84.448	7.448
	years			25,971.365	84.874	6.874
	years	272		23,314.352	85.715	6.715
80 ;	years	. 261		22,442.605	85.987	5.987
	years			17,639.007	87.756	6.756
82	years	183		16,178.594	88.408	6.408
00 , 04 .	years	165		14,698.681	89.083	6.083
	years years	$154 \\ 131$		13,783.183 $11,847.687$	89.501 90.440	5.501 5.440
	years	$\frac{131}{102}$		9,379.940	91.960	5.960
	years	92		8,518.505	92.592	5.592
,	years;	79		7,385.841	93.492	5,492
	years	77		7,209.758	93.633	4.633
90	years	71		6,674.593	94.008	4.008
91	vears	47		4,512.513	96.011	5.011
	vears	40		3,874.266	96.857	4.857
93	years	- 33		3,229.247	97.856	4.856
94;	years	25		2,484.498	99.380	5.380
	years	25		2,484.498	99.380	4.380
~ -	years	20		2.059.249	100.462	4.462
	years	15		1,529.000	101.933	4.933
	years	14		1,432.000	102.286	4.286
	years	10		1,040.000	104.000	5.000
	years	$\frac{9}{7}$		941.000	104.555	4.555
	years	6		$741.000 \\ 639.000$	105.857	3.857
	years	5		534.000	106.500 106.800	1.500 0.800
	years years	1	~	110.000	110.000	0.000
~ <b>_</b> ,	J CALS	1		110.000	110.000	0.000
To	tal	5,538	`	183,520.131	33.138	




Chart 20-A—Presumptive Expectation and Duration of Life—Total Maryland, 1915.




Chart 20-B—Presumptive Expectation and Duration of Life—Total Maryland, 1915.




Chart 20-C—Presumptive Expectation and Duration of Life—Total Maryland, 1915.

The succeeding tables (Table XXXIII and Table XXXIV) give the average age at death by counties and by months, respectively.

The county giving the lowest average age at death in 1915 was St. Mary's County (32.461); the highest average age at death was in Carroll County (50.536).

The next table (Table XXXIV) gives the average age at death by months.

The lowest average age at death is in August. The average age at death in this month is 34.882. It is evident that in this State with the high infant mortality in July, August and September the lowest average age at death will always fall in one of these three months.

By reference to the table it will be seen that the average age at death falls below the mean in the months of July, August, September and October. In all the other months the figures are above the mean.

#### TABLE XXXIII.

#### MEAN AGE AT DEATH BY COUNTIES-1915.

•	Total	$Total\ Age$	Average Age
Counties.	Deaths.	of Decedents.	at Death.
Allegany	822	34,663.716	42.170
Anne Arundel	736	26,583,747	36.119
Baltimore	2,687	115,908,523	43.137
Calvert	150	5,107.603	34.051
Caroline	278	10,250,735	36.873
Carroll	565	28,552,827	50.536
Cecil	365	17,238,452	47.229
Charles	267	9,189.813	34.419
Dorchester	497	16,506,325	33.212
Frederick	776	35,898.156	46.261
Garrett	224	8,614,946	38.460
Harford	386	17,595,121	45.583
Howard	199	8.907.712	44.762
Kent	272	10.633.874	39.095
Montgomery	436	19,230.802	44.107
Prince George's	518	20,330,444	39.248
Queen Anne's	272	10,407,071	38.261
Somerset	398	14,325.704	35,994
St. Mary's	266	8.634.587	32.461
Talbot	310	12,922,170	41.684
Washington	745	31,894,274	42.811
Wiconico	422	16,329,559	38.696
Worcester	351	13,932,291	39,693
Total Counties	11,942	493.658.452	41.338′
Baltimore City	9,327	374,958.480	40.201
Maryland		868,616.932	40.840
•			

TABLE XXXIV.

MEAN AGE AT DEATH BY MONTHS-1915.

RURAL MARYLAND.  Average Total Age of Age at	ND. Average	Total Value	<del></del>	=	Average	Potal	MARYLAND.	1verage
Deaths.	Decedents.	age ut Death.	Deaths.	Decedents.	Death.	Deaths.	Decedents.	Aye ar Death.
	41,895.096	43.325			44.368	1,816	79,563.837	43.813
	40,505.479	43.045			41.337	1.680	71,053.817	45.294
	53,789.070 \	45.584			42.956	2,132	94,683.523	44.411
	51,806.895	44.584			43.221	2,061	90,662.677	43.990
	2,216.549	44.863			40.364	1,705	73,054.926	42.847
	33,571.329	41.755			38.100	1,437	57,688.440	40.145
	38,259.432	37.657			35.100	1.780	65,075.620	36.559
	37,671.151	35.439			34.082	1,803	62,891.615	34.882
	34.368.462	35.875			35.206	1,638	58.308.820	35.598
	37,818.803	38.749			39.248	1,697	66,116.771	38.961
••	34,790.531	40.313			42.043	1,618	66,533.308	41,121
	46,965.655	43.852			43.343	1,902	82,983.578	43.630
4	93,658.452	41.338	9.327		40.201	21,269	868,616.932	40.840

#### Maryland Biometer.

This biometer is constructed on a table of survivorship, in which 10,000 persons born in a given year are traced throughout life, under the sanitary conditions of the year of computation, as indicated by the mortality returns.

If the death rates for the several age periods are determined for any given year, we may, by applying these rates to the estimated population of this year, construct a table of survivorship, showing the number surviving at certain periods thereafter among those born in the given year.

Thus, of 10,000 persons born in Maryland during 1915, we have to determine the number surviving at the end of five years, of ten years, etc., providing the death rate of 1915 remains constant. A chart constructed from this table will indicate the sanitary condition of the State during the year 1915, and, described in the manner presently to be mentioned, forms the "Maryland Biometer."

The Maryland biometer is shown in the chart in quinquennial periods for the year 1915, assuming the death rate of that year to remain constant. Instead of considering the whole population as a basis the scale is reduced to a population of 10,000 for the convenience of comparison with succeeding years. Of a population of 10,000 born in Maryland during 1915, how many will be living and how many dead at the end of five years, of ten years, etc.? This table and chart indicate survivorship and give both the number of living and dead at each quinquennial period after 1915.

Such charts readily admit of comparison with preceding or following years, as variations in the mortality at the various ages appear in the curve in their proper positions and do not (as in the expectancy tables) merely modify the form of the curve.

The only factors necessary in the construction of this curve are the mortality rates for the several age periods (0.5, 5.10, etc.), which may be applied first to the original population of 10,000, then to the remaining population, after deducting the deaths from 0.5, etc. Applying the death rates obtained from Table XIX-A to 10,000 persons born in 1915, the survivorship at succeeding quinquennial periods is shown in Table XXXV-A. In Table XXXV-B the survivorship with regard to color is shown.

#### TABLE XXXV-A.

#### SURVIVORSHIP IN MARYLAND, 1915, TOTAL POPULATION.

Survivors of 10,000 Persons Born in Maryland in 1915 at Succeeding Quinquennial Periods, Assuming the Death Rate of That Year to Be Constant.

Number born in	1915	10,000
Number reaching	age of 5 years (A. D. 1920)	8,228
Number reaching	age of 10 years (A. D. 1925)	8,118
Number reaching	age of 15 years (A. D. 1930)	8,025
Number reaching	age of 20 years (A. D. 1935)	7,862
Number reaching	age of 25 years (A. D. 1940)	7,636
Number reaching	age of 30 years (A. D. 1945)	7,371
Number reaching	age of 35 years (A. D. 1950)	7,070
Number reaching	age of 40 years (A. D. 1955)	6,729
Number reaching	age of 45 years (A. D. 1960)	6,343
Number reaching	age of 50 years (A. D. 1965)	5,881
Number reaching	age of 55 years (A. D. 1970)	5,305
Number reaching	age of 60 years (A. D. 1975)	4,530
Number reaching	age of 65 years (A. D. 1980)	3,640
Number reaching	age of 70 years (A. D. 1985)	2,681
Number reaching	age of 75 years (A. D. 1990)	1,593
Number reaching	age of 80 years (A. D. 1995)	583

#### TABLE XXXV-B.

COMPARATIVE TABLE OF SURVIVORSHIP IN MARYLAND, 1915, WHITE AND COLORED.

Number of Survivors of 10,000 Persons Born in Maryland in 1915 at Succeeding Quinquennial Periods, Assuming the Death Rates of That Year to Remain Constant.

	White.	Colored.
Number born in 1915	10,000	10,000
Number reaching age of 5 years (A. D. 1920)	8,517	6,908
Number reaching age of 10 years (A. D. 1925)	8,421	6,753
Number reaching age of 15 years (A. D. 1930)	8,350	6,584
Number reaching age of 20 years (A. D. 1935)	8,227	6,272
Number reaching age of 25 years (A. D. 1940)	8,052	5,883
Number reaching age of 30 years (A. D. 1945)	7,831	5,490
Number reaching age of 35 years (A. D. 1950)	7,567	5,070
Number reaching age of 40 years (A. D. 1955)	7,267	4,614
Number reaching age of 45 years (A. D. 1960)	6,924	4,110
Number reaching age of 50 years (A. D. 1965)	6,499	3,567
Number reaching age of 55 years (A. D. 1970)	5,970	2,887
Number reaching age of 60 years (A. D. 1975)	5,172	2,239
Number reaching age of 65 years (A. D. 1980)	4,199	1,685
Number reaching age of 70 years (A. D. 1985)	3,128	1,153
Number reaching age of 75 years (A. D. 1990)	1,881	631
Number reaching age of 80 years (A. D. 1995)	691	227

This biometer is graphically shown in Chart No. 21. It is evident that such a chart will indicate the life condition of the State, both in its form and in its area. To calculate the area it is only necessary to measure the length of the perpendiculars to the base line from each component point of the curve, and take the same measurement for the next succeeding point. The area of each trapezoid thus produced is calculated in the usual manner, by taking the product of one-half the sum of its parallel sides by its altitude.




CHART 21.




CHART 22—Illustrating the higher death rate in the colored population. In the upper biometer the total white living comprises the area marked colored living, plus the area marked white living. In the lower biometer the colored dead comprises the area marked white dead, plus the area marked colored dead.

#### TABLE XXXVI-A.

#### ESTIMATED POPULATIONS OF MARYLAND.

	1910.	1911.	1912.	1913.	1914.	1915.
0- 5 years	137,761	138,900	140,039	141,178	142.317	143,457
5-10 years	133,658	134,756	135,854	136,952	138,049	139,148
10-15 years	129,679	130,750	131,820	132,890	133,961	135,032
15-20 years	128,173	129,242	130,311	131,379	132,447	133,517
20–25 years	123,377	124,393	125,410	126,425	127,441	128,457
25-30 years	110,215	111,123	112,030	112,938	113,845	114,752
30–35 years	95,997	96,807	97,618	98,428	99,239	100,049
35–40 years	92,400	93,172	93,944	94,717	95,489	96,262
40-45 years	78,763	79,428	80,094	80,760	81,426	82,091
45-50 years	67,982	68,560	69,136	69,713	70,290	70,867
50-55 years	59,179	59,689	60,198	60,707	61,216	61,725
55-60 years	43,203	43,583	43,964	44,344	44,725	45,106
60-65 years	34,984	35,289	35,596	35,902	36,207	36,514
65-70 years	26,582	26,816	27,051	27,285	27,519	27,752
70–75 years	17,659	17,814	17,968	18,124	18,280	18,435
75–80 years	9,836	9,925	10,014	10,102	10,192	10,280
80 years and over	6,821	6,880	6,938	6,998	7,057	7,115
Unknown	1,341	1,349	1,357	1,366	1,374	1,381
Total	1,297,610	1,308,476	1,319,342	1,330,208	1,341,074	1,351,940

#### TABLE XXXVI-B.

#### DEATHS BY AGES.

	1910.	1911.	1912.	1913.	1914.	1915.
0- 5 years	5,873	5,467	5,415	5,896	5,419	5.084
5-10 years	421	426	373	436	397	371
10–15 years	372	320	272	289	299	308
15-20 years	606	574	556	591	573	544
20–25 years)	1,656	763	742	793	803	737
25–30 years}	1,000	850	843 ′	845	814	795
30-35 years	1,725	774	826	805	757	817
35-40 years}	1,120	857	- 885	818	912	930
40–45 years	1,764	842	904	892	1,006	942
45–50 years}	1,10%	921	940	964	1,029	1,032
50-55 years	2,120	1,077	1,127	1,101	1,156	1,210
55-60 years	2,120	1,017	1,056	1,095	1,250	1,317
60-65 years	2,640	1,215	1,276	1,186	1,322	1,435
65-70 years	2,040	1,285	1,377	1,293	1,403	1,462
70–75 years	2,401	1,325	1,370	1,413	1,483	1,497
75–80 years}	,	1,105	1,106	1,142	1,211	1,303
80 years and over	1,359	1,414	1,458	1,420	1,467	1,485
Unknown	34	58	71	69	82	81
Total	20,971	20,290	20,597	21,048	21,383	21,350

TABLE XXXVI-C.

#### DEATH RATE PER THOUSAND AT AGE PERIODS.

	1910.	1911.	1912.	1913.	1914.	1915.
0- 5 years	42.63	39.36	38.67	41.76	38.08	35.44
5–10 years	3.15	3.16	2.75	3.18	-2.88	2.67
10-15 years	2.87	2.45	2.06	2.17	2.23	2.28
15–20 years	4.73	4.44	4.27	4.50	4.33	4.07
20-25 years)	7.09	6.13	5.92	6.27	6.30	5.74
25-30 years	1.00	7.65	7.52	7.48	7.15	6.93
30-35 years (	9.16	8.00	8.46	8.18	7.63	8.17
35-40 years	9.10	9.20	9.42	8.64	9.55	9.66
40-45 years)	12.02	10.60	11.29	11.05	12.35	11.47
45–50 years	12.02	13.43	13.60	13.88	14.64	14.56
50-55 years	20.71	18.04	18.72	18.14	18.88	19.60
55-60 years	20.11	23.33	24.02	24.69	27.95	29.20
60-65 years)	42.88	34.43	35.85	-33.03	36.51	39.30
65-70 years	12.00	47.92	47.40	47.39	50.98	52.68
70-75 years)	87.33	74.38	76.25	77.96	81.13	81.20
75-80 years	01.00	111.33	110.45	113.05	118.82	126.75
80 years and over	199.24	205.52	210.15	202.92	207.88	208.71
Unknown	25.35	42.99	52.32	50.51	59.68	58.65
Total	16.16	15.51	15.61	15.82	15.94	15.79

#### TABLE XXXVII.

#### Survivorship by Ages—1910, 1911, 1912, 1913, 1914, 1915.

10,000 Born in	1910.	1911.	1912.	1913.	1914.	1915.
Number reaching 5 years	7.868	8,032	8.066	7.912	8,096	8,228
Number reaching 10 years	7,744	7,905	7,955	7,786	7,979	8,118
Number reaching 15 years	7,633	7,808	7,873	7,702	7,890	8,025
Number reaching 20 years	$\sqrt{7,452}$	7,635	7,705	7,529	7,719	7,862
Number reaching 25 years	6,924	7,401	7,477	7,293	7,476	7,636
Number reaching 30 years	5 0,524	7,118	7,196	7,020	7,209	7,371
Number reaching 35 years	$\{-6,290-$	6,833	6,892	6,733	6,934	7.070
Number reaching 40 years	5 0,290	$6,\!519$	6,567	6,442	6,603	6,729
Number reaching 45 years	5,534	6,173	$6,\!196$	6,086	6,195	6,343
Number reaching 50 years	3,004	5,758	5,775	5,665	5,742	5,881
Number reaching 55 years	$\frac{1}{4.388}$	5,239	5,234	5,151	5,200	5,305
Number reaching 60 years	4,000	4,628	4,605	4,515	4,473	4,530
Number reaching 65 years	2,506	3,831	3,780	3,769	3,656	3,640
Number reaching 70 years		2,913	2,884	2,876	2,724	2,681
Number reaching 75 years	318	1,830	1,784	1,755	1,619	1,593
Number reaching 80 years	( ,,10	811	799	763	657	583

Table XXXVIII gives the marriages in Baltimore and in the various counties by months. Of the 18,616 marriage returns, 6,202 were made to the clerks of the court of Baltimore City and returns for 12,414 were made to the clerks of the courts of the counties.

Table XXXIX gives the divorces for Baltimore City and the several counties by months. Out of the total of 822 divorces recorded in the State during the time covered by these returns, 554 were issued in the courts of Baltimore City.

TABLE XXXVIII.

MARRIAGES FOR MARYLAND-1915.

Total.	2,191	296 296	743	8.	169	256	2,743	105	260 2	505	738	197	723	175	504	179	147	200	116	177	1,320	313	219		12,414	6,202	18 616	10,010
Dec.	505	ਜ਼ ਜ਼	57	σo	41	334	245	19	41	85.	76	17	99	21	49	16	98 98	41 \$	17	88	166	<u> 20</u>	47		1,403	280	9 033	4,000
Nov.	211	95 65	0,	11	133	35	278	13	35	37	5	56	528	19	36	<u>8</u> ]	16	23	11	19	131	65 65	15	1	1,185	662	1 6/7	1,041
Oet.	191	31	2	10	1	31	268	<u>.</u>	07 07	<del>1</del>	55 SS	83	<u>6</u>	14	46	18	15	<u>6</u>	14	16	163	댦	20		1,189	204	1 700	1,100
Sept.	199	30	80	x	11	17	278	G:	17	33	83	17	3	18	20	18	Ξ	11	t-	Ħ	105 201	56	13		1,126	240	1 666	1,000
Aug.	211	16	50	C)	13	21	227	4	13	뜮	19	L-	83	10	4	15	10	25 25	ಣ	G	86	27	19		186	443	1 497	1,474
July.	177	021	52	<u>L-</u>	6	12	247	9	16	36	87	7	19	13	53	œ	<u>-</u>	20	4	G	76	<u>e.</u>	11			423	7	1,414
June.	203	65	8	ţ~	T.	13	277	10	19	48	74	24 47	69	18	33	13	15	33	10	20	103	28 87	18		1,190	90%	98	1,930
May.	164	91 85	75	ဗ	4	51	507	9	18	40	#	13	<u></u>	9	65 65	14	6	15	73	6	80 83	5	18		867	398	1 905	1,205
Apr.	158	119	S.	ဗ	1	7	506	#	16	38	<u>85</u>	16	55 Se	2	38	읞	L—	17	10	15	102	19	16		951	527	1 470	1,473
Mar.	117	19	43	10	<u>-</u>	16	154	4	21	40	52	5	48	10	83	9	S	18	ဗ	Π	88	15	17	] ;	747	352	60	1,034
Feb.	148	61	7	4	133	81	185	10	£1	50	23	. 14	20	Ħ	56	10	14	16	t-	l-	91	23	13		835	406	1 0 11	1,241
Jan.	150	57	4 55	23	19	61 80	177	oo	89	33	41	12	#	82	36	6	. 15	17	<u>8</u> 1	16	66	23	15		#06	474	016	T,513
	Allegany	Anne Armidel	Baltimore	Calvert	Caroline	Carroll	Cecil	Charles	Dorchester	Frederick	Garrett	Harford	Howard	Kent	Montgomery	Prince George's	Oueen Anne's	Somerset	St. Mary's.	Talbot	Washington	Wicomico	Woreester		Total Counties	Baltimore City	mater are males d	Total Maryland

# TABLE XXXIX.

DIVORCES FOR MARYLAND—1915.

- 101	7778
	ī
	9.
	3
жере то по	<u>.</u>
8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	<u>e</u>
	古
\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	33
######################################	62
4 7. 8. 4. 4. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0.	<u></u> 23
м в в в в в в в в в в в в в в в в в в в	14
	19
# m H at 0 at m 0 0 0 0 0 0 0 at 1 at 2	<del>\$</del>
Allegany Anne Arundel Baltimore Calvert Caroline Carroll Cecil Charles Dorchester Frederick Garrett Harford Howard Kont Montgomery Prince George's Queen Anne's. Somerset St. Mary's. Talbot Washington Wicomico Worcester Total Counties.	Total Maryland

#### Prosecutions.

W. M. Talbott, West River, Anne Arundel County, for interring the body of Robert T. Neal in Anne Arundel County, without first obtaining a burial permit. Hearing waived, jury trial prayed. The State's Attorney believed the action of the defendant justifiable and recommended that warrant be quashed.

Sol Levinson, 1107 E. Baltimore St., Baltimore, for refusal or neglect to properly obtain a burial permit before removing the body of Samuel Sanberg, who died November 9, 1914, in Baltimore County. Defendant fined \$10.00 and costs.

Sol Levinson, 1107 E. Baltimore St., Baltimore, for refusal or neglect to properly obtain a buirial permit before removing the body of Rachel Lazarus, who died January 16, 1915, in Baltimore County. Case dismissed by magistrate since defendant was authorized by State's Attorney to conduct interment and showed good faith on his part.

Dr. H. K. Peltekian, Preston and Eutaw Sts., Baltimore, for refusal or neglect to report the birth of the infant of Mrs. Walter Hermann, who was born February 22, 1914, in Baltimore County. Defendant plead guilty and was fined \$5.00 and costs.

R. N. Trabert, Freeland, Baltimore County, for interring the body of James White, in Baltimore County, without first obtaining a burial permit. Defendant plead guilty and was fined \$5.00 and costs.

M1s. Theresa Blankey, 535 S. Belnord St., Baltimore, for refusal or neglect to report the birth of the infant of Mrs. John J. Kerns, who was born January 10, 1915. Defendant plead guilty and was fined \$5.00 and costs.

Dr. James Craighill, The Walbert, Baltimore, for refusal or neglect to report the birth of the infant of Mrs. John J. Kerns, who was born January 10, 1915. Defendant plead guilty and was fined \$5.00 and costs.

Dr. George De Hoff, 2020 N. Charles St., Baltimore, for refusal or neglect to report the birth of the infant of Mrs. William Brown, who was born May 25, 1915, in Baltimore. Case dismissed upon recommendation of State. Certificate dated wrong and was reported by Dr. De Hoff's assistant.

Dr. George De Hoff, 2020 N. Charles St., Baltimore, for refusal or neglect to report the birth of the infant of Mrs. Henry Wildberger, who was born September 20, 1914, in Baltimore. Defendant plead guilty and was fined \$5.00 and costs.

Mrs. Yetta Feldman, Exeter St., Baltimore, for refusal or neglect to report to the Health Officer, or physician, a case of red, sore or swollen eyes in the infant of Mrs. Dora Cushner, in Baltimore, May 24, 1915. Defendant plead not guilty, but was fined \$5.00 and costs.

Dr. Raymond Glann, Mount Winans, Baltimore County, for refusal or neglect to report the birth of the infant of Mrs. Bernard C. Fink, who was born November 2, 1914, in Baltimore. Defendant plead not guilty but was fined \$5.00 and costs.

Mrs. Marie Kell, 1026 Denver St., Baltimore, for attending Mrs. John Monroe at childbirth, September 14, 1914, in Baltimore, without having been registered and licensed as provided by law. Case dismissed upon recommendation of State because of age and poverty of defendant.

Dr. Marshall G. Smith, 118 N. Calhoun St., Baltimore, for refusal or neglect to report the birth of the infant of Mrs. Harry Russell Miller, Sr., who was born May 12, 1915, in Baltimore. Defendant plead guilty. Case dismissed by magistrate.

Dr. James B. Wiltshire, 819 N. Eutaw St., Baltimore, for refusal or neglect to report the birth of the infant of Mrs. Leland Windsor, who was born September 19, 1914, in Baltimore. Case tried in Criminal Court. The evidence given by the State proved conclusively that defendant had not reported this birth but the Judge declared him not guilty.

Dr. Harry Fahrney, Myersville, Frederick County, for refusal or neglect to report the birth of the infant of Mrs. William Snyder, who was born January 14, 1915. Defendant plead guilty and was fined \$5.00 and costs.

H. G. Walker, Pleasantville, Harford County, for interring the body of Lida Hitchcock, without first obtaining a burial permit. Defendant plead guilty and was fined \$20.00 and costs.

Dr. T. R. Gough, Barnesville, Montgomery County, for refusal or neglect to report the birth of the infant (stillborn) of Mrs. Mamie Kuster, who was born dead December 10, 1914. Case stetted. Dr. Gough had made attempt to follow law.

Dr. T. A. Poole, 5611 Connecticut Ave., Chevy Chase, D. C., for refusal or neglect to report the birth of the infant of Mrs.

George F. Hobson, who was born February 27, 1915. Defendant plead guilty and was fined \$5.00 and costs.

Dr. George I. Eppart, Benning, D. C., for refusal or neglect to report the birth of the infant of Mrs. Jesse L. White, who was born February 3, 1915. Defendant plead not guilty, but was convicted and fined \$5.00 and costs.

Mrs. Mary Thomas, Cordova, Talbot County, for refusal or neglect to report the birth of the infant of Mrs. Jesse Suell, who was born October 22, 1914. Defendant plead not guilty and was fined \$5.00.

J. Radcliffe Farlow, Willards, Wicomico County, for interring the body of Mrs. Martha Ann Parker, December 22, 1914, in Wicomico County, without first obtaining a burial permit. Defendant plead guilty and was fined \$20.00 and costs.

(Sarah Keene, Madison, Dorchester County, for attending Mrs. Everett Bell at childbirth, November 14, 1914, without having been registered and licensed as provided by law. Found guilty and fined.) Case tried by County Registrar.

# REPORT OF BUREAU OF COMMUNICABLE DISEASES

#### C. HAMPSON JONES, Chief

Baltimore, Md., April 18, 1916.

Dr. John S. Fulton,

Secretary, State Department of Health,
- Baltimore, Md.

#### SIR:

I have the honor to transmit herewith the report of the Bureau of Communicable Diseases for the calendar year 1915. The data contained therein have been arranged under the following six sub-headings:

- I. Notifiable Disease Reports.
- II. Sanitary Surveys and Investigations.
- III. Public Health Conditions by Counties.
- IV. Occupational Diseases.
- V. Cases Examined for Diagnosis.
- VI. Prosecutions.

#### I. NOTIFIABLE DISEASE REPORTS.

In 1915, in the counties of Maryland,\* there were 9,954 cases of notifiable diseases reported, as contrasted with 6,631 cases in 1911, 5,821 cases in 1912, 12,113 cases in 1913, and 9,061 cases in 1914.\*\* Typhoid fever, diphtheria, scarlet fever, measles, chicken-pox, whooping cough, influenza and mumps were the prevailing epidemics.

<sup>\*</sup>The State of Maryland, exclusive of Baltimore City.

<sup>\*\*</sup>The above figures do not include the county tuberculosis cases.

The following is a detailed statement of the 1915 figures, arranged in numerical order, with comparative figures for 1911, 1912, 1913 and 1914.

# CASES OF SICKNESS FROM NOTIFIABLE DISEASE IN THE COUNTIES OF MARYLAND FOR 1915, 1914, 1913, 1912 AND 1911.

Name of Disease.	1915.	1914.	1913.	1912.	1911.
Typhoid fever	2.249	1.860	2.983	1.791	1,999
Diphtheria	1,390	790	1.049	785	783
Scarlet fever	1.141	1.655	822	607	656
Measles	1.049	2,000	5,352	1,675	1,706
Chickenpox	984	732	549	251	274
Whooping cough	906	768	812	441	453
Influenza	854	31	13	61	41
Mumps	602	470	287	103	605
German measles	206	259	29	11	13
Erysipelas	- 95	42	30	18	15
Acute dysentery	89	30	. 1	1	3
Smallpox	82	216	132	20	31
Malaria	64	96	13	14	27
Meningitis	54	39	11	6	8
Septicemia	49	16	1	4	1
Anterior poliomyelitis	40	12	9	32	12
Septic sore throat	39	18	5	0	0
Pellagra	13	2	5	0	3
Para-typhoid fever	9	0	_ 0	0	0
Ophthalmia neonatorum	7	2	$^2$	0	0
Trachoma	3	0	0	1	0
Cervical adenitis	3	0	0	0	0
Scabies		4	0	0	0
Foot-and-mouth disease	1	0	0	0	0
Conjunctivitis	1	0	0	0	0
Rabies		0	0.	0	0
Anthrax	1	0	0.	0	0
Impetigo contagiosa	0	13	0	0	0
Puerperal sepsis	0	3	0	0	0
Tetanus	0	2	. 0	0	1
Ring-worm	0	1	0	0	0
Catarrhal conjunctivitis	0	0	8	0	0
Yearly Totals	9,933	9,061	12,113	5,821	6,631
Belated returns of typhoid					
fever	21				
Total	9,954	9,061	12,113	5,821	6,631

In the foregoing summary of cases of sickness from notifiable disease, the total is 9,954. There were reported during the year 3,215 cases of tuberculosis, as compared with 3,302 cases reported in 1914, distributed as follows:

- (a) Baltimore City, 2,051.
- (b) Counties of Maryland, 1,164.

By including the 1,164 county cases of tuberculosis, the figures for the year are raised to 11,118.

Compared with the preceding year there is a very noticeable increase in the total number of cases of notifiable disease reported. This is especially due to a high attack rate in typhoid fever, diphtheria, scarlet fever and measles. Good returns were received from seven counties; fair returns from ten; in different returns from five; while the returns from one—Cal vert County—are very poor, totaling only 73 cases for the entire year.

The following table will give a condensed summary of the status of the 23 counties of Maryland in regard to the reporting of notifiable diseases:

STATUS OF THE COUNTIES OF MARYLAND, NOTIFIABLE DISEASE REPORTS, 1915.

Pe	pulation.		
Allegany	67.008	962	146.0
Anne Arundel	39,518	388	91.8
	139,012	2,131	153.0
Calvert	10.379	73	70.3
Caroline	20,781	196	94.3
Carroll	33,974	748	229.0
Cecil	23,283	260	112.0
Charles	15,713	169	108.0
Dorchester	29,042	367	126.0
Frederick	53,070	554	104.0
Garrett	21,373	117	54.7
Harford	27,805	285	103.0
Howard	15,785	244 -	155.0
Kent	15,992°	338	211.0
Montgomery	32,953	466	141.0
Prince George's	39.443	554	140.0
Queen Anne's	16.034	154	96.0
Somerset	26,736	$352^{\circ}$	131.0
St. Mary's	16,950	205	121.0
Talbot	19,239	311	161.0
Washington	51.981	664	128.0
Wicomico	28,905	180	62.3
Worcester	22,356	215	96.1

## STATUS OF NOTIFICATION OF INFECTIOUS DISEASE IN THE COUNTIES OF MARYLAND, 1915.

		Cases	Classifi-
Rank.	County.	Reported.	cation.
1st	Baltimore County	. 2,131	"Good"
2nd	Allegany County		"Good"
3rd	Carroll County	. 748	"Good"
4th	Washington County	. 664	"Good"
5th	Frederick County	. 554	"Good"
6th	Prince George's County	. 554	"Good"
$7 \mathrm{th}$	Montgomery County	. 466	"Good"
8th	Anne Arundel County	. 388	"Fair"
9th	Dorchester County	. 367	"Fair"
10th	Somerset County	. 352	"Fair"
11th	Kent County		"Fair"
12th	Talbot County	. 311	"Fair"
13th	Harford County	. 285	"Fair"
14th	Cecil County	. 260	"Fair"
15th	Howard County	. 244	"Fair"
16th	Worcester County		"Fair"
17th	St. Mary's County		"Fair"
18th	Caroline County	. 196	"Poor"
19th	Wicomico County	. 180	"Poor"
20th	Charles County	. 169	"Poor"
21st	Queen Anne's County	. 154	"Poor"
22nd	Garrett County	. 117	"Poor"
23rd	Calvert County		"Very poor"

The reported cases of notifiable disease, rural Maryland (the State of Maryland, exclusive of Baltimore City), for 1915, by months and diseases, will next be shown in the form of a table. A glance at these figures will convince one that some of the minor notifiable diseases are not properly reported.

Three of the rarer communicable diseases appear in the list for this year. These are: A case of foot-and-mouth disease in February; a case of rabies in April, and a case of anthrax in July. Each is an example of a disease of animals communicable to man.

REPORTED CASES OF NOTIFIABLE DISEASES—RURAL MARYLAND, BY MONTHS AND DISEASES, 1915.\*

. Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Measles Chickenpox Whooping cough Influenza Mumps German measles Erysipelas Acute dysentery Smallpox Malaria Meningitis Septicemia Anterior poliomyelitis. Septic sore throat Pellagra Para-typhoid fever. Ophthalmia neonatorum. Trachoma Cervical adenitis Scabies Foot-and-mouth disease. Conjunctivitis Rabies	100 103 180 50 194 175 31 104 2 2 2 2 2 2 2 1 0 0 0 1 1 0	62 45	47   105   98   78   243   111   17   111   1   3   0   4   4   8   1   2   2   0   0   0   0   0   0   0   0   0   0	76 88 70	74 30 70 218 53 66 72 24 92 10 4 17 1 4 7 3 0 0 0 0 0 0 0	944 355 899 213 700 1555 19 32 477 7 1 6 4 4 5 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	96 20	446   255   122   644   0   3   0   7   17   0   144   3   4   10   0   0   0   0   0   0   0   0   0 	4 8 2 4 4 31 0 9 6 6 1 1 6 6 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2277 511 566 499 411 22 15 06 17 01 18 3 6 3 3 1 2 0 0 0 0 0 0 0 0 0 0 0 0 0	355 140 466 961 81 80 26 1 100 2 2 4 4 11 3 10 0 0 0 0	214 115 95 80 83 21 8 6 2 2 1 1 1 7 2 0 4 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2,249 1,390 1,141 1,049 984 906 854 602 206 95 89 82 64 49 40 39 77 3 3 1 1
Anthrax		873						$\frac{0}{636}$		862	1,016	968	
			939		750						1,018		

<sup>\*</sup>Tuberculosis separately compiled.

The statistics of tuberculosis, a disease of State-wide importance, will next be displayed in a separate table. These figures, indicating especially sex and color, include Baltimore City. The total number of cases of tuberculosis registered in 1915 in the entire State is 3,215. The reporting is better in Baltimore City than in the counties. Taking population for population, the attack rate is considerably higher in the colored race than in the white race, the proportion being as 36 to 21 per 10,000.

#### TUBERCULOSIS, MORBIDITY, MARYLAND, 1915.

	Male. White.		Male, Colored.	/	Total.
Baltimore City Counties of Maryland		$\frac{632}{471}$	$\frac{278}{142}$	$\frac{251}{152}$	$2,051 \\ 1,164$
Entire State	1,289	1,103	420	$\frac{-}{403}$	3,215

Tuberculosis morbidity statistics, arranged according to color and age-periods, will appear in the next table. In both races the attack rate is highest in the quadrennium from 20-25, gradually declining thereafter up to and including extreme old age. This disease is more frequent in early life than was formerly supposed.

TUBERCULOSIS MORBIDITY FOR 1915, ARRANGED ACCORDING TO COLOR AND AGE-PERIODS.

on 4 08 4 0 order: 0 4 4 0 order: 0 4 4 0 order: 0 4 4 0 order: 0 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	ig	•	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	81		5 29 2,392 0 18 823	47
.08-57 - ≎	1-		रू न			12	13
.87-07 I o	=		15	16		26	27
.05-35 # co	37		19	13		53	59
,68–08 <sub>75</sub>	40		90 4	34		ලි ල	74
를 잃 모든-60.	62		35 80	43		38 8	105
29 20-22.	87		39	57		100 35	1#1
35, H = 50.	112		51 14	8		161 49	210
.61-01 5 5 40-45.	168		60 17	2.2		186 59	245
.01-68 55 40.	221		22	115		258 78	336
67 20–35.	268		$\begin{array}{c} 115 \\ 28 \end{array}$	143		316 95	411
.08-32 52 4	276		124 27	151		$\begin{array}{c} 326 \\ 101 \end{array}$	427
& to 20–25.	333		135 62	197		$\frac{380}{150}$	530
5 g 15-20.	192		22.23	134		$\frac{213}{113}$	326
.čI-01 작 등	19		23 41	13		88 88	113
.01-3 # F	29		8 8 8	87		88	87
12 13 0-2.	46		4 9	10		33 53	26
Baltimore City.           White	Total	COUNTIES OF MARYLAND.	Vhite	Total	STATE OF MARYLAND.	Vhite	Total

In the next table the tuberculosis morbidity figures will be arranged according to counties, with the sex and color also indicated. The Baltimore City figures are not included in this table.

The largest number of cases (253) was received from Baltimore County; the next largest number (118) was received from Allegany County, while Washington County, with reports of 100 cases, ranks third. Only five reports were received from Garrett County, all five cases being white persons, one male and four females.

Among white persons the female cases predominate in fifteen of the counties, while in seven the male cases are in the ascendency. In one county—Howard—the number of white cases in each sex is even, being seven males and seven females. In the colored race the attack rate is about even for both sexes throughout the counties. In Kent County all the white cases, seven in number, are females.

TUBERCULOSIS MORBIDITY FOR 1915, ARRANGED BY COUNTIES.

	White Male.	White Female.	Total White.	Colored. Male.	Colored. Female.	Total Colored.	Total.
Allegany	51	55	106	8 ″	4	12	118
Anne Arundel	14	17	31	18	- 17	35	66
Baltimore	117	113	230	9	14	23	253
Calvert	2	5	7	2	2	4	11
Caroline	9	7	16	$\frac{2}{2}$	0	2	18
Carroll	13	19	32	1	3	4 5	36
Cecil	7	6	13	. 7	1	5	18
Charles	6	5	11	. 7	7	14	25
Dorchester	. 11	12	23	8	14	22	45
Frederick	$\sqrt{22}$	26	48	5	11	16	64
Garrett	1	4	5	0	0	0	5
Harford	9	12	21	4	5	. 9	30
Howard	7	7	14	3	5	8	22
Kent	0	7	7	7	8	15	22
Montgomery	22	29	51	, 8	6	14	65
Prince George's	13	11	24	17	5	22	46
Queen Anne's	6	4	10	5	5	10	20
Somerset	14	23	37	8	10	18	55
St. Mary's	5	4	9	4	, 5	9	18
Talbot	9	10	19	7	12	19	38
Washington	39	54	93	4	3 8 8	7	100
Wicomico	14	26	40	3	8	11	51
Worcester	8	15	23	7	8	15	38
Total	399	471	870	141	153	294	1,164

The tuberculosis morbidity according to months, displayed in the subjoined table, shows that the largest number of cases of this disease in the entire State occurred in March, with July next. In Baltimore City the largest number of cases reported in any one month was in July.

#### TUBERCULOSIS MORBIDITY ACCORDING TO MONTHS, YEAR 1915.

	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Counties of Maryland Baltimore City State of Maryland	185	148	202		174	178	213	193	176	151	130	107	

#### II. SANITARY SURVEYS AND INVESTIGATIONS.

A total of 35 investigations were made in 1915 under the direction of this Bureau. These cover a wide range of topics. A summary of the results of the most important of these investigations will next be given, arranged in chronological order.

On January 22nd and 23rd an investigation was made of the smallpox situation at Pocomoke City, Worcester County. For two months smallpox had prevailed in Pocomoke City and vicinity, due largely to failure on the part of the authorities to enforce the vaccination law. Ten cases of sickness were reported, one terminating fatally. This figure, however, only includes the well-defined cases, a number of light ones going on to recovery unrecognized early in the outbreak.

On February 5th an investigation was made of the scarlet fever situation at Havre de Grace, Harford County. A total of nine cases of sickness occurred-3 in January, 5 in February and 1 in March. The public school buildings and moving-picture parlors were disinfected by the

spray method, using a watery solution of pyxol, 1 in 500. On February 6th a scarlet fever outbreak at St. Mary's Industrial School for Boys, Wilkens Avenue extended, was investigated. A total of 19 cases developed before the outbreak was checked, 9 occurring in February and 10 in March. The disease was probably introduced into the school from Virginia.

On March 1st an investigation was made of an epidemic of mumps at the Crownsville State Hospital for the Colored Insane, a total of 38

cases occurring-2 in February and 36 in March.

On March 2nd an investigation was made of an outbreak of diphtheria at St. Mary's Orphan Asylum, Cold Spring Lane, Baltimore County. The first case, a fatal one, in a white child, aged 10 years, occurred at the end of February. Cultures were taken from all the children and the Sisters in attendance, 190 cultures in all, 20 of which

proved to be positive for diphtheria, making a total of 21 clinical cases and "carriers" at the institution.

On April 15th an investigation was made of a case of scarlet fever in a white male student, aged 21 years, at Western Maryland College, Westminster, Carroll County. A rigid quarantine was established, no other cases occurring at the institution. The source of the disease was unknown.

In May and June of the present year (1915) there was an outbreak of typhoid fever at the Maryland Tuberculosis Sanatorium, which attacked the several groups of patients and employes using milk from one of the seven dairymen supplying the institution. Upon investigation it was found that an unrecognized case of typhoid fever had occurred in April, in the wife of the dairyman furnishing the infected milk. The cutbreak subsided when the sale of milk from this dairy was stopped. The outbreak consisted of 27 cases, 19 of them tuberculous patients and 8 healthy persons. Three deaths occurred in the former group, and none in the latter.

On June 7th an investigation was made of the scarlet fever outbreak at Elk Ridge, Howard County, this disease having invaded a dairy supplying milk to the residents of the town. Pupils attending the public school also were attacked. The outbreak consisted of 15 cases.

On July 24th an investigation was made of a house outbreak of typhoid fever at Perryman, Harford County, where six cases occurred in one family. The infection was traced to a polluted subsoil well. The patients were colored—four males and two females. The first patient, a boy, 15 years old, died of the disease.

On July 31st an investigation was made of a case of ophthalmia neonatorum at St. Michaels, Talbot County. This baby, a colored twin, developed the disease when three days old. Both a physician and a midwife were in attendance. The child, almost entirely blind in both eyes, was brought to the Johns Hopkins Hospital for treatment.

Typhoid fever at Bel Air, Harford County. Date of inspection, September 16th. Five cases of this disease were investigated and traced to a dairy supplying milk to the town. One of the five cases, a colored boy, aged 13 years, had been employed in this dairy, delivering milk to customers. He developed typhoid fever on August 15th, and continued to deliver milk until August 20th.

On October 12th an investigation was made of a diphtheria outbreak in the public schools at Hancock, Washington County. A total of 239 cultures were taken, of which 221 proved to be negative and 18 positive. In addition, 15 clinical cases occurred in the town and its environs.

On October 24th, an investigation was made of typhoid fever in Hagerstown. Information was obtained concerning 33 cases. The cases were scattered, and no general cause could be assigned. With two or three exceptions, all of these cases occurred in October.

On November 5th and 8th an investigation was made of scarlet fever conditions at Havre de Grace, Harford County, eight cases occurring within the past month, all white. Later on, another white case was discovered, and also a case in a colored boy attending school.

On November 17th to 19th an investigation was made of scarlet fever and diphtheria conditions at St. Inigoes, Ridge and Scotland, in St. Mary's County. Ten cases of diphtheria occurred at Scotland and one at Ridge; at the latter place 3 cases of scarlet fever occurred.

On December 3rd an outbreak of typhoid fever, 11 cases in all, was investigated at Chase, Baltimore County. The disease was attributed to pollution of the subsoil wells, some of them only eight feet deep, by the surface privies.

On December 20th—21st an outbreak of typhoid fever, eight cases in all, was investigated at "The Rocks," Harford County. These cases, occurring in three households, all intimately related, were due to contact. All were white persons of the poorer class, with insanitary home surroundings.

#### III. Public Health Conditions by Counties.

The State of Maryland is arbitrarily divided into two public health jurisdictions—the one composed of Baltimore City alone, the other comprising the State of Maryland; exclusive of Baltimore City. The former has an area of 31.64 square miles, with an estimated (1915) population of 584,608; the latter—the special province of the State Department of Health—has an area approximating 10,000 square miles, with an estimated (1915) population of 767,332. It is ordinarily spoken of as "rural Maryland," or the "counties, of Maryland."

A series of 24 tables will follow. The first, a summary for the entire 23 counties, portrays notifiable disease conditions throughout rural Maryland, by months and counties, for 1915. This will be succeeded by a table for each of the counties, arranged alphabetically, giving the number of reported cases by month and disease. Epidemiological notes are also appended.

At the head of each county table the area in square miles of the county is given, along with the estimated (1915) population. The estimated (1915) population of the entire State of Maryland is 1,351,940.

The tuberculosis morbidity figures are not included in any of these tables.

Following the county morbidity tables is one entitled "Reported Cases of Notifiable Diseases—Rural Maryland, by Counties and Diseases, 1915." In this table, a comprehensive one compiled from the county figures, the totals of the various communicable diseases for the entire year are shown by counties and diseases. In the typhoid fever column are included 21 belated reports of cases of this disease, raising the grand total for typhoid from 2,249 to 2,270 cases.

### REPORTED CASES OF NOTIFIABLE DISEASES, RURAL MARYLAND, BY MONTHS AND COUNTIES, 1915.

Counties.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Allegany Anne Arundel Baltimore Calvert	82 30 243 9	$\begin{vmatrix} 68 \\ 26 \\ 319 \\ 1 \end{vmatrix}$	67 67 230 4	60   36   186   2		73 43 146 11	42 23 86 10	48 27 134 17	87 25 128	121 18 113 8	125 21 156	32	962 388 2,131 73
Caroline Carroll Cecil	26 43 23	13 64 20	11 148 6	$10 \\ 218 \\ 12$	38 53	0 44 61	1 51 8	17 26 6	19 26 15	27 26 33	46 23 6	22	196 748 260
Charles	11   31   64   10	6 47 32 9	7   23   33   5	7   18   38   11	$\begin{vmatrix} 3 \\ 10 \\ 56 \\ 2 \end{vmatrix}$	13 72 8	$\begin{vmatrix} 9 \\ 21 \\ 27 \\ 3 \end{vmatrix}$	27   42   35   8	14 30 43 9	41 25 46 14	$egin{array}{c c} 26 \\ 72 \\ 57 \\ 24 \\ \end{array}$	35 51 14	367 554 117
Harford	55 22 15 20	39 15 13 66	19   12   29   23	14 52 7 53	16   10   95   38	8 21 82 30	19 17 22 81	8 30 15 21	24 25 23 26	21 13 17 42	40   9   10   41	22   18   10   25	285 244 338 466
Prince George's	21 16 3 62	33 14 2 28	34 7 50 24	42 6 43 17	48 1 40 33	40 12 14 19	42 16 10 15	38 27 2	67 15 3 24	70 22 5 23	69 9 30 49	50 9 3 40	554 154 205 352
Talbot Washington Wicomico	33 33 12	14 32 7	92 35 10	30 40 20	11 45 3	51 10	11 19 18	24 23 18	30 29 21	$15 \\ 114 \\ 21$	$\begin{array}{ c c c }\hline 34 \\ 126 \\ \hline 21 \end{array}$	12 117 19	311 664 180
Worcester	33     897	5     873	3     939	$\frac{5}{ }$ $ 927 $	$\frac{6}{748}$	$\frac{17}{782}$	$\frac{19}{570}$		$\frac{26}{715}$	$\frac{27}{862}$	$\begin{vmatrix} 22 \\ \\ 1,016 \end{vmatrix}$	$\frac{27}{968}$	$\frac{215}{9,933}$

#### Belated Returns of Typhoid Fever from Frederick City Hospital, 1915.

Carroll	0	0	0	0	0	0	1	0	0	0	0	0	1
Frederick	1	0	$  0 \rangle$	0	<b>2</b>	0	3	1	5	0	1	0	13
Montgomery	0	0	0	0	0	0	0	1	2	0	0	0	3
Washington	0	0	0	0	0	0	0	0	3	0	0	0	3
			ĺ	<b></b>	ĺ	Í			<b></b> -		i		
Total	897	873	939	927	748	782	570	636	715	862	1,016	968	9,933
Complete Total	898	873	939	927	750	782	574	638	725	862	1,017	968	9,953

#### Belated Returns of Typhoid Fever from a Death Certificate, 1915.

Charles	0	0	0	0	0	0	0	0	0	0	1	0	1
Complete Total	  898	873	<del></del>    939	$ 927 $	${750}$	782	574	 638	725	862	1.017	  968	9,953
Grand total	898	873	939	927	750	782	574	638	725	862	1,018	968	9,954

#### ALLEGANY COUNTY.

(Area in Square Miles, 442; Population, 67,008.) REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever	24	17	17	3	4	9	20	25	39	20	22	17	217
Diphtheria	5	6	11	2	3	4	2		28		73		260
Scarlet fever	25	26	19	24	40	26	$\frac{2}{9}$	8 7	4			2	194
Whooping cough	0	0	0	0	0	1	$\tilde{2}$	4	3	$egin{array}{c c} 5 \\ 3 \\ 2 \\ 0 \\ 2 \\ 1 \\ 0 \\ \end{array}$	5	$1\overline{2}$	
Chickenpox	17	- 6	4	6	4	9	2 2 0	$\begin{vmatrix} 4 \\ 2 \end{vmatrix}$	1	2	12	11	
Measles		1 9	4 7	3	11	16	0	0	3	0	2	0	
Mumps	1 8	9	7	10	7	7	2 1	1 1	3	2	0	2	58
Erysipelas	0	0	2	1	$^{2}$	0	1	1	0	1	2	0	10
German measles	1	0	0	0	0	0	1	0	0	0	0	1	3
Influenza	1	0	1	6	3	1	0	0	0	0	0	28	40
Septicemia	0	1	1	$\begin{array}{c} 6 \\ 2 \\ 0 \end{array}$	$\frac{3}{2}$	0	3	0	0	0		1	10
Septic sore throat	0	$egin{array}{c} 1 \ 2 \ 0 \end{array}$	0		0	0	0	0	3	0	0	0	5
Meningitis	0		1	0	0	0	0	0	1	0	0	$\begin{vmatrix} 2\\0 \end{vmatrix}$	4
Poliomyelitis	0	0	0	3	0	0	0	0	0	$\begin{array}{c c} 2 \\ 3 \\ 2 \end{array}$	0	0	5
Dysentery	0	- 0	- 0	0	0	0	0	0	$\frac{2}{0}$	3	0	0	5
Trachoma	0	0	0	0	0	0	0	0		2	0	0	2
Para-typhoid	0	0	0	0	0	0	0	0	0	0	2	0	2
Total	82	68	67	60	76	73	42	48	87	121	125	113	962

Typhoid fever prevailed to an undue extent in Westernport, with 16 cases in January, 12 in February and 10 in March. In July there were 14 cases of this disease reported from Cumberland; in August, 9 from Cumberland, 7 from Westernport and 5 from Luke; in September, 13 from Cumberland and 7 from Frostburg; in October, 8 from Cumberland and 4 from Luke; in November, 7 each from Westernport and Cumberland; in December, 3 each from Luke and Eckhart.

Several mild outbreaks of diphtheria occurred during the year. In March there were 5 cases in Cumberland and 3 in Lonaconing; in August, 3 each in Cumberland and Lonaconing; in September, 12 from Frostburg and 5 in Lonaconing; in October, 26 in Frostburg, 12 at Moscow Mills, 10 each at Lonaconing and Eckhart, and 6 in Cumberland; in November, 25 at Frostburg, 17 at Cumberland, 11 at Lonaconing and 5 at Grahamtown. In December there were 14 cases in Frostburg and 10 in Cumberland.

Scarlet fever cases occurred, in undue numbers, as follows: In January there were 8 cases at Frostburg and 5 at Eckhart; in February, 6 at Frostburg, 4 at Lonaconing and 3 at Eckhart; in March, 8 at Frostburg and 3 at Lonaconing; in April, 16 cases at Lonaconing; in May, 18 at Lonaconing, at Cumberland 8, and 4 each at Westernport and Gilmore. In June there were 12 cases at Lonaconing, 7 at Cumberland, 5 at Benfield and 4 at Frostburg.

In December there were 8 cases of whooping cough reported from Cumberland and 4 from Morantown.

A few cases of chickenpox were present in Cumberland, 9 cases having been reported in January, 7 in June, 5 in November and 7 in December. In January 6 cases were reported from Lonaconing.

In May, 6 cases of measles were reported at Keyser, R. F. D.; in June, 12 cases were reported from Cumberland.

During the first half of the year mumps was reported from Cumberland, as follows: In January, February, March, May and June, 7 cases in each month; in April, 10 cases.

In December, 28 cases of influenza were reported from Cumberland.

In February, 2 cases of septic sore throat were reported from Lonaconing; in September, 2 from Cumberland and 1 from Allegany Grove. A few scattered cases of infantile paralysis occurred. In April, 1 case each was reported from Cumberland, Lonaconing and Flintstone; in October, 1 case each from Cumberland and Lindnerville.

Two imported cases of trachoma were treated at the Western Maryland Hospital early in the year. Both were white men, Italian laborers, brought to the hospital from West Virginia.

#### ANNE ARUNDEL COUNTY.

(Area in Square Miles, 425; Population, 39,518.) REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Whooping cough. Chickenpox Measles Septicemia Mumps Influenza German measles. Pellagra Meningitis Dysentery Erysipelas Malaria Poliomyelitis Total	0 7 13 1 7 1 1 0 0 0 0 0 0 0 0	$\begin{array}{c} 3 \\ 0 \\ 13 \\ 2 \\ 4 \\ 2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	$\begin{bmatrix} 0 \\ 2 \\ 10 \\ 0 \\ 0 \\ 36 \\ 36 \\ 34 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ $	$\begin{array}{c} 4 \\ 2 \\ 6 \\ 3 \\ 9 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ - \\ 36 \end{array}$	$\begin{bmatrix} 6 \\ 2 \\ 1 \\ 15 \\ 4 \\ 0 \\ 0 \\ 0 \\ 4 \\ 0 \\ 0 \\ 0 \\ 1 \\ 7 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	$\begin{bmatrix} 7 \\ 1 \\ 7 \\ 4 \\ 8 \\ 10 \\ 1 \\ 1 \\ 2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \\ 43 \end{bmatrix}$	$ \begin{array}{c} 12 \\ 3 \\ 0 \\ 0 \\ 3 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	21 0 0 0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0	$ \begin{array}{c} 23 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	5 4 2 0 3 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0	8 3 6 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	5 5 11 0 7 2 0 0 1 1 0 0 0 0 0 0 0 0 0 0 0 0	59 18

In May, 4 cases of typhoid fever occurred at Annapolis; in July, 3 cases each at Deal and Churchton. In August, 6 cases were reported from Annapolis, R. F. D., 3 from Glenburnie and 2 from Churchton; in September, 6 cases from Annapolis and 3 from Mayo; in November, 3 from Curtis Bay.

Four cases of diphtheria were reported from Severn, R. F. D., in January; in December, 2 cases were reported from Curtis Bay.

In January, 7 cases of scarlet fever were reported from Brooklyn and 4 from Glenburnie; in February, 5 from Brooklyn and 3 from Glenburnie; in March, 5 from Waterbury and 3 from Eastport; in April, 5 from Brooklyn; in June, 5 from Benfield, R. F. D.; in November, 4 from Brooklyn; in December, 4 from Annapolis and 2 from Brooklyn.

Seven cases of whooping cough were reported from Riverview; in May, 3 from Bayard.

Nine cases of chickenpox were reported from Dorsey, R. F. D., in March, and 6 in April; in June, 6 cases of this disease were reported from Annapolis.

In March an epidemic of mumps prevailed at the Crownsville State Hospital, 36 cases being reported.

Two cases of pellagra were reported during the year, one in April, at Bridewell, the other in October, at Annapolis. The former was a colored male laborer 32 years old; the latter a white male 33 years old, a merchant by occupation.

Seven cases of erysipelas occurred at the Maryland House of Correction in May.

Two cases of infantile paralysis occurred in August, one at Brooklyn and one at Harwood.

#### BALTIMORE COUNTY.

(Area in Square Miles, 656; Population, 139,012.)
REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Whooping cough. Chickenpox Measles Mumps Erysipelas German measles. Influenza Septic sore throat. Pellagra Meningitis Septicemia Ophthalmia neonatorum. Cervical adenitis. Malaria Dysentery Poliomyelitis Para-typhoid Smallpox	15   31   35   5   105   9   32   1   1   7   1   0   0   0   0   0   0   0   0	$ \begin{vmatrix} 244 & 66 & 66 & 67 & 67 & 67 & 67 & 67 & $	13 39 7	5 17 10 15 1 19 55 0 2 0 0 0 1 0 0	10 11 12 1 12 34 11 1 12 7 0 0 1 0 0 0 0 1	20 10 21 18 17 28 10 3 3 12 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	27 4 13 14 2 14 3 0 0 0 0 0 0 1 0 0 0 1 0 0 0 0 1 0	$egin{array}{c} 0 \ 2 \ 0 \ 0 \ 1 \ 1 \ 0 \ 0 \ 5 \ 0 \ 6 \ \end{array}$		$ \begin{array}{c} 466 \\ 277 \\ 53 \\ 35 \\ 59 \\ 44 \\ 20 \\ 00 \\ 00 \\ 11 \\ 10 \\ 00 \\ 66 \\ 44 \\ 00 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	$egin{array}{c c} 9 & 6 & 25 \\ 19 & 1 & 0 \\ 4 & 0 & 2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \\ \end{array}$	16 18 39 87 13 1 2 10 1 0 0 0 0 0	116 346 242 262 17 57 124 3 6 13 6 2
Total	$ {243} $	  319 	  230 	186	  118 	  146  	86	134	128	113	<del></del>  156 	—  272 	2.131

In January, 3 cases of typhoid fever occurred at Highlandtown and 2 at Catonsville; in February, 4 cases at Highlandtown and 3 at Notre Dame College; in April, 3 cases at Highlandtown and 2 at Catonsville; in June, 5 cases at St. Mary's Industrial School and 2 at Raspeburg; in July, 7 cases at Highlandtown, 3 each at Riderwood and St. Mary's Industrial School and 2 at Mt. Winans. In August there were 10 cases of typhoid fever at Highlandtown, 5 at Granite and 3 each at Roland Park, Lauraville and St. Mary's Industrial School. In September there were 4 cases each at Turners Station, Pikesville, Highlandtown and Arlington, 3 each from Catonsville and Govans and 2 from St. Mary's Industrial School. In October there were 5 cases at Sparrows Point and Towson, 4 each at Granite and Mt. Winans and 3 at Halethorpe; in November.

11 cases were reported from Highlandtown; in December, 7 from Chase, 6 from Catonsville, R. F. D., and 3 from Mt. Winans.

Diphtheria cases increased in number in several localities. In January there were 6 cases at Hamilton, 5 at Roland Park and 3 at Lauraville; in February, 8 at Highlandtown and 4 at Roland Park; in March, 21 cases occurred at St. Mary's Orphan Asylum, 8 at Highlandtown and 3 at Edgemere; in April, 7 at Highlandtown. In September, 14 cases occurred at Kernan Hospital and 4 at Holbrook; in November, 13 at Highlandtown and 3 at Monkton; in December, 10 cases of this disease at Highlandtown and 2 at St. Mary's Industrial School.

In January, 7 cases of scarlet fever were reported from Sparrows Point and 6 each from Highlandtown and Canton; in February, 10 cases of this disease were reported from Highlandtown and 9 from St. Mary's Industrial School; in March, 10 cases from St. Mary's Industrial School and 5 from Canton; in April, 8 from Catonsville, 5 from Highlandtown and 1 from St. Vincent's Orphan Asylum; in May, 5 cases from Catonsville; in June, 5 from Highlandtown, 4 from Ilchester, R. F. D., and 2 from St. Mary's Industrial School. In July, 7 cases were reported from Highlandtown; in November, 5 from Arlington and 1 from St. Mary's Industrial School.

Eight cases of whooping-cough occurred at Catonsville in March, and also in July. In June, 6 cases of this disease were reported from Relay and 5 from Grange. In August, 15 cases were reported from Catonsville; in December, 6 cases each from Fork and Arlington.

Several small outbreaks of chickenpox occurred. In January, 44 cases occurred at Catonsville, 17 at Relay and 7 each at Mt. Washington and Raspeburg. In February there were 28 cases of this disease at Catonsville, 20 at Mt. Washington and 10 at Highlandtown; in March, 8 cases at Loch Raven and 7 at Maryland School for Boys; in December, 17 cases at Rosewood State Training School and 11 at Catonsville.

Outbreaks of measles occurred at several places. In May, 13 cases were reported from Relay and 5 from Bengies; in June, 9 cases from Roland Park and 5 from Arlington; in November, 18 from Highlandtown; in December, 67 from Highlandtown and 9 from Canton.

In January, 13 cases of mumps were reported from Arling-

ton; in February, 80 cases from Arlington, 13 from Monkton, R. F. D., and 6 from Mt. Washington. In March, 24 cases of this disease were reported from Arlington; in April, 6 cases from Cockeysville; in November, 9 from Sparrows Point; in December, 8 from Roland Park.

Ten cases of German measles were reported from Lansdowne in April.

Influenza was epidemic at Sparrows Point, 30 cases being reported in March and 41 in April. In December, 10 cases of this disease were reported from Highlandtown.

A few scattered cases of infantile paralysis occurred. In July, 2 cases were reported from Highlandtown and 1 from Cockeysville; in August, 1 case each occurred at Highlandtown, Rognel Heights, Hamilton, Sparrows Point, Texas and Overlea; in September, 1 case occurred at Halethorpe.

One case of smallpox was reported, in May, from Loch Raven. The patient was an unvaccinated colored man working on the Loch Raven dam.

#### CALVERT COUNTY.

(Area in Square Miles, 222; Population, 10,379.) REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Whooping cough Chickenpox Measles Influenza Meningitis Total	$\begin{bmatrix} -4 \\ 1 \\ 3 \\ 0 \\ -1 \\ 0 \\ - \\ 9 \end{bmatrix}$	$\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ \\ 1 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 3 \\ 0 \\ 0 \\ 0 \\ - \\ 4 \end{bmatrix}$	$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 2 \\ 0 \\ - \\ 2 \end{bmatrix}$	$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -2 \\ -2 \end{bmatrix}$	1 8 0 2 0 0 0	5 0 3 0 0 0	15 0 0 2 0 0 0 17	6 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	$egin{bmatrix} 7 \\ 1 \\ 0 \\ 0 \\ 0 \\ - \\ 8 \\ \hline \end{bmatrix}$	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{bmatrix} 2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ \hline 3 \end{bmatrix}$	41 16 3 7 5 1 7

In January 3 cases of typhoid fever were reported from Wallville. In July, 2 cases of this disease were reported from Chesapeake Beach; in August, 4 from Broome's Island; in September, 4 from Willows.

Seven cases of whooping-cough were reported from Sollers in June.

In December a case of meningitis, male, white, was reported from Broome's Island.

#### CAROLINE COUNTY.

(Area in Square Miles, 320; Population, 20,781.)
REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	Scptember.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Chickenpox Septic sore throat Dysentery Foot-and-mouth disease Erysipelas Meningitis Influenza Malaria	0 12 4 0 8 1 1 0 0 0 0	0 9 1 0 2 0 0 1 0 0 0 0	0 0 8 0 0 0 0 0 1 2 0	1 6 0 0 0 0 0 0 0 0	3 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0	0 0 1 0 0 0 0 0 0 0	13 3 0 0 0 0 1 0 0 0 0 0	11 6 0 1 0 0 1 0 0 0 0	10 13 1 0 0 0 1 0 0 0 1 1	9 37 0 0 0 0 0 0 0 0	3 17 2 0 0 0 0 0 0 0 0 0	50 98 23 1 10 1 4 1 2 3 2

In August, 4 cases of typhoid fever were reported from Goldsboro, R. F. D.; in October, 3 cases of this disease were reported from Henderson, R. F. D.

Nine cases of diphtheria were reported from Denton in January and 4 cases in February. In September, 3 cases of this disease were reported from Ridgely; in October, 4 cases were reported from Preston; in November, 13 from Greensboro, 9 from Preston, R. F. D., and 4 from Preston.

In March, 6 cases of scarlet fever were reported at Preston, R. F. D.; in April, 3 cases of this disease occurred at Bethlehem. Seven cases of chickenpox were reported from Preston, in January.

In February a case of foot-and-mouth disease was reported in a white male, aged 39 years, residing at Denton, R. F. D. This man, E. C., a farmer, was taken ill about the middle of December, 1914. He was exposed to foot-and-mouth disease about Thanksgiving Day, 10 of his cattle and 7 of his hogs dying of it. The patient was isolated and treated. Neither his wife nor any of his children contracted the disease.

#### CARROLL COUNTY.

(Area in Square Miles, 437; Population, 33,974.)
REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever	5	1	0	1	2	0	6	15	13	12	5	6	6€
Diphtheria	1	1 8	$\frac{2}{4}$	1 4	$\frac{2}{0}$	ŏ	ő	0	5		10	3	
Scarlet fever	12		4	3	1	0	o	0	1		0	1	35
Whooping cough	0	8	6	31	19	39	36	4	3	1 5	0	8	159
Chickenpox	6	7	27	14	5		0	0	2	0	3	1	65
Measles	0	0	1	1	0	$\begin{bmatrix} 3\\2\\0 \end{bmatrix}$	5 2 0	0	0	0	1	1	9
Mumps	17	21	14	25	2	2	5	0	0	0	3	0	89
Erysipelas	1	$\frac{2}{0}$	0	0	0		2	1 4	0 2	0	1	2	9
Dysentery	1	0	0	0	0	0		4	2	0	0	0	7
Influenza	0	1		139	8		0	0	0		0		260
German measles	0	3	<b>0</b>	0	[ 1]	0	0	0	0	0	0	1	5
Septicemia	0	1	0	0	0		0	0	0		0	0	1
Meningitis	0	. 0	0	0	0	0	0	1	0	0	0	0	1
Poliomyelitis	$ \cdot $ 0	0	0	0	0	0	0	1	0	0	0	0	1
m 1												4.7	
Total	43	64	148	218	38	44	51	26	26	26	23	41	748

Belated Returns of Typhoid Fever from Frederick City Hospital, 1915.

Taneytown	0	0	0	0	0	0	1	0	0	0	0	0	1
Total	43 43	64 64	148 148	218 218	38 38	44 44	51 52	26 26	26 26	26 26	23 23	41 41	  748  749

In August, 5 cases of typhoid fever were reported from Sykesville, R. F. D.; in September, 3 cases were reported from Oakland Mills.

In February, 3 cases of diphtheria were reported from Finksburg; in October, 3 cases were reported from Finksburg, R. F. D.

In January, 4 cases of scarlet fever were reported from Lineboro and 3 from Taneytown; in February, 3 cases of this disease were reported from Taneytown.

In April, 16 cases of whooping cough were reported from Harney, R. F. D.; in May, 6 cases were reported from Harney and 5 from Key-Mar, R. F. D. In June, 15 cases were reported from Taneytown and 10 from Taneytown, R. F. D.; in July, 14 cases were reported from Taneytown and 10 from Sykesville.

In March, 12 cases of chickenpox occurred at Union Mills and 7 at Westminster; in April, 9 cases occurred at Oakland.

In January, 7 cases of mumps occurred at Oakland and 6 at Eldersburg; in February, 8 cases occurred at Oakland and 7 at Sykesville; in March, 4 cases at Oakland; in April, 8 at Berrett and 6 at Freedom.

Influenza prevailed, in March, to the extent of 28 cases at Union Mills, 24 at Manchester, 8 at Mt. Airy, 7 at Uniontown, 6 at Union Bridge and 5 at Hampstead and Cranberry. In April an excessive number of cases of this disease occurred, as follows: At Taneytown, 22 cases; at Springfied State Hospital, 21 cases; 17 cases occurred at Sykesville, 13 at Manchester, 8 at Key-Mar, R. F. D.; 7 at Uniontown, 6 each at Union Bridge and Westminster and 5 at Melrose. In May, 4 cases occurred at Woodbine.

One case of poliomyelitis (infantile paralysis) was reported in August, at Westminster, R. F. D., in a colored male.

#### CECIL COUNTY.

(Area in Square Miles, 360; Population, 23,283.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Chickenpox Measles Malaria Influenza Erysipelas Septicemia Meningitis Mumps Para-typhoid Dysentery	3 3 7 4 4 4 1 1 0 0 0 0 0 0	2 0 4 5 7 0 0 0 0 0 0 0	1 1 1 0 2 0 0 0 0 0 0 0 0 0 0	5 0 3 0 0 1 0 2 0 1 0 0 0 1 0 0 0 0 0 0 0	1 2 0 0 0 48 0 1 0 1 0 0 0 0 0 48 0 0 0 0 0 0 0 0 0 0 0 0 0	1 0 1 6 0 52 0 0 0 0 0 0	1 0 1 5 0 0 0 0 0 1 0 0 0 0	4 0 0 0 0 2 0 0 0 0 0 0 0 0 0	7 2 0 2 0 4 0 0 0 0 0 0 0 0	2 0 2 0 0 24 0 0 1 1 1 1 1	3 0 2 0 0 0 0 0 0 0 0 0 0 0	3 2 4 0 1 5 0 0 0 0 1 0 1 0	33 10 25 22 14 137 1 5 2 4 2 1 2
Total	23	20	6	12	53	61	-8	6	15	33	6	17	260

Two cases of typhoid fever were reported at Elkton, in April, and 2 at Elkton, R. F. D., in September.

Four cases of scarlet fever were reported at Chesapeake City, in January, 3 cases at Elkton, R. F. D., in February, and 2 at Elkton, in December.

In June, 4 cases of whooping cough occurred at Blythedale; in July, 5 cases occurred at Childs, R. F. D.

In May there were 36 cases of measles at Perryville and 6 at Northeast. In June, 21 cases occurred at Perryville, 6 at Childs and 4 at Rising Sun. In October there were 21 cases at Chesapeake City.

#### CHARLES COUNTY.

(Area in Square Miles, 451; Population, 15,713. REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Chickenpox Measles Influenza Poliomyelitis Dysentery Malaria Erysipelas	3 2 0 0 4 1 1 0 0 0	1 1 0 0 0 2 0 1 0	1 3 0 0 0 3 0 0 0	0 0 1 0 4 1 1 0 0 0	0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0	6 0 0 0 2 0 0 1 0 0	15 5 0 0 5 0 0 1 1	11 1 0 0 1 1 1 0 0 0	30 1 0 0 8 0 0 1 0	7 1 2 4 11 1 0 0 0	8 2 0 0 5 1 0 0 0	82 16 3 4 40 15 2 4 1
Total	 11	6	$$ $\frac{1}{7}$	   7	3		$\frac{}{9}$		 14	41	<del>26</del>	16	169

Belated Returns of Typhoid Fever from Death Certificate, 1915.

Riverside, R. F. D	ÍI				I							1	
Total	11 11	6 6	7	7 7	3 3	$\frac{2}{2}$	9	27 27	14 14	41 41	26 27	16 16	169 170

In August, 4 cases of typhoid fever occurred at Hughesville; in October, 9 cases of this disease occurred at La Plata, and 3 each at Port Tobacco and Pomfret. In December, 5 cases occurred at White Plains.

Three cases of diphtheria were reported from Berry, in August.

Four cases of measles were reported from Pisgah, in October; in November, 9 cases were reported from Marbury.

One case of poliomyelitis (infantile paralysis) was reported from Wicomico, in January. The patient was a colored male aged 16 years, attending the Wicomico public school. In April a case of this disease was reported from Grayton. The patient was a colored female infant, one year old.

#### DORCHESTER COUNTY.

(Area in Square Miles, 618; Population, 29,042.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fovor	11	4		7	4	7	15	29	20	14	$oxed{20}$	10	145
Typhoid fever	3	4	4	- 1	1	0	3	3	9	7	44	17	93
Diphtheria	7	16	4	ō	$\begin{bmatrix} \hat{\mathbf{o}} \end{bmatrix}$	5	1	0	0	- 1		5	43
Chickenpox	1	3	6	1	0	1	0	1	0	1	$\frac{4}{3}$	0	17
Measles	1	ő	- öl	ō	ŏ	0	ő	ō	ŏ	0	0	ŏ	1
Mumps	8	20	6	6	ŏ	ŏ	ŏ	1	ŏ	ŏ	ő	0	41
Septicemia	0	ő	1	o	ŏ	ŏ	o	1	ŏ	ő	ŏ	ŏ	2
Influenza	ŏ	ŏ	ō		4	ő	ŏ	$\hat{0}$	ŏ	ŏ	ĭ	ŏ	7
Conjunctivitis	Ö	Õ	ő	$-\frac{2}{1}$	ō	o o	- 0	ő	ő	ő	õ	0	1
Dysentery	Ö	ŏ	ŏ	ō	ĭ	ŏ	1	4	ŏ	o	ŏ	ŏ	$\tilde{6}$
Erysipelas	0	0		0	0	0	1		1	0	0	0	3
Malaria	0	0	0	0	0	0	0	$egin{array}{c} 0 \ 2 \ 1 \end{array}$	0	0	0	0	2
Meningitis	0	0	0	0	0	0	0	1	0	0	0	3	4
Para-typhoid	0	0	0	0	0	0	0	0	0	1	0	0	1
Ophthalmia neonatorum	0	0	0	0	0	0	0	0	0	1	0	0	1
		[	[						_				
Total	31	47	23	18	10	13	21	42	30	25	72	35	367

Typhoid fever prevailed in Cambridge to the extent of 9 cases in January, 8 in July, 7 in August, 4 in October and 9 in November. In August, 5 cases of this disease were reported at Castle Haven; in September, 3 at East New Market; in November, 5 at Cambridge, R. F. D.

In September, 7 cases of diphtheria occurred at Church Creek, R. F. D.; in October, 5 at Church Creek. In November there were 14 cases of this disease at Williamsburg, 9 at Cambridge, 6 at Hurlock, R. F. D., and 5 at Madison. In December, 8 cases occurred at Cambridge.

In February there were 3 cases of scarlet fever each at Linkwood, Golden Hill and Golden Hill, R. F. D. In June, 3 cases occurred at Church Creek; in December, 4 at Cambridge.

Four cases of chickenpox were reported from Church Creek, R. F. D., in March.

At Cambridge there were 8 cases of mumps in January, 20 in February and 5 in March.

One case of ophthalmia neonatorum occurred in October. The patient, a colored female two weeks old, resided at Linkwood.

#### FREDERICK COUNTY.

(Area in Square Miles, 662; Population, 53,070.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Discuses.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Whooping cough. Chickenpox Measles Mumps Erysipelas German measles. Influenza Septicemia Meningitis Para-typhoid	3 9 23 19 3 1 2 1 2 1 0 0	4 5 7 7 6 0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 1 18 4 0 1 0 0 4 2 0 0	1 1 2 11 5 0 0 1 0 17 0 0 0	21 1 3 21 1 0 0 0 8 1 0	7 2 4 49 7 1 1 0 0 1 0 0 0 0 0	15 0 3 8 0 0 0 0 0 0 0	1 3 10 0 0 0 1 0 0 0 0 0	1 8 6 2 0 2 0 0 1 0 1	3 7 7 2 0 5 0 0 0 0 0	8 18 9 4 17 0 0 1 0 0 0 0	5 9 9 13 10 1 1 0 0 0	79 173 57 3 12 6 2 34 3 2
Dysentery	$\begin{array}{c c} 0 \\ 0 \\ \hline -64 \end{array}$	$\begin{vmatrix} 0 \\ 0 \\ -32 \end{vmatrix}$	$\begin{array}{c} 0 \\ 0 \\ \hline 33 \end{array}$	0 0 38	$\begin{array}{c} 0 \\ 0 \\ \hline - \\ \hline 56 \end{array}$	$0 \\ 0 \\ -72$	$\frac{0}{0}$	$\frac{0}{0}$ $\frac{0}{35}$	$\begin{array}{c} 0 \\ 0 \\ \hline 43 \end{array}$	$\begin{bmatrix} 2 \\ 0 \\ -46 \end{bmatrix}$	$\frac{0}{0}$	$\begin{array}{c} 0 \\ 2 \\ \hline 51 \end{array}$	$\frac{2}{2} = 554$

Belated Returns of Typhoid Fever from Frederick City Hospital, 1915.

Enadonisk	,		_	_		_	_		_	_			4
Frederick	1 1	U	U	l v	l v	U	l v	U	U	ļ v	0	U	1
Sabillasville	0	0	0	0	2	0	0	0	1	1 0	0	0	3
Brunswick	0	0	0	0	0	0	2	0	0	0	1	0	3
Buckeystown		0	0	0	0	0	1	1	0	0	0	0	2
Myersville	0	0	0	0	0	0	0	0	1	0	0	0	1
Frederick, R. F. D	0	0	0	0	0	0	0	0	1	0	0	0	- 1
Walkersville	0	0	0	0	0	0	0	0	1	0	0	0	1
Mt. Airy	0	0	0	0	0	0	0	0	1	0	0	0	1
	ÍI					[ <del></del> -	<b> </b>				II		
Total	64	32	33	38	56	72	27	35	43	46	57	51	554
							_						
Complete Total	65	32	33	38	58	72	30	36	48	46	58	51	567

In May the milk outbreak of typhoid fever began at the Maryland Tuberculosis Sanatorium, near Sabillasville, 18 cases being reported from the institution in this month. In September, 3

cases each, of typhoid fever, occurred at Frederick and Emmitsburg. In October, 6 cases of this disease occurred at Emmitsburg, and 3 cases each at Frederick and Brunswick.

In November, 7 cases of diphtheria occurred at Sabillasville. In January there were 8 cases of whooping cough reported from Woodsboro and 7 from Middletown. In March, 12 cases of this disease were reported from Emmitsburg and 6 from Jefferson; in May, 9 cases from Middletown and 5 from Myersville.

Nine cases of chickenpox were reported from Middletown, R. F. D., in November.

In April, 9 cases of influenza were reported from Myersville. In December, 2 cases of septic sore throat developed at the Maryland School for the Deaf, in Frederick.

In January there were 7 cases of scarlet fever at Mt. Airy, R. F. D., and 3 at Buckeystown.

#### GARRETT COUNTY.

(Area in Square Miles, 660; Population, 21,373.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Measles Septicemia Influenza Malaria Meningitis Para-typhoid Erysipelas Mumps Chickenpox Total	1 1 8 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 3 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0	0 6 2 1 0 0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0	1 1 2 3 0 0 0 0 0 0 1 0 0	0	1 0 0 0 0 0 0 0 0	3 6 15 0 0 0 0 0 0 0 0 0 0 0	0 7 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	26 36 30 6 3 2 5 1 1 1 1 1 4 

At Kitzmiller there were 3 cases of typhoid fever in each September and October. In September there were 2 cases of this disease at Accident; in October, 2 at Crellin.

In February, 3 cases of diphtheria occurred at Friendsville; in March there were 3 cases at Sines; in April, 5 at Davis; in June, 4 at Swanton, R. F. D.; in November, 5 at Sutton.

In January there were 3 cases of scarlet fever at Accident, R. F. D.; in November, quite a localized outbreak of this disease occurred at Jennings, 14 cases being reported.

#### HARFORD COUNTY.

(Area in Square Miles, 388; Population, 27,805.) Reported Cases of Notifiable Diseases, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough. Chickenpox Measles Mumps Influenza Septic sore throat Meningitis German measles Septicemia Pellagra Dysentery Erysipelas	2   5   15   1   4   0   21   3   4   0   0   0   0   0   0   0   0   0	0 5 11 10 1 0 0 11 0 0 0 0 0 0	1 7 2 5 0 0 0 0 0 2 0 0 0 0 0 0 0 0 0 0 0 0	2 1 1 4 1 1 0 0 0 0 0 0 0 0 0	1 1 1 5 0 2 2 2 2 0 1 1 0 0 0 0	0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		7 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	177 0 66 0 0 0 0 0 0 0 0 0 0 0 0 0 0 1 1 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	15 2 19 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4 4 11 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	71 28 78 34 6 4 38 9 2 2 1 1 1
Total	55	39	19	14	16	8	19	8	24	21	40	22	285

In July there were 7 cases of typhoid fever reported from Perryman; September, 4 from Bel Air; in November, 7 from The Rocks.

In January there were 5 cases of scarlet fever reported from Bel Air, R. F. D., 3 from Havre de Grace; in February, Havre de Grace, 5; Aberdeen, 3; in October, Havre de Grace, 3; Forest Hill, 3; in November, Havre de Grace, 7; Coopstown, 3; The Rocks, 3, and in December, Havre de Grace, 7.

In February there were 6 cases of whooping cough reported from Van Bibber.

In January there were 5 cases of mumps reported from Bel Air and 4 from Perryman; in February, 4 from Abingdon.

In January there were 2 cases of septic sore throat reported from Forest Hill.

#### HOWARD COUNTY.

(Area in Square Miles, 365; Population, 15,785.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

* Diseases.	January.	February.	March.	April.	Мау.	June.	July.	August.	September.	Oetober.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Chickenpox Measles Mumps Influenza Meningitis Erysipelas Septic sore throat Ophthalmia neonatorum Poliomyelitis Dysentery Septicemia	6 3 4 4 3 1 1 0 0 0 0 0 0 0	3 2 0 1 6 0 0 1 1 1 1 0 0 0	1 1 1 3 3 2 0 0 0 1 0 0 0 0 0	0 1 1 7 5 0 1 34 0 1 2 0 0	1 0 1 0 0 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 0 10 0 2 1 5 0 0 0 0 0	3 0 3 0 6 2 0 0 0 0 0 0 0 0 0	28 0 0 0 0 1 0 0 0 0 1 0 0 0	14 1 0 0 1 3 0 2 1 0 2 0 0	7 2 0 0 3 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4 1 0 0 3 1 0 0 0 0 0 0 0 0	2 3 2 0 4 4 4 0 0 0 0 0 0 0 0	70 14 22 15 36 18 8 42 4 2 6 1 3 2
Total	   22  	15	12	52	10	21	17	 ′ 30	25	13	9	18	244

In January there were 3 cases of typhoid fever reported from Ellicott City; in August, 5 from Ellicott City, 4 from Ellicott City, R. F. D.; in August, 7 from Glenelg; in September, 8 from Ellicott City; in October, 4 from Ellicott City.

In June there were 9 cases of scarlet fever reported from Elk Ridge.

In May there was 1 case of ophthalmia neonatorum reported from Ellicott City. The patient was a colored infant.

In May there was a case of poliomyelitis reported from Laurel, R. F. D., a colored male. In June, 2 cases, colored, one male and the other female, were reported from Gibson.

#### KENT COUNTY.

(Area in Square Miles, 281; Population, 15,992.)
REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Whooping cough. Chickenpox Measles Erysipelas Influenza Malaria Septicemia German measles. Mumps Anthrax Septic sore throat Dysentery	2 1 1 2 1 1 4 2 0 0 0 0 0 0	0 0 0 0 2 11 0 0 0 0 0 0	2 0 0 0 5 13 2 0 0 1 6 0 0 0	1 0 1 1 0 0 0 0 3 0 0 0 0	2 0 0 0 0 15 0 0 0 77 1 0 0	$\begin{bmatrix} 0 \\ 1 \\ 5 \\ 39 \\ 0 \\ 1 \\ 0 \\ 35 \\ 1 \\ 0 \end{bmatrix}$	0 1 0 1 3 15 0 0 0 0 0 0	9 0 0 0 0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0	11 1 0 0 0 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 5 0 0 0 0 0 0 0 0 0 0 0 0 0	5 3 0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 6 0 0 1 0 0 0 0 0 0 0 0 0	36 17 1 4 19 114 5 6 2 1 121 4 1 5
Total	   15	   13 	29	   7 	95	82	22	15	23	17	10	10	338

In August there were 4 cases of typhoid fever reported from Rock Hall and 3 from Millington; in September, 5 from Rock Hall.

In March there were 7 cases of measles reported from Kennedyville, R. F. D.; in February, 6 from Kennedyville; in May, 6 from Worton and 6 from Betterton, R. F. D.; in June, 11 from Still Pond, R. F. D.; from Coleman, 10; Betterton, 5, and Worton, 4. In July, 6 from Kennedyville, R. F. D., and 5 from Massey.

In May there were 37 cases of German measles reported from Coleman; from Still Pond, 19, and Worton, R. F. D., 11. In June, 17 from Coleman and 13 from Still Pond.

In August there were 4 cases of septic sore throat reported from Worton, all colored, 2 males and 2 females.

In July there was a case of anthrax reported from Chestertown, R. F. D., a colored man who contracted pustular anthrax from a cut on his finger inflicted while flaying a cow dead of this disease. The patient recovered.

#### MONTGOMERY COUNTY.

(Area in Square Miles, 490; Population, 32,953.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	Oetober.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Whooping cough. Chickenpox Measles Influenza Meningitis Trachoma German measles. Mumps Poliomyelitis Dysentery Malaria Septicemia Para-typhoid Erysipelas Septic sore throat	0 3 1 1 0 10 3 1 1 1 1 0 0 0 0 0 0 0	2 6 6 24 5 5 14 4 10 0 0 0 0 0 0 0 0 0 0 0 0 0 1	0 2 5 1 4 0 10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0   1   9   0   4   2   34   0   0   0   0   0   0   0   0   0	0 1 0 2 10 15 6 0 0 2 0 1 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0	1 0 0 6 4 14 11 2 0 0 0 1 1 1 0 0 0 0 0 0 0 0 0 0 0	10 11 5 19 1 28 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	6 0 1 8 2 2 0 0 0 0 0 0 1 1 1 0 0 0 0 0 0 0 0 0	21 0 0 1 0 0 0 0 0 0 1 1 1 1 0 0 0 0 0 0	$ \begin{vmatrix} 9 \\ 5 \\ 4 \\ 2 \\ 2 \\ 17 \\ 3 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	3 1 16 0 17 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	7 4 9 9 0 3 1 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	59 34 74 44 86 73 67 3 1 5 3 4 1 2 1
Total	20	66	23	53	38	30	81	21	26	42	41	25	466

Belated Returns of Typhoid Fever from Frederick City Hospital, 1915.

	1									-		1	
Mt. Ephraim	0	0	0	0	0	o	0	1	0	0	0	0	1
Clarksburg		0	0	0	0	0	0	0	1	0	. 0	0	1
Dickerson		0	0	0	0	0	0	0	1	0	0	0	1
Total	20	66	23	53	38	30	81	21	26	42	41	25	466
· ·	<u>  </u>	i		i									
Complete Total	20	66	23	53	38	30	81	22	28	42	41	25	469

In July there were 6 cases of typhoid fever reported from Damascus; in September, 4 from Silver Springs and 4 from Olney, R. F. D.

In July there were 6 cases of diphtheria reported from Chevy Chase.

In February there were 7 cases of scarlet fever reported from Chevy Chase and 6 from Bethesda; in November, 6 from Buck Lodge.

In July there were 5 cases of whooping-cough reported from Chevy Chase and 4 from Dickerson.

In October there were 14 cases of chickenpox reported from Kensington; in November, 6 from Kensington and 4 from Silver Spring.

In May there were 11 cases of measles reported from Takoma; in June, 9 from Takoma; in July, 7 from Burdett and 6 from Colesville.

In April there were 9 cases of influenza reported from Ashton and 6 from Olney.

PRINCE GEORGE'S COUNTY.

(Area in Square Miles, 486; Population, 39,443.

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

					:								
Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever. Diphtheria Scarlet fever. Whooping cough. Chickenpox Measles Mumps Influenza Septicemia Ophthalmia neonatorum Pellagra Malaria Septic sore throat Poliomyelitis Dysentery Erysipelas Meningitis German measles.	5 3 3 5 0 2 2 2 1 1 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 3 3 5 0 4 3 1 15 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3 6 5 2 8 3 0 5 2 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	34 0 2 0 4 34 34 0 0 1 0 0 0 0 0 0 0 0	7 1 3 0 3 23 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	100 0 3 1 1 1 222 0 1 1 0 0 0 2 1 1 1 0 0 0 0 0	20 1 1 1 9 0 4 4 0 0 0 0 0 1 1 0 0 1 0 1 0 0 1 1 0 0 0 1 1 0 0 0 1 0 0 0 1 0 0 0 1 0 0 0 1 0 0 0 0 1 0	37 31 31 11 00 00 00 00 00 00 00 22 14 11 11	30 6 6 6 17 0 2 2 2 0 0 0 0 0 6 6 1 0 0 0 0 0 0 0 0 0 0 0 0	8 3 7 42 2 2 4 0 0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0	13 19 2	63 93 38 118
Total	21	33	34	42	48	40	42	38	67	70	69	50	554

At Nottingham there were 3 cases of typhoid fever reported in July; in September, 7 cases from Forestville and 5 from Aquasco. In October there were 7 cases reported from Forestville.

In April there were 3 cases of scarlet fever reported from Seat Pleasant; in December, 5 from Laurel.

In October there were 9 cases of whooping cough reported from Mt. Rainier; in November, 27 cases from Mt. Rainier and 15 from Brentwood. In December there were 6 cases reported from Brentwood and 6 from Woodmoor.

In April there were 9 cases of chickenpox reported from Mt. Rainier.

In April there were 10 cases of measles reported from Forest-ville. In May, 12 cases from Ritchie, 7 from Landover and 6 from Largo. In June, 7 from Mt. Rainier, 5 from Ritchie; in July, 5 from Brentwood, 5 from Capitol Heights, 5 from Mt. Rainier and 4 from College Park.

In February there were 12 cases of influenza reported from Mt. Rainier.

In April there was one case of ophthalmia neonatorum reported from Hyattsville. The patient was a white male. In November, one case from College Park, the patient being a white male.

In September there were 13 cases of acute dysentery reported from Aquasco.

QUEEN ANNE'S COUNTY.

(Area in Square Miles, 352; Population, 16,034.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever	3	2	2	2	1	7	6	20	12	16	4	0	75
Diphtheria	0	0		$\frac{2}{0}$	0	1	0	0	1	3	5	4	14
Scarlet fever	1	$\begin{bmatrix} 2 \\ 0 \\ 5 \end{bmatrix}$	1	1 0	0	1 1 3	0	0	0	0	0	0	9
Whooping cough	0	0	0		0	3	10	5	0	0	0	0	18
Chickenpox	2 2	0	0	1	0	0	0	0	0	0	0	1	4
Measles	2	$\frac{4}{2}$	0	0	0	0	0	0	1	0	0	0	7
Mumps	6	2	3	0	0	0	0	0	-1	0	0	2	14
Erysipelas	1	0	0	1	0	0	0	0	0	0	0	0	2
Smallpox	1	0	ŏ	0	0	0	0	0	0	0	0	0	1
Influenza	0	1	0	0	0	0	0	0	0	0	0	0	1
Poliomyelitis	0	0	1 0	0	0	0	0	0	0	0	0	0	1
Malaria	0	0		1	0	0	0	2	0	0	0	0	3
Dysentery	0	0	0	0	0	0	0	0	0	2	0	1	3
Septic sore throat	0	0	0	0	0	0	. 0	0	0	1	0	0	1
Septicemia	0	0	0	- 0	0	0	0	0	0	0	-0	1	1
Total	   16	14	7	6	1	12	16	27	15	22	9	9	154

In August there were 5 cases of typhoid fever reported from Stevensville; 4, in September, from Sudlersville, R. F. D.; in October, 4 from Ford's Store.

In November there were 3 cases of diphtheria reported from Queen Anne.

In July there were 5 cases of whooping cough reported from Centreville, R. F. D.

In February there were 5 cases of scarlet fever reported from Carmichael.

In January there was one case of smallpox reported from Centreville. The patient was a colored female.

In March there was one case of poliomyelitis reported from Chester. The patient was a colored female.

ST. MARY'S COUNTY.

(Area in Square Miles, 372; Population, 16,950.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Chickenpox Measles Poliomyelitis Erysipelas Influenza Malaria Meningitis Mumps Dysentery Pellagra German measles Total	0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 1 0 0 0 49 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 8 0 35 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{bmatrix} 2 \\ 0 \\ 1 \\ 3 \\ 0 \\ 32 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	0 0 0 3 0 10 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 4 0 4 0 0 0 0 1 1 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 12 7 0 6 1 0 0 1 0 0 0 0 0 0 1 1 0 0 0 1 1 0	0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	

In October there were 3 cases of typhoid fever reported from Valley Lee.

In November there were 10 cases of diphtheria reported from Scotland.

In November there were 3 cases of scarlet fever reported, each from Ridge and California.

In April there were 8 cases of whooping cough reported from Mechanicsville.

In March there were 42 cases of measles reported from Mechanicsville and 7 from Huntersville; in April, 35 from Mechanicsville; in May, 10 from Charlotte Hall, 8 from Mechanicsville, R. F. D., and 7 from Mechanicsville.

In January there was one case of poliomyelitis reported from Great Mills; in May, one case from Clements. The former was a white female; the latter, a colored female.

In October there was one case of pellagra reported from Charlotte Hall. The patient was a colored female.

#### SOMERSET COUNTY.

(Area in Square Miles, 362; Population, 26,736.)
REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Chickenpox Measles Mumps Smallpox Influenza Dysentery Erysipelas Septicemia Malaria Meningitis Poliomyelitis Septic sore throat Total	$\begin{bmatrix} 2 \\ 2 \\ 9 \\ 24 \\ 3 \\ 0 \\ 22 \\ 20 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	5 1 0 10 10 0 0 0 0 0 0 0	$\begin{bmatrix} 1 \\ 0 \\ 3 \\ 12 \\ 4 \\ 0 \\ 0 \\ 3 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	0 0 1 1 0 3 3 0 3 6 0 0 1 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 0 4 4 0 0 23 1 0 0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{array}{c} 1 \\ 0 \\ 0 \\ 1 \end{array}$	1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	12 1 0 0 0 2 0 0 0 0 2 0 0 0 0 0 0 0 0 0	15   0   3   1   0   0   0   0   0   0   0   0   0   0	2 3 1 3 0 0 0 0 0 0 0 1 2 0 0	$egin{array}{c c} 3 \\ 7 \\ 0 \\ 2 \\ 0 \\ 0 \\ 2 \\ 0 \end{array}$	8 7 3 0 0 0 16 0 1 0 0 0 0	55 25 42 5 36 26

In July there were 5 cases of typhoid fever reported from Marion; in August, 3 from Marion; in September, 8 from Marion; in October, 3 from Marion; in November, 4 from Crisfield.

In November, there were 6 cases of diphtheria reported from Westover.

In January there were 5 cases of scarlet fever reported from Tylerton; in November, 4 from Crisfield.

In January there were 16 cases of whooping cough reported from Westover; in March, 4 from Westover.

In May there were 23 cases of measles reported from Chance; in June, 10 cases from Chance.

In January there were 12 cases of smallpox reported from Princess Anne, 5 from Marion, 1 from Crisfield and 2 from Pocomoke, R. F. D.

TALBOT COUNTY.

(Area in Square Miles, 286; Population, 19,239.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.  Typhoid fever. 1 Diphtheria 0 Scarlet fever. 1 Whooping cough. 15	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Diphtheria         0           Scarlet fever         1		3	1	1								
Chickenpox         10           Measles         0           Mumps         3           Erysipelas         1           Influenza         2           Ophthalmia neonatorum         0           Poliomyelitis         0           Septicemia         0           Dysentery         0           Pellagra         0           Total         33	1 1 4 5 0 1 0 2 0 0 0 0 0 0 0 0 1 1 1 1 1 1 1 1	3 2 2 2 0 0 0 82 1 0 0 0 0 0 0	1 2 3 0 0 0 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 7 0 0 1 1 0 0 0 1 0 0 1 0 0	2 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0	7 2 0 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	18 4 0 0 0 0 0 0 0 0 0 0 0 1 1 1 0	16 4 7 0 0 3 0 0 0 0 0 0 0 0 0 0 0	10 0 0 0 0 0 1 0 0 0 0 0 0 0	1 0 1 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	69 59 15 21 24 5 7 4 101 1 2 1 1

In July there were 3 cases of typhoid fever reported from Tilghman; in August, 5 from Easton and 4 from Trappe, R. F. D.; in September, 5 from St. Michaels, 4 from Trappe; in November, 6 from Avalon and 4 from Easton.

In May there were 7 cases of diphtheria reported from Easton; in October, 6 from Easton; in November, 5 from Wittman, R. F. D.

In September there were 3 cases of scarlet fever reported from Oxford.

In January there were 8 cases of whooping cough reported from McDaniel.

In March there was one case of ophthalmia neonatorum reported from Trappe, the patient being a white female.

In January there were 7 cases of chickenpox reported from Cordova, R. F. D.

In April there was one case of poliomyelitis reported from the Emergency Hospital, Easton, the patient being a colored female.

#### WASHINGTON COUNTY.

(Area in Square Miles, 458; Population, 51,981.) REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	Junc.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever	4	6	1	8	9	10	-4	17	15	47	17	56	194
Diphtheria	9		10		1	5	6	4	10	45	59	31	191
Scarlet fever	8	$\frac{9}{2}$	0	2 2 1	4	5	3	1	2	5	17	19	68
Whooping cough	0	7	12	1	0	15	0	0	0	1	14	0	50
Chickenpox	2		5	8	12	12	2	0	1	13	14	6	78
Measles	2 1 3	$\frac{3}{2}$	0	1	0	0	0	0	0	1	0	3	8.
Mumps	3	2	0 2 3	0	0	1 1	1	0	0	0	0	0	9
Erysipelas	3			0	0	1	0	0	0	0	1	1	9
German measles	1	0	0	0	0	0	0	0	0	0	0	1	2
Septicemia	1	0	0	0	0	0	0	1	0	0	1	0	3
Meningitis	0	-1 0	0	0	0	0		0	1	1 1	1	0	4
Influenza	0		2	4	3	1	0	0	0		1	0	12
Smallpox	0	0	0	13	16		3	0	0	0	0	0	33
Malaria	0	. 0	0	1	0	0			0	0	0	0	1
Scabies	1	0	0	0	0	0		0	0	0	0	0	1
Dysentery	0	0	0	0	0	0	0	0	0	0	1	0	1
Total	33	32	35	40	45	51	19	23	29	114	126	117	664

Belated Returns of Typhoid Fever from Frederick City Hospital, 1915.

Highfield	0	0	0	0	0	0	0	0	3	0	0	0	3
Total		í I								i i	_ 1		l
Complete Total	33	32	35	40	45	51	19	23	32	114	126	117	667

In June there were 6 cases of typhoid fever reported from Hagerstown; in August, 8 from Hagerstown and 4 from Clear-spring; in September, 8 from Hagerstown and 3 from Hancock; in October, 33 from Hagerstown and 4 from Smithsburg; in November, 10 from Hagerstown; in December, 29 cases from Hagerstown and 4 from Smithsburg.

In March there were 5 cases of diphtheria reported from Hancock; in September, 10 from Hagerstown; in October, 15 from Hancock, 14 from Hagerstown and 6 from Charlton; in November, 29 from Hagerstown, 6 from Smithsburg, R. F. D., and 4 from Smithsburg; in December, 15 from Hagerstown and 3 from Funkstown.

In January there were 4 cases of scarlet fever reported from Hancock, R. F. D., and 2 from St. James' School; in October, 4 from Hagerstown; in November, 11 from Hagerstown and 2 from Big Pool; in December, 9 from Hagerstown and 4 from Big Pool.

In March there were 11 cases of whooping cough reported from Hagerstown; in June, 4 from Smithsburg and 3 from Big Pool; in November, 9 from Hagerstown and 5 from Smithsburg, R. F. D.

In May there were 10 cases of chickenpox reported from Hagerstown; in June, 7 from Hagerstown and 5 from Funkstown; in October, 13 from Hagerstown; in November, 11 from Hagerstown and 3 from Smithsburg.

In April there were 9 cases of smallpox reported from Hagerstown, 3 from Sharpsburg and 1 from Sharpsburg, R. F. D.; in May, 7 from Sharpsburg, 4 from Sharpsburg, R. F. D.; 3 from Hagerstown, 1 from Mondel and 1 from Boonsboro, R. F. D.; in June, 1 from Hagerstown; in July, 2 from Hagerstown and 1 from Downsville, R. F. D.

#### WICOMICO COUNTY.

(Area in Square Miles, 365; Population, 28,905.) REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever	3	0	4 0	$\frac{4}{2}$	1 0	5 2 0	14 0	13 1 2	12 3	$\frac{14}{2}$	6 <b>1</b> 5	8 10	84 35
Scarlet fever	0	0	0	4 0	0	1	1 1	0	0 2	0	0	0	9
Chickenpox	$\begin{array}{c} 1 \\ 0 \\ 6 \end{array}$	0 1 6	$\begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix}$	0 1 8	$\begin{bmatrix} .0\\0\\2 \end{bmatrix}$	$0 \\ 0 \\ 1$	0 1 0	0 0	0	0 0	0	0	$\begin{array}{c} 4\\3\\25\end{array}$
Meningitis	1 0	0	0	0	0	$0 \\ 1$		0	0	0	0	0	$\frac{1}{2}$
Erysipelas	0	0	0	0	0	0	1 0	$\begin{vmatrix} 0 \\ 2 \\ 0 \end{vmatrix}$	0	0 3	0	0	1 6
Malaria Poliomyelitis Pellagra	0 0 0	0 0 0	0 0	0	0 0	0	0 0 0	0	1 0	$0 \\ 1$	0	0 0	3 1 1
Smallpox	0 	0	0	0	0	0	0	0	0	0	0		1
Total	12	7	10	20	3	10	18	18	21	21	21	19	180

In July there were 5 cases of typhoid fever reported from Pittsville, R. F. D., and 7 from Salisbury; in August, 9 from Salisbury; in September, 6 from Salisbury, 2 from Pittsville, R. F. D., and 2 from Fruitland; in October, 8 from Salisbury; in December, 4 from Salisbury.

In November there were 13 cases of diphtheria reported from Salisbury; in December, 8 from Salisbury.

In April there were 4 cases of scarlet fever reported from Parsonsburg.

In September there was one case of poliomyelitis reported from Sharptown. The patient was a white male.

In October there was one case of pellagra reported from Tyaskin. The patient was a white female.

In December there was one case of smallpox reported from Salisbury. The patient was a colored male.

#### WORCESTER COUNTY.

(Area in Square Miles, 475; Population, 22,356.)

REPORTED CASES OF NOTIFIABLE DISEASES, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Typhoid fever Diphtheria Scarlet fever Whooping cough Chickenpox Measles Influenza Smallpox Rabies Erysipelas Malaria Poliomyelitis Septic sore throat Septicemia Meningitis	3 1 0 0 1 22 1 5 0 0 0 0 0 0 0	2 0 0 0 2 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0	0 1 0 0 0 0 0	0	25 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	23 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	19 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 1 0 0 0 5 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3 1 3 1 2 1
Total	33	5	3	5	6	17	19	25	26	27	22	27	215

In July there were 3 cases of typhoid fever reported from Snow Hill, R. F. D., 2 from Pocomoke City, 2 from Snow Hill and 2 from Berlin; in August, 7 from Pocomoke City, 6 from Berlin, 3 from Snow Hill, 2 from Stockton and 2 from Showell; in September, 6 from Pocomoke City, 5 from Berlin, 3 from Showell and 3 from Berlin, R. F. D.; in October, 8 from Pocomoke City, 5 from Stockton, 2 from Snow Hill, R. F. D., 2 from Girdletree and 2 from Berlin; in November, 5 from Pocomoke City, 3 from Snow Hill, 2 from Berlin, 2 from Pocomoke City, R. F. D., and 2 from Whaleyville; in December, 4 from Snow Hill, 4 from Snow Hill, R. F. D., 3 from Pocomoke City and 3 from Bishopville.

In January there were 9 cases of measles reported from Showell, 6 from Bishop and 5 from Bishopville.

In January there were 5 cases of smallpox reported from Pocomoke City. The patients were all colored, 4 being males and 1 female. In June, 5 cases were reported from Snow Hill. The patients were all white, 2 males and 3 females.

In April there was one case of rabies reported from Bishop. The patient was a white male aged 45 years, a farmer by occupation.

In July there were 2 cases of poliomyelitis reported from Snow Hill, and in September 1 case from Ocean City.

[[u,to,L]	962 388 733 733 734 748 860 760 760 760 760 760 760 760 760 760 7	9,933 21 9,954
8itivitənninoD	оооооооноо Фооооооооо	HO H
Poot-and-Mouth Disease	<u> </u>	HO   H
ршоцэх.іД	поососососос	00 0
Septic Sore Throat	ю о ю о н о о о о и о с о ю н н н о 4 о о о н	68 0 68
Septicemia	<u> </u>	64 64
Er ysipelas	<u>оргоноиниянии 4 4 они</u>	95 0
Dysentery	рг-0-4-04-04-00-0-00-0-0-0-0-0-0-0-0-0-0-	68 C   68
German Measles	00000000000000000000000000000000000000	206
Malaria	001 F 001 0 L L 20 L Q 001 4 C 10 4 8 0 L 10 L	20 2
La Grippe	42121	854 0 854
murotono9N vimlahthqO	00000000000000000000000000000000000000	10 1
$p_{\mathcal{A}}$	000000000000000000000000000000000000000	13
Rabies	00000000000000000000000000000000000000	HO   H
xodypung	10 mm o m o co c	80 0 80 B
Landy suddal	0000000000000000000	00 0
sitignin9M	4484040004040404000000444	20 2
Cervical Adenitis	00000000000000000000	m O m
Soubies	0000000000000000000	HOTH
Poliomyelitis		80 8
sdun <sub>I</sub> g	88.0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	602
Сһіскепрох	57 50 50 50 50 50 50 50 50 50 50	984 0 984
Agnod gniqooAAI	0.52 11 1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2	906
કગુકળગુ	112 2 2 2 1 1 1 1 1 1 2 2 2 2 2 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
Searlet Fever	1 6 7 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	$\begin{vmatrix} 1,141 & 1, \\ 0 & 0 \\ \hline 1,141 & 1, \\ 1,141 & 1, \end{vmatrix}$
vironthqia	84 5 0 8 4 5 5 8 8 4 7 4 8 8 4 5 7 8 8 8 4 8 8 4 5 7 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	065,1
bionquT-nnad	000000000000000000000000000000000000000	60 6
19:19A biohqyT	21 22 23 24 25 25 25 25 25 25 25 25 25 25	2,249
xv.ujuy	00000000000000000000	HO H
County	Allegany Anne Arundel Baltimore Calvert Caroline Carroline Carroll Charles Dorchester Frederick Garrett Harford Harford Howard Kent Montgomery Prince George's Chaeen Anne's Someer Ame's Someer Ame's Somerset Talbot Washington Washington	Total

#### IV. OCCUPATIONAL DISEASES.

The reporting of occupational diseases and accidents is still in a very unsatisfactory condition. This remark applies alike to the number of cases of sickness or disability reported and to the terms by which such indispositions are designated. In a majority of instances it is a reporting of symptoms rather than of definite diseases.

In 1915 a total of 36 reports of occupational diseases and disabilities, chiefly from the Bureau of Child Labor, was received. The following table displays these cases, arranged by months and diseases.

A case of pustular anthrax is reported as occurring in March. As the report did not reach this office until after the patient had recovered, and the nature of the disease had not been verified bacteriologically, this case is not included in the list of communicable diseases.

#### OCCUPATIONAL DISEASES-MORBIDITY, 1915.

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Ophthalmic neurosis. Toxemia Infection of right hand. Occupational asthenopia. Exhaustion Pustular anthrax Bronchitis Brass poisoning. Muscular strain. Lacerated hands. Leg cut by broken glass. Irritation of respiratory tract. Miner's asthma. Incised wound of thumb. Amenorrhæa Myalgia Neurosis Heat exhaustion. Byssinosis Muscular cramps.  Total	2	0 2 1 2 2 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 3 3 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0							000000000000000000000000000000000000000	0 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 8 1 3 3 1 1 1 1 2 1 1 2 1 1 1 1 1 1 1 1 1

#### V. CASES EXAMINED FOR DIAGNOSIS.

In 1915, in the counties of Maryland, a total of 31 cases of exanthematous disease was examined for the purpose of assisting the physician or health officer in making a diagnosis. In most instances, smallpox was suspected.

Of the 31 cases of sickness examined, 27 were white persons and 4 were colored; 17 were males and 14 were females. Eight of the Maryland counties were represented.

The distribution of these cases, by months and diseases, was as follows:

Diseases.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October	November.	December.	Total.
Scarlet fever. Smallpox Diphtheria Chickenpox Foot-and-mouth disease. Mumps Typhoid fever. Pellagra Furunculosis Total	2   2   0   0   0   0   0   0   0   0   0   0	3 0 1 1 1 1 0 0 0 0	0 0 0 0 0 0 1 0 0	0 14 0 0 0 0 0 0 0	0 1 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 1 1 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 1 0 0 0 0 0	0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0	0 0 0 1 0 0 0 0 0	0 0 0 0 0 0 0	5 17 2 2 1 1 1 1 1 1 31

The following table indicates the distribution of these cases by color and sex:

Months.	White.	Colored.	Male.	Female.	Total.
January	3	1	3	1	4
February		0	5	2	7
March		0	0	1	1
April	14	0	6	8	14
May	0	1	1	0	1
June	1	0	0	1	1
August	0	1	0	1	1
September	1	0	1	0	1
November	0	1	1	0	1
Totals	27	4	17	14	31

The following table shows the distribution of these cases by counties:

Counties.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Total.
Baltimore Caroline Carroll Harford Kent Somerset Washington Worcester	0 0 1 0 1 1 1	$\begin{bmatrix} 1 \\ 1 \\ 0 \\ 4 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$	0 0 0 0 1 0 0 0	0 0 0 0 0 0 14 0	1 0 0 0 0 0 0	0 0 0 1 0 0 0 0	0 0 0 0 0 0	1 0 0 0 0 0 0	1 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 1 0 0 0 0	0 0 0 0 0 0 0	4 1 1 6 1 1 16 1
Totals	4	7	1	14	1	1	0	1	1	0	1	0	31

#### VI. PROSECUTIONS.

Dr. A. H. S., Baltimore, Md.; for failure to notify scarlet fever (3 cases), February, 1915. Guilty; fined \$10.00 and costs.

Dr. D. E. H., Hagerstown, Md.; for failure to notify chickenpox or smallpox (1 case), June, 1915. Guilty; fined \$2.00.

Dr. Wm. M. S., Towson, Md.; for failure to notify tuberculosis (1 case), July, 1915. Case dismissed.

Dr. Wm. J. R., Baltimore, Md.; failure to notify typhus fever (1 case), August, 1915.

Dr. M. L., Hagerstown, Md.; for failure to notify smallpox (1 case), September, 1915. Guilty; fined \$2.00.

· Dr. H. E. Z., Baltimore, Md.; for failure to notify typhoid fever (5 cases), September, 1915. Case dismissed.

Report made by Dr. C. W. G. Rohrer, Assistant Chief.

### Respectfully submitted,

C. Hampson Jones, Chief, Bureau of Communicable Diseases.

# Report of the Bureau of Bacteriology. WM. ROYAL STOKES, M. D., Chief Bacteriologist

I hereby respectfully report upon the work carried out in the bacteriological laboratory during the year 1915. Table No. 1 shows that there were 6,022 examinations made for physicians, and this is almost double the amount of examinations for 1914, since 3,866 examinations were made for physicians during this year. We made 5,126 examinations of water, milk and other foods, which is also an increase over the year 1914, when 3,187 such examinations were made. A total of 11,184 examinations was made, and 5,458 complete doses of antityphoid vaccine were given out to citizens of the State. The work in detail can be found in Table No. 1, which follows:

TABLE No. 1.

EXAMINATIONS FOR THE DETECTION OF DISEASES AND INSANITARY
CONDITIONS.

	COMBIL	ions.		
Positive.	Negative.	Suspicious.	Unsatisfactory.	Total.
Typhoid fever 611	813	<sup>-</sup> 88	5	1,517
Malaria 4	203	$^2$	64	273
Tuberculosis 309	888	4	4	1,205
Diphtheria 582	2,237	6	31	2,856
Miscellaneous (Examinati	ons of R	abies, Gono	coccus, Typhoid	
Blood Culture, Feces and	1 Urine Sp	pecimens)		171
Examination of Water				4,320
Examination of Milk				472
Examination of Cream				2
Examination of Ice Crean	1			1
Examination of Oysters				259
Examination of Canned G	oods			30
Examination of Tomato Pr	ulp			12
Examination of Ice				30
Total Number of Exa	ninations.			
Complete Doses of Antityp	hoid Vacci	ine Distribut	ed	5,458

In commenting on the various miscellaneous examinations made, it might be noted that the practical application of these examinations is allotted to the Bureaus of Sanitary Engineering and of Food and Drugs. Much of the bacteriological work for the former Bureau is used in the regulation of the various water supplies and sewage-disposal systems.

## REPORT OF THE CO-OPERATIVE LABORATORY FOR ALLEGANY AND GARRETT COUNTIES FOR THE MONTHS OF SEPTEMBER, OCTOBER, NOVEMBER AND DECEMBER, 1915.

	C	umber Count			lllegan County	~		Farre Coun		
· ·	P.	N.	s.	P.	$N$ . $\cdot$	s.	$\stackrel{\leftarrow}{P}$ .	N.	s.	Total.
Diphtheria (sw. from throat)	59	200	2	160	1,066	5	15	129	0	1,636
Typhoid fever (Widal test)	26	13	14	19	20	11	6	4	0	113
Typhoid fever (bile cult.,										
feces, urine and blood)	0	4	0	4	8	0	0	4	0	20
Typhoid fever (urine)	0	0	0	$^2$	$^{2}$	0	0	0	0	4
Para-typhoid (Widal test).	3	8	1	1	8	1	1	1	0	24
Malaria	0	5	0	0	2	0	0	0	0	7
Tuberculosis (sputum)	9	55	0	12	44	0	. 0	10	0	130
Tubercular meningitis (Spg.	0	_	0	0	0	0	0	_	_	
Fl.)	0	1	0	0	0	0	0	0	0	1
Control cultures	0	31	0	0	0	0-	0	0	0	- 31
	Gd.	Bd.	Sus.	Gd.	Bd.	Sus.	Gd.	Bd.	Sus.	
Milk Examinations	23	14	53	5	0	17	4	1	7	124
Water Examinations	21	14	0	12	9	1	29	19	3	108
Ice Cream Examinations	4	7	0	0	0	0	0	0	0	11
Cream Examinations	5	3	0	0	_0	0	0	0	0	8
Cond. Milk Examinations	0	1	0	0	. 0	0	0	0	0	1
Butter Examinations	0	0	0	2	0	0	0	0	0	2
Total Number Examina	tion	s		• • • •			• • • • •			2,220
Typhoid Vaccine Distributed	7				2	0	46		17	83
Smallpox Vaccine Distribute						$\overset{\circ}{0}$	1		50	<b>6</b> 1
							_		00	O1

## Respectfully submitted,

W. A. Günther,

Bacteriologist.

<sup>\*</sup>NOTE—"P," "N" and "S" in above headings represent "Positive," "Negative" and "Suspicious," respectively.

## Report of Bureau of Chemistry.

W. B. D. PENNIMAN, Chief

Dr. John S. Fulton,
State Department of Health,
Baltimore, Maryland.

DEAR SIR:

Below you will find the list of the substances analyzed by this Bureau during the year 1915. We have included in this list only those analyses made for other Bureaus. The analyses made in connection with our investigating work are not included in this list.

The law requires us to use the methods of the General Government, but we are still continuing to follow out your regulations that fundamental data regarding all methods and standards must be obtained in this laboratory as a necessary preliminary to prosecution. The principal investigations which have been carried on during the year are as follows:

Water and Sewage Analyses—This work has been completed and standard Laboratory methods adopted during the year. The plan proposed in our report of 1915 of using small containers for samples that can be sent by Parcel Post has been put into effect. In part, the results from a practical standpoint have been entirely satisfactory so far as convenience to the inspectors, accuracy of results and expense. New methods will have to be experimented with as they are proposed, and it is our intention to keep at least one trained chemist at work on this very important part of our work.

Tin in Canned Goods—A method for tin in canned goods has been devised in this laboratory.

Aspirin—A new method for the separation and estimation of this important drug has been devised.

Toxicological Work—Improvements in existing methods have been made. The more common general methods for the detection and estimation of poisonous substances have been studied and a new method suggested by our work is now under investi-

gation. The details of this method naturally require a good deal of additional study, and the work will be continued.

Investigations of minor importance have been made on the following subjects:

Evaporated Milk,
Nitrated Oils,
The Effect of Formaldehyd on Milk Serum,
Saccharin in Soft Drinks,
Canned Tomatoes,
The Amount of Sugar in Ripe and Artificially Ripened
Oranges,
Spirit of Camphor,
Chicory in Coffee.

### W. B. D. Penniman, Chemist of the Board of Health.

#### DAIRY PRODUCTS.

Butter—within the limit	25	analyses.
Butter—not within the limit	3	66
Butter—for information	3	66
	7	66
Butter—checks	1	66
Condensed Milk—within limit		66
Evaporated Milk—within limit	-2	
Evaporated Milk—for information	3	66
Evaporated Milk—checks	- 6	44
Milk—within the limit	272	66
Milk—not within the limit	43	+6
Milk—checks	40	66
MITK—checks	40	
		66 .
Human Milk	4	., .
- 1		
OILS AND FATS.		
Lard—within the limit	3	66
Lard—not within the limit	1	66
Salad Oil—within limit	1	66
	$\frac{1}{2}$	- "
Olive Oil—within limit	2	
GLANNED GOODS		
CANNED GOODS.		•
Strawberries—for information	1	46
Beaus—for information	$\frac{1}{2}$	66
as course and a second	- 1	66
Apples—for information		66
Canned Tomatoes—within limits	43	"
Canned Tomatoes—not within limits	10	**
SOFT DRINKS.		
Lemon Soda—within limits	45	66
Lemon Soda—not within limits	- 6	6.6
Sarsaparilla—within limits	40	66
Sarsaparma—within mints	10	٠.

Sarsaparilla—not within limits	3	analyses.
Ginger Ale—within 'limits	42	"
Ginger Ale—not within limits	7	44
	10	4.6
Vanilla Soda—within limits		
Orange Soda—within limits	15	44
Orange Soda—not within limits	1	
Birch Beer—within_limits	14	"
Birch Beer—not within limits	5	"
Volco—within limits	1	- 66
Wil Mar Kola—within limits	1	"
Mo Ro Soda—within limits	1	66
	1	66
Wild Cherry—within limits		
Cherry Soda—within limits	5	"
Cherry Soda—not within limits	1	
Coca-Cola—within limits	1	44
Coca-Cola—not within limits	1.	"
Grape Soda—within limits	7	44
Grape Soda—not within limits	$\dot{2}$	44
To Ko Kola—within limits	$\bar{2}$	44
		44
Cider Champagne—within limits	1	
Limko—within limits	1	"
Peach Soda—within limits	$^4$	
Chocolate Soda—within limits	$^{2}$	4.
Soda Water—within limits	1	44
Raspberry Soda—within limits	4	66
Root Beer—within limits	$\hat{6}$	46
	$\stackrel{0}{\sim} 2$	44
Cream Soda—within limits	_	44
Cream Soda—not within limits	1	"
Gayola—within limits	1	
Strawberry Soda—within limits	25	44
Strawberry Soda—not within limits	1	44
Sherbet—within the limits	1	44
Blackberry Cider—within limits	$\hat{1}$	44
Disconderly Class within minus		
VINEGARS.		
X71	004	- "
Vinegars—within the limits	681	
Vinegars—not within the limits		
	116	- "
	116 1	"
Vinegars—for information		
Vinegars—for information	1	"
Vinegars—for information	1	"
Vinegars—for information	1	"
Vinegars—for information	1 128	"
Vinegars—for information Vinegars—check  EXTRACTS.  Extract Rose—not within limits	1 128 1	"
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.	1 128 1 2	
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.	1 128 1 2 2	 
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.	1 128 1 2	  
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.  Extract Peach—not within limits.	1 128 1 2 2	   
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.  Extract Peach—not within limits.	1 128 1 2 2 1	  
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.  Extract Peach—not within limits.  Extract Raspberry—not within limits.	1 128 1 2 2 1 1	   
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Orange—not within limits. Extract Lemon—not within limits. Extract Peach—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits.	1 128 1 2 2 1 1 1 2	   
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Orange—not within limits. Extract Lemon—not within limits. Extract Peach—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits.	1 128 1 2 2 1 1 1 2 1	    
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.  Extract Peach—not within limits.  Extract Raspberry—not within limits.  Extract Strawberry—not within limits.  Extract Vanilla—not within limits.  Extract Vanilla—not within limits.  Extract Pineapple—not within limits.	1 128 1 2 2 1 1 1 2 1 2	    
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Orange—not within limits. Extract Lemon—not within limits. Extract Peach—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits.	1 128 1 2 2 1 1 1 2 1	    
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.  Extract Peach—not within limits.  Extract Raspberry—not within limits.  Extract Strawberry—not within limits.  Extract Vanilla—not within limits.  Extract Vanilla—not within limits.  Extract Pineapple—not within limits.	1 128 1 2 2 1 1 1 2 1 2	    
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits.  Extract Almond—not within limits.  Extract Orange—not within limits.  Extract Lemon—not within limits.  Extract Peach—not within limits.  Extract Raspberry—not within limits.  Extract Strawberry—not within limits.  Extract Vanilla—not within limits.  Extract Vanilla—not within limits.  Extract Pineapple—not within limits.	1 128 1 2 2 1 1 1 2 1 2	    
Vinegars—for information Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Orange—not within limits. Extract Lemon—not within limits. Extract Peach—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits. Extract Pineapple—not within limits. Extract Peppermint—not within limits. Extract Peppermint—not within limits.	1 128 1 2 2 1 1 2 1 2 1 2 1	    
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Lemon—not within limits. Extract Lemon—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits. Extract Pineapple—not within limits. Extract Peppermint—not within limits.  Extract Peppermint—not within limits.  DRUGS.  Aspirin	1 128 1 2 2 1 1 1 2 1 2 1 2 1 2 1 7	    
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Orange—not within limits. Extract Lemon—not within limits. Extract Raspberry—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits. Extract Pineapple—not within limits. Extract Peppermint—not within limits.  Extract Peppermint—not within limits. Extract Strawberry—not within limits. Extract Strawberry—not within limits. Extract Peppermint—not within limits. Extract Peppermint—not within limits.  DRUGS.	1 128 1 2 2 1 1 1 2 1 2 1 2 1 2 1 2 1 2	
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Lemon—not within limits. Extract Lemon—not within limits. Extract Peach—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits. Extract Pineapple—not within limits. Extract Peppermint—not within limits.  DRUGS.  Aspirin Fowler's solution. Landanum	1 128 1 2 2 1 1 1 2 1 2 1 1 2 1 1 2 1 1 2 1 1 1 2 1	
Vinegars—for information. Vinegars—check  EXTRACTS.  Extract Rose—not within limits. Extract Almond—not within limits. Extract Orange—not within limits. Extract Lemon—not within limits. Extract Raspberry—not within limits. Extract Raspberry—not within limits. Extract Strawberry—not within limits. Extract Vanilla—not within limits. Extract Pineapple—not within limits. Extract Peppermint—not within limits.  Extract Peppermint—not within limits. Extract Strawberry—not within limits. Extract Strawberry—not within limits. Extract Peppermint—not within limits. Extract Peppermint—not within limits.  DRUGS.	1 128 1 2 2 1 1 1 2 1 2 1 2 1 2 1 2 1 2	

#### MISCELLANEOUS FOOD PRODUCTS.

Coffee—within the limits	121	analyses.
Coffee—not within the limits	21	"
Coffee—for information	3	"
Vanilla Ice Cream—within limits	1	66
Oysters—within the limits	20	66
Whiskey—within the limits	4	44
Whiskey—within the filmits	1	66
Ground Beef—within the limits	-	66
		44
Ground Beef—not within limits	4.0	66
Ground Beef—check		66
Flour—for information		"
Honey—for information	4	"
Egg Lyn—for information	1	
Rye Flour—within limits	$^{2}$	46
Ice Cream Cones—within limits	1	66
Cornmeal—for information	1	4.6
Chickory—for information	1	66
Rye_for information	1	44
Wheat—for information	1	66
Bread—for information	1	56
Beets—for information	1	66
	1	66
Peas—for information	1	46
Beans—for information	1	"
Peanuts—for information	1	**

#### MISCELLANEOUS.

Special test for lead—Dr. W. W. Ford. Report on stone jar.

#### WATERS FOR THE YEAR 1915.

WATERS FO	, 11111	1 EMI 1010.	Special	
Waters	Water.	Scwage.	Analysis.	Ice.
Allegany County	67		56	1
Anne Arundel County	146		30	
Baltimore County	419	41	178	2
Baltimore City	7		7	1
Calvert County	94		43	
Caroline County	22		18	1
Carroll County	208	157	284	
Cecil County	45		30	1
Charles County	37		3	
Dorchester County	112		19	
Frederick County	183		104	2
Garrett County	24		25	1
Harford County	88		<b>7</b> 0	
Howard County	40		3 ~	
Kent County	108		76	3
Montgomery County	126		75	2
Prince George's County	41		20	
Queen Anne's County	31		<b>1</b> 2	1
St. Mary's County	15		3	1
Somerset County	35		31	2
Talbot County	78	6	15	2
Washington County	30 (		14	1
Wicomico County	265		18	2
Worcester County	20		18	3
Table Waters	21	• • •		
Total	2,062	204	1,152	26

## Report of the State Food and Drug Commissioner.

To the State Board of Health of Maryland.

#### GENTLEMEN:

Herewith I beg to submit my report for the twelve months ending December 31, 1915.

As shown by the schedule below, 1,910 samples of food and drugs were purchased and analyzed during the year 1915, of which number 1,701 were passed and 209 were alleged to be in violation of the Pure Food and Drugs Act.

Increased work has been performed by our Veterinarian and Dairy Inspector, whose report is hereto attached, inspection of dairies and slaughter houses during the past year being confined to the following counties: Anne Arundel, Baltimore, Carroll, Frederick, Harford, Howard, Kent, Prince George's, Queen Anne's, Somerset, Talbot, Washington, Wicomico.

The milk supply of Baltimore City being carefully looked after by the City Health Department, the inspection of these dairies and dairy farms by the State authorities is not considered necessary, as it would be a mere duplication of work without added results. It is gratifying to note a gradual improvement in the sanitary conditions of the dairies and slaughter houses in the State, although in some instances repeated visits were necessary, and even a few prosecutions, in order to impress upon the owners of the establishment the importance and necessity of conforming to the requirements of the Sanitary Inspection Law. Investigations will be continued during the present year, and it is hoped that ere long we shall be able to record a satisfactory score for every dairy and slaughter house in Maryland, especially as far as proper provision against contamination by flies, the great carriers of filth and disease, is concerned.

I beg to call your attention to the report, hereto appended, of the Cannery Inspector attached to this Department, which shows to a marked degree the satisfactory results obtained in the enforcement of the sanitary requirements for canning establishments. The employment of more inspectors and consequent expenditure of more money, with the view of bringing the canneries of Maryland up to a condition beyond criticism and second to none in the country, is absolutely necessary, and it is earnestly hoped that the Legislature now in session will find it possible to appropriate the money asked for, in order to place this, one of the leading industries of the State, in the foremost ranks in respect to sanitary conditions, both in equipment and methods.

During the past summer the crab canneries at Crisfield and Cambridge were thoroughly inspected, with the result that nearly all of them were found to be operated in compliance with the Sanitary Inspection Law. Considerable attention has also been given during the year to the conditions surrounding the shucking and canning of oysters, and while a number of establishments required special instruction and warning, the situation as a whole was found rather satisfactory.

The bacteriological examination of oysters reported below was made with the view of ascertaining possible contamination with bacteria of the colon bacillus type, the numerical valuation being made in accordance with the scoring method adopted by the American Public Health Association and the Agricultural Department at Washington. A score of less than 50 in the examination of the liquid found in five oysters in shell, taken from the same lot, or of three samples of the liquid in the case of shucked oysters, is considered at present insufficient to warrant condemnation, and naturally, the lower the score, the better the quality of the oysters. If the colon bacillus has been isolated from 1 Cc. of the liquid, a value of 1 is given that particular sample; if isolated from 0.1 Cc. a value of 10 is given, and if isolated from 0.01 Cc. a value of 100 is given, etc.

The examination of a single set of oysters in any locality does not, of course, justify any definite conclusion, as subsequent examinations may indicate better or worse conditions. Since the Health Department of Baltimore City has undertaken to investigate the oyster supply furnished in this city, all examinations of intra-urban oysters by the State Department of Health ceased with the close of the year 1915.

OYSTERS EXAMINED IN 1915.	Score for Isolated Colon Bacillus.
	or 2d Bac
	e f late on
Name and Locality.	202
Geo. W. Sheckels, 201 N. Pine St., City:	w ~
Oysters in shell from Mill Creek, Va	. 1
Shucked oysters from Mill Creek, Va	1
Oysters in shell from Eastern Bay	0
Shucked oysters from Eastern Bay	3
T. A. Robinson & Co., High and Low Sts., City:	
Shucked oysters from South River, Md	120
Shell stock	14
E. H. Burrows & Son, 514 W. Lexington St., City: Shucked oysters from West River, Md	
Shell stock	1
Shell stockOysters in shell from Patuxent River	14 0
Shucked oysters from Patuxent River	21
MacKenthun & Bach, 522 W. Lexington St., City:	
Shucked oysters from Bay Shore, West River	2,100
Shell oysters from Bay Shore, West River	32
Gerahty & McWilliams, Pratt and Hollingsworth Sts., City:	
Shucked stock from Tangier Sound	21
Shell stock from Tangier Sound	50
Oysters in shell from Mobjack Bay. Shucked oysters from Mobjack Bay.	$\frac{2}{21}$
John G. Wagner & Co., 104 W. Dover St., City:	21
Shucked stock from North End Bar, Rappahannock River	30
Shell stock from North End Bar, Rappahannock River	3
Oysters in shell from Hampton, Va Shucked oysters from Hampton, Va	$\frac{2}{12}$
Atlantic Packing Co., President and Stile Sts., City:	12
Shucked oysters from Magothy River	3
Shell oysters from Magothy River	1
W. H. Kirkwood & Sons, 114 Dover St., City:	
Shucked oysters from Water View, Va	120
Wm. B. McCaddin & Co., 26 Market Space, City:	2
Shucked oysters from Tangier Sound	0
Shell oysters from Tangier Sound	1
Oysters in shell from Tangier Sound	1
Shucked oysters from Tangier Sound	12
Shucked oysters from Old Point, Va	0
Shell oysters from Old Point, Va	1
Oysters in shell from Old Point, Va. Shucked oysters from Old Point, Va.	4
J. Geo. Pertsch, 521 W. Lexington St., City:	21
Shucked oysters from York River	1 200
Shell oysters from York River	0
Shucked oysters from Chesapeake Bay	3
Shell oysters from Chesapeake Bay	0
F. F. East & Co., Frederick and Water Sts., City: Shucked oysters from Hollins Point, Md	3
Shell oysters from Hollins Point, Md	$\frac{\mathfrak{d}}{2}$
Oysters in shell from Holland Point	1
Shucked oysters from Holland Point	3

OYSTERS EXAMINED IN 1915—Continued.	oillus
. A Tanakka	Score for Isolated
Name and Locality.	ž, T
Shall Packing Co., Atlantic Wharf, City: Oysters in shell from Patuxent River	1
Shucked oysters from Patuxent River	$2\tilde{1}$
C. L. Applegarth Co., Webster St., City:	
Oysters in shell from Patuxent River	5
Shucked oysters from Patuxent River	21
Storey & Bunnell Packing Co., foot of West St., City: Oysters in shell from Rhoad River	3
Shucked oysters from Rhoad River	1
Chas. W. Reddish, Clement and Boyle Sts., City:	
Oysters in shell from Cedar Point	1
Shucked oysters from Cedar Point	2
R. Pohler. 514 W. Lexington St., City:	0
Oysters in shell from Patuxent River	3
Wm. Heyser, Pratt and Grant Sts., City:	U
Oysters in shell from Hampton, Va	1
Shucked oysters from Hampton, Va	22
B. Hayden, 734 W. German St., City:	0
Oysters in shell from York River	$\frac{0}{3}$
Shucked oysters from York River	, 0
Oysters in shell from Herring Bay	1
Shucked oysters from Herring Bay	3
J. Langrall & Bro., Inc., 2105 Aliceanna St., City:	
Oysters in shell from Potomac River	0
Shucked oysters from Potomac River	3
H. McWilliams & Co., 110 Hughes St., City: Oysters in shell from South River	14
Oysters shucked from South River	12
W. H. Kirkwood & Son, 114 Dover St., City:	
Oysters in shell from Patuxent River	0
Shucked oysters from Patuxent River	300
T. E. Jones & Co., Cambridge, Md.: Shucked oysters from Chesapeake Bay	120
Shell oysters from Chesapeake Bay	2
W. G. Winterbottom, Cambridge, Md.:	
Shucked oysters from Annapolis, Md	1
Shell oysters from Annapolis, Md	1
I. L. Leonard & Ço., Cambridge, Md.: Shucked oysters from Potomac River, Md	1
Shell oysters from Potomac River, Md	4
H. L. Harris & Co., Cambridge, Md.:	_
Shucked oysters from Choptank and Potomac Rivers	120
Shell oysters from Choptank and Potomac Rivers	4
T. M. Bramble & Co., Cambridge, Md.:	21
Shucked oysters from Potomac and Severn Rivers  Shell oysters from Potomac and Severn Rivers	0
A. Phillips, Cambridge, Md.:	J
Shucked oysters from Choptank River, Md	1
Shell oysters from Choptank River, Md	0
M. V. Smith & Son. 310 W. Mulberry St., City:	4
Oysters in shell from Lynn Haven Bay	$\frac{1}{12}$
which offices from the first faction bufferest stressesses	

## LIST OF SAMPLES OF FOODS AND DRUGS ANALYZED FROM JANUARY 1, 1915, TO DECEMBER 31, 1915.

	Passed.	Not Passed.
Agninin Mahlota	88	3
Aspirin Tablets	27	$\frac{3}{2}$
Butter	11	$\frac{2}{2}$
Birch Beer		16
Coffee	120	
Coffee (Information)	4	0
Extract of Vanilla	1	1
Extract of Almond	1	1
Extract of Lemon	0	1
Extract of Orange	0	1
Extract of Peppermint	0	1
Extract of Peach	0	1
Extract of Pineapple	0	2
Extract of Rose	0	1
Extract of Raspberry	0	1
Extract of Strawberry	0	2
Fowler's Solution	122	12
Flour (Rye)	1	0
Hamburg Steak	21	9
Ice Cream	1	0
Ice Cream Cones	1	0
Information Samples	9	0
Jamaica Type Rum	0	1
Lard	š	$\tilde{3}$
Laudanum	83	21
Milk	245	95
Oysters (Information)		0
Olive Oil	$\overset{-\circ}{2}$	ŏ
Oleomargarine	ĩ	ŏ
Salad Oil	î	ő
Soft Drinks	221	0
Spirit of Camphor	2	ő
Spirit of Camphol	1	- 0
	41	· ·
Syrup of Iodide of Iron	1	4
Tincture Chloride of Iron	1	0
Tincture of Iodine	_	0
Vinegar	674	100
Witch Hazel	8 `	0
Whiskey	4	0
Total	1,701	209

As shown by the following list, our efforts to bring alleged violations of the Pure Food and Drug Law to trial met with better results than in former years, and the small number (25) of prosecutions may be accepted as evidence that violations of the law are annually growing less.

10	
$\overline{}$	
~	
==	
_	
rk	
$\simeq$	
N	
-	
-	
-	
-	
$\sim$	
DUR	
$\overline{}$	
$\sim$	
$\vdash$	
7	
~	
< €	
- 4	
$\vdash$	
-	
1.3	
-	
OF MARYLAND I	
Z	
-	
5-	
1	
0	
_	
-	
-	
V	
. 7	
$\vdash$	
7.5	
(J2	
75	
$\sim$	
=	
$\overline{}$	
AND DRUG	
_	
-	
4	
_	
~4	
$\vdash$	
$\overline{}$	
$\sim$	
$\perp$	
-	
-	
$\sim$	
-	
THE FOOD	
د.	
R 7	
BR 1	
ER C	
DER 1	
VDER 1	
NDER 7	
UNDER	
UNDER	
UNDER	
T UNDER	
T UNDER	
AT UNDER	
HT UNDER	
GHT UNDER	
IGHT UNDER 5	
UGHT UNDER	
DUGHT UNDER	
OUGHT UNDER	
ROUGHT UNDER	
SROUGHT UNDER	
BROUGHT UNDER	
ROSECUTIONS BROUGHT UNDER 5	
BROUGHT UNDER	
BROUGHT UNDER	
PROSECUTIONS BROUGHT UNDER	
PROSECUTIONS BROUGHT UNDER	
PROSECUTIONS BROUGHT UNDER	
PROSECUTIONS BROUGHT UNDER	
BROUGHT UNDER	
PROSECUTIONS BROUGHT UNDER	
OF PROSECUTIONS BROUGHT UNDER	
PROSECUTIONS BROUGHT UNDER	

	9	9	9	\$5.		\$10	\$5	810	ع ا	, ,	Of		~	$^{\rm of}$	è	3	\$25		\$25	5		\$5	\$25	
	Cas	q	5	of	3	of	of	of	fine of		ine			ine	•	ot	of		of	of		of	of	
	of O			fine		ne	fine	ne	, ,-		=			H H	,	nne	ne		ne.	fine		fine	ne	
	on	7	=	id	,	id H	id	d fi	paic		Dai			paid			id fi		id fi	iid		paid fine of \$5	id fi	
	siti	\$	:: :-:	Da	į	pa	ba	nai	. ;		:-			-	sts	Da	paj		paj	30:			paj	
	ous		, S S	ty;	vi.	٠. د	; ;	zi 2	. s.		guilty; paid inne of			ilty	ည် `	S	 G	ķ	ty;	<u> </u>	σ'n	ty;		ņ
	Final Disnosition of Gase.		and costs.	gui	and costs.	E)	and costs. lead guilty	and costs.  Plead guilty: paid fine of \$10	and costs.	0	gu E	,		S	\$10 and costs	ead guilty and goets	Plead guilty; paid fine of \$25	and costs.	uil	and costs. ound cuilt	and costs.	lead guilty		and costs.
	nal	7	1 T	pr	nd ,	달.	ה הקים הקים	100 200 200	ng	\$10.	Z 5	ted.		nd	9,	ב ב ב	ad p	nd	ad g	g g E	nd	7	ad g	pr
	Fi	ī	riead guilty, pard line or 45 and costs.	Plead guilty; paid fine of \$5	ت :	Plead guilty; paid fine of \$10	and costs. Plead guilty; paid fine of \$5	a Ple	a For	<del>95-</del>	Found	\$10. Stetted.		Found guilty; paid fine of	<b>≎</b> ≎-	Plead guilty; paid nne of \$5	Ple	ä	Plead guilty; paid fine of \$25	and costs.  Found guilty: paid fine of \$1	ದ	Plead guilty;	Plead guilty; paid fine of \$25	ਕ
								14	ije								ė,							
		4	ion.	Je	٠	trio E	tion.	on. trio	ion.		e e	ohi		trio	on.		Office.		offe	utte		trio	utte	
	cd.		ide in Fowler's Solution.	hir		]C	eficiency of arsenic tric	ide in Fowler's Solution. eficiency of arsenic tric	ide in Fowler's Solution		<u> </u>	nor			ide in Fowler's Solution.	eficiency of arsenic tric ide in Fewler's Solution			E.	q s		eficiency of arsenic tric	o s	
	Why Prosecuted.		So	orl		sen	sen so	$\frac{80}{2}$	So T		orī	÷.		sen	8	sen Sen	ry i		ry	ig 3		sen	ر الا	
	280.	,	$^{ m cm}_{ m S's}$	Ξ		ar '	ar ar	$^{ m er's}$	er's	-:		it 0	_;	ar	$\mathbf{s}'$ 19	ar,	ico		ico	so		ar or'e	$^{\circ}$	
	P	9		of	'n.	ot	of.		W	in landanum.	10	n. our	in laudanum.	of	M.	10 K	(F)		f ch	tter		of	tter	
	7100	,	ير الم	ıcy	laudanum.	2,5	). 10.4 10.4	E E		ıda	Ey.	taudanum Keess amo	lda	lcy	Ĕ	<u>ان</u> ال	e 0		6 О	pm		NC.V	ng.	
	=		ciei e ir	cien	apn	ciei:	e eier	e cier	e in	뎚.	nen	ncia SS	lat	eien.	∄.	cien iii	enc		enc	ess		cier	ess	
		Doffermen of engants trion	įġ	Deficiency of morphine in	ج ا	Deficiency of arsenic triox-	Deficiency of arsenic triox-	ide in Fowler's Solution. Deficiency of arsenic triox-	ide in Fowler's Solution.  Excess amount of morphine	ii	Denciency of morphine in	Landanum. Excess amount of morphine	Ë	Deficiency of arsenic triox-	<u>ک</u> ر ک	Deficiency of arsenic triox- ide in Fewler's Solution	Presence of chicory in coffee.		res	1.00		)efi	100	
62																			ы П	2 I		5 I	10	
nple	sed.	5	191	20, 1915	7	20, 1915	21, 1915	21, 1915	27. 1915	7	28, 1915	4. 1915		1915	7	crar	1915		10, 1915 Presence of chicory in coffee.	24, 1915 Process butter sold as butter.		1915 Deficiency of arsenic triox-	25, 1915 Process butter sold as butter.	
Date Sample	was Purchased.	00	Ś	20,										4,		ń	10,		10,	24,		5 <u>7</u>	25,	
Date	Pun	Ton	Jan. 20, 1919	Jan.	,	Jan.	Jan.	Jan.	Williamson's Drug Store Jan.	,	Jan.	Feb.		Feb.		ren.	Feb.		Feb.	Feb.		Apr.	Feb.	
									ore															
									Š			Ç.					Š.							
	f nt		:				ins	tein	rug	)		ä		zó.			ore					ï.		
	Name of Defendant.	- 70	מנופ	ızeı	7	īe.	wk	enst	$-\frac{\mathbf{z}}{\mathbf{z}}$	(	ໝໍ	Che		3ros	ع	j.	š		<u>.</u>	. •		ollie		
	Van	- 5	5	Ę		Ĭ	H	ebţ	son		111	ĥу		er J	7	in in	rug		age	ook		ŭ	ber	
	~ ~	-	į	Б.	2	2	رن درد	ij	ama	11		ort		uey.	Ė		Α.		Br	Ü		Ü.	We	
		Š	THOS. A. GORDEL,	obt.	Ē	<u>-</u>	nire	A. M. Lichtenstein.	Tilli	ζ	5	otw		aldn	5	e c	ssoc		Ą.	Þ.		vin	Ъ.	
		É	7	E	2	ń	52		=	•	۲	Ġ.		Fe	È	1	As		Ą	H		Ľ	J.	
	ora-	0602	000	7865 Robt. B. Frazer.	0.07	iszi S. F. Schindel.	7856 Shirey & Hawkins.	7858	7951	5	(301 A. G. HOFILLE.	7992 Clotworthy Chem. Co.		8008 Feldmeyer Bros.	9	8055 Kose Drug Co.	8042 Assoc. Drug Stores.		8041 A. A. Brager.	8066 J. W. Crook.		8316 Levin D. Collier.	8067 J. P. Weber.	
	Labora-	, r	•	[-			[-	رع	(-	ť	•	[~		00					<i>o</i> o					
			:	:		Hagerstown	:				:			:		:	:		:	:		:	:	
	0 4		:	:		:	:	:			:	:		:		п.,	:		:	:			:	
	y, Tour		:	:		II.M	pun	put	:		4	٥.		· ·	4	1001	, e		е.	е.		· Fa	ه.	
	7. T	7	H	=	4	sto	erle	erls	rick	•	W.IC	nor		poli		3110	nor		nor	nor		ury	nor	
	City, Town or	7	Oakiand	Elkton		ıgeı	Cumberland	Cumberland	Frederick		Brunswick	Baltimore		Annapolis	-	ingniandtown	Baltimore		Baltimore	Baltimore		Salisbury	Baltimore .	
		Ċ	Š	E		Ï.	Cu	Çn	F	Ė	121	Ba		7.			Ba		B	Be		ŭ	B	

# LIST OF PROSECUTIONS BROUGHT UNDER THE FOOD AND DRUGS LAW OF MARYLAND DURING 1915.

Continued.

		Ŧ	<del>J</del>	4	<del>4</del>					10	4	10
	se.	0	0	0	0					e, €,	0	€€ ±
	Final Disposition of Case.	Apr. 28, 1915 Presence of preservatives in Found guilty; paid fine of	Apr. 29, 1915 Presentatives in Found guilty; paid fine of growned beef. \$10 and costs.	Apr. 29, 1915 Presence of preservatives in Found guilty; paid fine of ground beef.	Apr. 30, 1915 Presence of preservatives in Found guilty; paid fine of	7728 American Meat Market. Nov. 27, 1914 Presence of cottonseed oil Dismissed by Grand Jury.				Apr. 16, 1914 Presence of cottonseed oil in Plead guilty; paid fine of \$5 lard.	Mar. 26, 1914 Presence of cottonseed oil in Found guilty; paid fine of	Sept. 29, 1914 Deficiency of butter fat in Plend guilty; paid fine of \$5 ice cream.
	to of	iid	tid	tiđ	uid	nd .				ı Ü	aid	ı fin
	tion	gd s	3d 'S	3d %	od s	ran				paid	g.	paic
	osi	ty;	ty;	ty;	ty;	y 0		-		· ·	ty;	, i
	Sist	ound guilty;	ound guilty;	ound guilty;	ound guilty;	٦ چ				ead guilty	ound guilty;	ead guilt; and costs.
	al J	1d	5 E	10 E	ld .	iiss		ed.	ed.	විශ <u>ව</u> පැප	ام ع	ಮರ ಆರ
	Fin	our \$18	our \$10	oni \$10	our.	isn		tett	tett	lea	mo's	lea an
		E E	- T	J U	H O			χ <u>ς</u> 	SO SO	n P	n E	n F
		ii.	ž ii	: ::	i	resence of cottonseed oil and other substances in		tine	d a	ii ii	il ii	t E
	d.	tive	tive	tive	tive	eed		in	sol	d o	q o	fa
	ute	гvа	rva	rva	rva	ons		ne	ed;	see	ısee	ter
	sec	ese	ese.	ese.	ese.	sub		iodi e.	nug	tto.	ttoı	pmq
	Why Prosecuted.	resence of pr	resence of preround beef.	resence of preround beef.	resence of pr	# #		eficiency of iod ture of iodine.	spr	cider vinegar. resence of cott lard.	00	of 1.
	'nу	9 9	100 100 100	2 G ⊆ 5 o 5	0 C	e cotho		cy f ic	Ē.	e d	e 01	eficiency ice cream
	Ħ	enc	enc	enc	enc	enc	ď.	zien re c	gar	enc G	esenc	cien
		res	res er	res gr	res	res	lard	)efic	/ine	resen Presen Jard.	res	)efi
0		10 11	νο Γ	5 F	5 I	4 F		4. H	4	4 E	4 I	4. I
npl	sed.	191	191	191	191	191		191	191	191	191	191
Date Sample	veas Purchased.	28, 38,	29,	29,	30,	27,		Jan. 14, 1914 Deficiency of iodine in tinc- Stetted. ture of iodine.	Apr. 14, 1914 Vinegar misbranded; sold as Stetted.	16,	26,	29,
ate	Pur	pr.	pr.	pr.	pr.	ον.		an.	pr.	pr.	far.	ept.
T		A	A	A	A	Z.		F	A	A	7	202
						ket						
	t.					Maı					e.	
4	Name of Defendant.	n.	nan	nan		at		son.	re.		sno.	
	fen	omi	rnr	rnr	aaf	Me		ouls	Moc	ms.	5	on.
,	De	Se	) B	BC	$\mathbf{s}_{\mathrm{ch}}$	can.		H :	C.	Ada	el J	Mas
		D.	nest	nest	is:	ieri		S	ar	J.	cha	괱
		¥.	E	E	Cp	An		Juc	Osc	24	Mi	Ъ.
	vo.	8324 W. D. Seamon.	8329 Ernest Bornman.	8337 Ernest Bornman.	8338 Chas. Schaaf.	238		6093 Juo. M. Toulson.	6869 Oscar C. Moore.	6926 R. J. Adams.	6807 Michael J. Crouse.	597
	City, Town or Labora- County. tory No.											Crisfield 7597 P. R. Mason.
}	75	Baltimore	-:	:	:	Baltimore		Salisbury	Cambridge	:	Baltimore	:
	or	:	:	:	:	:			:		:	
	y, Town or County.	:	:	:	:	:			e	:	:	:
	, Te	ore	ore	ore	ore	ore		ury	idg	Id	юге	pld
	ity C	Itim	Baltimore	Baltimore	Baltimore	ltin		lisb	mbr	Crisfield	ltin	isfie
	0	Ba	Ba	Ba	Ba	Ba		Sa	Ca	C.T.	Ba	Cr

# TOWNS VISITED BY FOOD AND DRUG INSPECTORS DURING THE YEAR 1915.

Aberdeen (3), Adamstown (2), Arlington (3), Annapolis (5), Andrews, Abels (2), Allen, Bethlehem, Bradshaw, Betterton, Birdsville, Barstow, Buena Vista, Bowens, Broomes Island, Brentwood, Boone, Buckeystown (2), Brunswick (3), Bellevue (2), Belair (4), Bethesda, Berwyn (3), Berlin (3), Bowie (2), Brooklyn, Bishop Head, Bivalve, Barton (2), Boonsboro (2), Beaver Creek, Baden (2), Brandywine (2), Birdsville, Battle Swamp, Beltsville (2), Branchville (2), Bladensburg, Blackistone, Bushwood, Bryantown, Beachville, Belle Alton (2), Beauvue (2), Baldwin (2), Bengies (2), Bar Neck, Black Rock, Butler, Chesapeake City (2), Cumberland (4), Cooksville (2), Centreville (4), Churchpeake City (2), Cumberland (4), Cooksyllle (2), Centreville (4), Churchville (2), Chestertown (3), Cockeysville (2), Corbett, Catonsville (2), Canton (2), Cambridge (4), Crisfield (4), Capitol Heights, Curtis Bay, Carney, Claiborne (2), Church Creek, Crapo, Clara, Cedarville, Croome Station (2), Conowingo (2), Colora, Cranberry Station, College Park, Clements (2), Crompton (2), California (2), Charlotte Hall, Cox's Station, Chaptico (2), Colgate, Cowenton, Chase (2), Corbett, Cob Hill, Cordova, Chester, Church Hill, Chance, Crumpton, Chaneyville, Charlotte Hall, Derwood (2), Denton (3), Deals Island (2), Delmar, Deer Park (2), Downeyville, Davidsonville, Dryden (2), Delight, Durkirk, Ellston (4) Hall, Derwood (2), Denton (3), Deals Island (2), Delmar, Deer Park (2), Downesville, Davidsonville, Dryden (2), Delight, Dunkirk, Elkton (4), Emmitsburg (2), Easton (3), East New Market (2), Elkridge (2), Ellicott City (2), Eastport (3), Ellwood (2), Essex, Edgemere, Eckhart, Edgewood, Edgewater, Funkstown (2), Frostburg (3), Frederick (6), Federalsburg, Fairfield, Fruitland (2), Fork (2), Freeland (2), Fairbank, Frizzelberg, Fallston, Forrest Hill, Fowblesburg, Fords Store, Franklinville, Fullerton, Fairlee, Friendship, Faulkner, Gaithersburg (2), Gryons Clyndon (2), Cordonville, Caldon Hill, Greet Greensboro (2), Govans, Glyndon (2), Gardenville, Golden Hill, Great Mills (2), Girdletree, Grange, Golden Ring, Glencoe, Glenarm (2), Garrison, Granite, Gorsuch Mills, Glenburnie, Galena, Galloway, Grove, Gallant Green, Hamilton (4), Hagerstown (4), Half Way (2), Havre de Grace (3), Hancock (2), Hampstead (3), Hillsboro (2), Hurlock (2), Highlandtown (2), Hyattsville (4), Hebron (2), Hopewell, Hooper Island, Harwood (2), Hughesville (2), Holly Wood (2), Hurry, Harrimansville, Hombergsville, Hyde, Hebbville (2), Hillsdale, Harrisonville (2), Hickory, Herford, Henderson, Hess, Huntington, Helen, Ironshire, Indian Head, Issue (2), Island Creek, Insley, Jesterville, Jefferson (2), Jarboeville, Jacksonville, Jarrettsville, Jessups, Kensington (2), Keymar, Kingston, Keedysville, Kingsville (2), Kennedysville, Lauraville (3), Lonaconing (2), Lisbon (2), Laurel (4), Lakesville, Linterville, Lime Kiln, La Plata (2), Leeland, Locust Grove (2), Lakeland, Leonardtown, Lothair (2), Long Green, Lewistown, Linwood, Lansdown, Lower Marlboro, Loveville, Laurel Grove, Mt. Airy (4), Myersville (2), Middletown (3), Manchester (3), Millington (2), Monkton, Mt. Winans (2), Mt. Ranier (2), Mardella Springs (2), Marion Station, Milton, Madison, Midland (2), Mason's Corner, Mitchellsville, Mulligan, Medford (2), Mt. Savage, Muirkirk, Milestown (2), Millstone, Maddox, Mechanicsville (2), Middle River (2), Mt. Washington, Monkton, Mountain Lake Park, Meadows, Malcolm, Millers Wharf, Mattox, Mapleville, Mt. Carmel, Manor, McDaniels, Marydel, Madonna, Morrisville, Maryland Line, Mar-Manor, McDaniels, Marydel, Madonna, Morrisville, Maryland Line, Marley, Millersville, Morrell Park, Magnolia, Mutual, New Market, North East (3), New Windsor (2), Newark (2), Neavitt, Nanticoke, Nottingham, Naylors, North Keys, Newberg (2), Newport, North Keys, Oxford (2), Overlea (3), Oakland (3), Ocean City (2), Owings Mills (2), Oakwood, Odenton, Owing Station, Oakville, Oraville, Popes Creek, Perryville (2), Port Deposit (3), Poplar Springs (2), Pikesville (2), Preston (2), Parkton (2), Princess Anne (2), Pocomoke City (2), Parkville (2), Parsonsburg (2), Pittsville (2), Port Tobacco, Pomonkey, Pomport, Perry Hall (2), Perryman (3), Piney Point (2), Pearson (2), Park Hall, Point Ridge, Phoenix, Poplar (2), Poplar Heights, Pumphrey, Prince Frederick, Port Republic, Palmer, Queenstown (2), Queen Anne's (3), Quantico, Raspeburg (2), Rising Sun (2), Ridgeville (2), Rockville (2), Reisterstown (3), Ridgeley (3), Royal Oak (2), Riderwood (2), Riverton, Riverdale (2), Rowlandville, Redgate (2), Rock Point, Rosedale (2), Rossville (2), Ruxton, Rockdale, Randallstown (2), Relay, Sykesville (3), Sudlersville, Sharpstown, Salisburg (4), Snow Hill (3), Seat Pleasant, St. Mary's City, St. Inigoes, St. George, Scotland (2), St. Helena, Sweet Air, Shawsville, Shipley, Severn, Still Pond, St. Leonards, Solomon's Island, Sherwood, Smithsburg, St. Michaels (3), Sparrows Point (2), Sharpsburg, Stockton, Sparks, Silver Springs, Stevensville, Taneytown (2), Thurmont (3), Trappe, Towson (3), Tilghman's (2), Taylor's Island, Toddville, Tyaskin, Tilghmanton, T. B. Talbot, Tomkinsville, Turner, Texas, Timonium, Union Bridge (3), Upper Marlboro (2), Uniontown, Upperco, Upper Falls, Valley Lee (2), Vienna, Williamsport (2), Westernport (2), Woodsboro, Walkersville, Westminster (3), Westport (2), Westover (2), Winona (2), Wingate, White Haven, Whaleyville (2), Williards, Westwood (2), Woodlawn (2), Washington Grove, Wakeville, Woodenberg, Winchester, Warren, Walston, Whitemarsh, Worton, White Plain.

During the year 1915, 265 hearings were held in connection with alleged violations of the law, as follows: 16 in January, 23 in February, 36 in April, 30 in May, 16 in September, 16 in October, 99 in November and 29 in December.

Besides repeated inspections of markets, stores and wharves in Baltimore City, 653 towns were visited by the inspectors, resulting in 10,805 inspections of places where food products are manufactured, stored or sold, more especially with the view of correcting insanitary conditions. This does not include the inspections made by the Cannery and Dairy inspectors, which are separately reported upon elsewhere.

# INSPECTIONS MADE IN 1915.

Butter Inspections	70
Bottling Plant	1
Bakeries	69
Crab Houses	43
Canned Goods Inspections	4
Depots	4
Fruit Exchanges	1
General Stores	8.450
Ice Cream Plants	118
Markets	390
Oyster Inspections	202
Produce Yards	4
Sanitary Inspections	358
Special Inspections	49
Soda Water Inspections	10
Terminal Warehouses	78
Wharves	954
	- JOX
Total	10.805

The following lists will show the character and quantity of food products condemned and destroyed by the inspectors of this Department during the past year, and also the number of samples of food products and drugs examined and not passed after analysis.

# CONDEMNATIONS OF FOOD PRODUCTS FROM JANUARY 1, 1915, TO DECEMBER 31, 1915.

Apples	1,908 gal. cans.
Bananas	2 bunches.
Beef	5½ lbs.
Butter, Peanut	2 lbs.
Beans	972 cans.
Chickens	18 lbs.
Currants	282 lbs.
Cantaloupes	1 basket & 7 crates.
Carrots	1 bushel.
Cherries	15 quarts.
Cranberries	97,831 lbs. (98 barrels).
Cabbage	40 lbs.
Dates	20 lbs.
Flavoring Extracts	8 bottles.
Fish	680 lbs.
Grapes	12 lbs.
Ham	7 lbs.
Horse Radish	4 bottles.
Haddock	1,400 lbs.
Hake	500 lbs.
Liver	15 lbs.
Lettuce	1 basket.
Laudanum	64 ounces.
Milk, Condensed and Evaporated	280 cans.
Onions	5½ bushels.
Oranges	225 lbs.
Pudding	8½ lbs.
Peaches	2 baskets.
Prunes	15 lbs.
Potatoes	18 bushels.
Pumpkin	3 lbs.
Rabbits	21
Raisins	94 lbs.
Sausage	20 lbs.
Sweet Potatoes	100 lbs.
Strawberries	53 quarts.
Syrup Iodide of Iron	$1\frac{1}{2}$ pints.
Turnips	½ bushel.
Tomatoes	2,139 cans.
Tomato Pulp	38,941 cans.

# LIST OF FOODS AND DRUGS NOT PASSED DURING 1915.

	I	Representation. Cause for Rejection.
3	samples	Aspirin TabletsAdulterated with large quantities of
		foreign material.
2	samples	ButterMisbranded; renovated butter.
2	samples	Birch BeerAdulterated; contain saccharin.
16	samples	CoffeeAdulterated; contain chickory.
12	samples	Flavoring ExtractsMisbranded; artificial product.
12	samples	Fowler's SolutionAdulterated; deficient in arsenic tri-
	•	oxide.
9	samples	Hamburg SteakAdulterated; contain preservative.
1	sample ?	Jamaica Type RumMisbranded; imitation product.
3	samples	LardAdulterated; mixture of cottonseed
_		oil and other substances.
21	samples	LaudanumAdulterated; deficient in morphine.
95	samples	MilkLow in butter fat and bacteriolog-
		ically bad.
2	samples	Syrup Iodide of IronContain excessive amount of ferrous
	,	iodide.
2	samples	Syrup Iodide of IronAdulterated; deficient in ferrous
	1	iodide.
79	samples	VinegarAdulterated; deficient in acetic acid.
21	samples	VinegarMisbranded; sold as cider vinegar
	`	when not cider vinegar.

It is most unfortunate that as yet the Legislature at present in session has taken no favorable action on our application for a special appropriation to continue the work of sanitary inspection of the canneries of this State, and should the Legislature fail to make an appropriation, the work, so successfully inaugurated last year, will be seriously crippled. It is absolutely impossible to continue successful inspection of the large number of canneries in this State with the services of but one man at our disposal, since frequent visits during the short season are necessary and this would require at least five or six additional inspectors to be stationed in different parts of the State.

The Maryland Canners' Association has promised us their support in this matter, and I have no reason whatever to doubt their sincerity. This makes me somewhat hopeful of a final favorable outcome in our efforts to conserve one of the greatest industries in the State of Maryland, and if sufficient aid is not forthcoming from the State Treasury, I fear that the canning industry will suffer seriously.

Below please find the annual report of our Meat Inspector, whose activities during the many years of his connection with the State Board of Health have undoubtedly been of considerable value to the public.

### REPORT OF MEAT INSPECTOR FOR THE YEAR 1915.

Inspected at slaughter-houses and abattoirs — 84,378 cattle, 33,716 calves, 145,286 sheep, 203,587 hogs. Visited 2,756 slaughter-houses. Condemned:

Beef, tainted	450 lbs.
Beef, corned	450 lbs.
Beef, tubercular	10,125 lbs.
Calves, under age and weight	37
Cows, tubercular	6
Hogs, cholera	13
Mutton	1,400 lbs.
Veal	700 lbs.
Veal and Mutton	310 lbs.

CHARLES N. MITTEN, Inspector.

I also desire to call your attention to the appended letter of the Department's chemist and to the report of examinations of canned tomatoes made in the State Laboratory. The laboratory work of this investigation was performed by Clifford O. Miller, assistant chemist, whose thorough and very creditable work has supplied us with much valuable information. Printed copies of this report are being mailed to all the canners of this State, and those whose products appear in Group 3 will receive special letters notifying them of the conditions found in the canned tomatoes obtained at their establishments.

MARCH 6, 1916.

Dear Sir:

I am sending you herewith a statement of the results obtained, during the past six months, in the chemical examination of packed tomatoes.

In accordance with your suggestion, we have divided the samples analyzed into three groups:

Group I contains all those which apparently are not adulterated through the addition of water.

Group II are those tomatoes which fall below the standards adopted, but whose aduleration may not have been intentional, and may have arisen from the condensation of steam used in the packing processes.

Group III contains the samples in which the extent to which water has been added excludes the possibility of such addition having been accidental.

With respect to the tentative standards suggested by you, we may say that their fairness is fully confirmed by our analytical results, there being no case where the intentional addition of water was charged by our Inspectors, and in which this charge was not fully established in the laboratory. These standards are as follows:

Can contents pulped:—Total solids not less than 5%.

Filtered juice:—Specific Gravity—(25° C.)—not less than 1.0190.

- —Refractometer reading—(20° C.)—not less than 31.0.
- -Total solids-not less than 4.0%.
- -Reducing sugars—not less than 2.0%.

It will be noticed that there are really two sets of standards: one having to do with the amount of solid material, which should be and is naturally present in properly packed tomatoes, while the other has to do more particularly with the physical and chemical constants of the juice of tomatoes. We are of the opinion that the standard of 6.4% solids, suggested by the U. S. Government, is too high for the 1915 pack. The minimum of 5% is perhaps a little low, and therefore may in fairness be required of the packers, particularly if the examination of the filtered juice yields results which fall below the standard figures set forth above.

The drained solids vary greatly in amount, depending to a large extent upon the methods of packing. They are most likely lower in plunger-packed goods than in hand-packed; also in ripe or over-ripe tomatoes than in green ones. In addition to adulteration through addition of water, the occasional presence of green or of soft fruit was noted. The ash also varies, depending upon the amount of seeds present, and upon whether salt has been added. The amount of sugar in green tomatoes is lower than in ripe tomatoes, and a similar low sugar content is found when the tomatoes were over-ripe and had begun to ferment.

We also suggest for your consideration the possibility of fixing a limit for the acid present. Excessive acid generally

indicates that the tomatoes were over-ripe when packed, and had begun to ferment; however, a high acid content may be due in some instances to an excessive amount of seeds present.

Yours very truly,

W. B. D. PENNIMAN, Chemist, State Dept. of Health.

CHAS. CASPARI, JR.,

State Food and Drug Commissioner.

# RESULTS OF EXAMINATIONS OF CANNED TOMATOES. GROUP I.

GROUP 1.													
Vo.		rids 19.		ENTS OF				F	LITERE	d Juice			
Laboratory No.	Net Weight.	Per Cent. Solids by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Aeid-Sugar Ratio.
10,072	34.74	33.00	5.73	0.430	0.032	4.30	0.410	0.037	32.80	1.0202	0.398	2.68	1:6.7
10,064	35.62	50.49	5.63	0.480	0.030	4.49	0.408	0.017	34.20	1.0207	0.365	2.64	1:7.2
10,066	33.29	54,50	5.42	0.430	0.004	4.23	0.380	0.008	33.05	1.0195	0.309	2.51	1:8.1
9,802	18.69	16.98	5.80	0.344	0.0	4.29	0.390	0.00	33.00	1.0210	0.421	2.02	1:4.3
10,076	10.58	61.66	5.54	0.405	0.016	4.30	0.403	0.009	32.45	1.0203	0.360	2.84	1:7.9
8,975	19.57	45.04	5.15	0.375	0.0	4.01	0.243	0.0	32.65	1.0202	0.376	2.25	1:6.0
8,975	20.10	45.59	5.25	0.460	0.010	4.09	0.331	0.006	31.25	1.0200	0.386	2.23	1:5.7
8,861	34.39	36.41	5.40	0.237	0.004	4.08	0.301	0.003	33.31	1.0206	0.384	2.65	1:6.9
9,819	34.39	42.57	5.19	0.379	0.009	4.12	0.264	0.0	32.05	1.0199	0.337	2.70	1:8.0
9,810	29.98	59.41	5.10	0.314	0.0	3.98	0.344	0.0	32.15	1.0197	0.405	2.28	1:5.6
9,807	35.27	38.50	5.05	0.32	0.002	4.13	0.270	0.0	33.00	1.0199	0.324	2.66	1:8.2
10,083	32.62	43.31	5.36	0.470	0.005	4.26	0.450	0.017	33.20	1.0212	0.384	2.48	1:6.4
10,085	33.15	35.63	5.54	0.430	0.027	4.31	0.410	0.031	33.10	1.0204	0.390	2.19	1:5.6
10,092	20.45	35.34	5.75	0.467	0.025	4.90	0.413	0.025	35.68	1.0238	0.350	3.29	1:9.5
9,824	33.86	55.20	5.32	0.372	0.006	4.20	0.355	0.003	34.30	1.0214	0.625	2.36	1:3.8
10,097	34.38	40.10	5.83	0.368	0.002	4.34	0.379	0.003	34.00	1.0212	0.430	2.43	1:5.6
10,089	34.56	37.24	5.64	0.966	0.420	4.60	1.06	0.510	34.18	1.0234	0.404	2.14	1:5.5
9,813	34.39	31.79	5.33	0.437	0.028	4.26	0.413	0.028	33.15	1.0210	0.374	2.16	1:5.8
9,803	18.69	16.98	5.50	0.292	0.004	4.05	0.228	0.006	33.10	1.0199	0.350	2.67	1:7.6
10,071	33.29	47.09	5.17	0.483	0.040	3.98	0.425	0.040	32.65	1.0200	0.326	2.41	1:7.3
10,078	32.62	43.24	5.32	0.440	0.018	4.43	0.430	0.017	33.15	1.0211	0.380	2.93	1:7.7
9,822	35.09	42.28	5.01	0.390	0.004	3.99	0.288	0.006	33.00	1.0205	0.317	2.85	1:9.0
8,973	33.12	45.74	5.33	0.439	0.02	4.06	0.161	0.0	32.50	1.0202	0.363	2.55	1:7.0
9,816	34.74	36.46	5.83	0.393	0.025	4.37	0.410	0.041	33.80	1.0213	0.392	2.55	1:6.5
9,820	34.21	44.34	5.18	0.411	0.010	4.08	0.333	0.0	35.00	1.0222	0.370	3.03	1:8.2
10,086	31.74°	44.44	6.37	0.590	0.024	5.12	0.400	0.028	36.65	1.0242	0.403	3.24	1:8.4

# GROUP I-Continued.

No.		Solids ning.		ENTS OF				F	ILTERE	d Juice			
Laboratory No.	Net Weight.	Per Cent. Solid by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Acid-Sugar Ratio.
8,873	34.09	38.38	5.38	0.480	0.001	4.22	0.309	0.002	33.33	1.0212	0.470	2.41	1:5.1
10,065	32.27	43.95	5.09	0.620	0.140	3.98	0.520	0.110	31.35	1.0196	0.359	2.12	1:5.9
9,812	20.63	40.17	5.28	0.25	0.02	4.25	0.19	0.005	32.35	1.0206	0.246	2.17	1:8.8
9,837	33.33	35.34	4.82	0.263	0.0	4.21	0.342	0.0	31.38	1.0197	0.351	2.30	1:6.5
9,817	33.50	55.26	5.52	0.336	0.027	4.30	0.280	0.017	33.55	1.0209	0.350	2.70	1:7.7
9,818	34.74	40.61	5.94	0.380	0.022	4.53	0.440	0.009	34.25	1.0228	0.443	2.87	1:6.5
9,838	33.86	40.10	5.11	0.234	0.004	4.17	0.278	0.003	31.40	1.0194	0.355	2.22	1:6.6
9,805	20.63	39.31	5.18	0.382	0.018	3.99	0.222	0.0	32.28	1.0196	0.382	2.28	1:5.0
9,841	34.21	42.80	5.11	0.418	0.026	3.98	0.330	0.017	32.00	1.0201	0.350	2.40	1:6.9
10,095	19.38	44.54	6.12	0.413	0.037	4.90	0.456	0.039	36.70	1.0233	0.330	3.06	1:9.3
10,087	32.27	66.36	5.75	0.340	0.00	4.56	0.270	0.0	34.35	1.0217	0.360	3.19	1:8.6
9,804	20.81	27.12	5.92	0.278	0.0	4.61	0.189	0.0	34.90	1.0220	0.324	3.11	1:_9.6
10,075	20.11	36.00	5.25	0.45	0.017	4.18	0.417	0.022	32.55	1.0205	0.380	2.40	1:6.4
10,067	33.29	56.61	5.47	0.370	0.034	4.16	0.290	0.020	32.60	1.0199	0.322	2.80	1:8.7
10,077	102.29	39.14	5.10	0.430	0.015	4.14	0.430	0.014	33.00	1.0202	0.390	2.32	1:6.0
9,809	33.86	39.58	5.02	0.24	0.0	4.29	0.372	0.0	32.00	1.0190	0.360	2.24	1:6.2
9,806	33.65	39.78	5.96	0.37	0.009	4.67	0.340	0.009	34.80	1.0222	0.417	3.02	1:7.2
9,811	33.50	33.25	5.08	0.411	0.016	3.99	0.332	0.020	32.00	1.0202	0.376	2.40	1:6.8
9,814	34.56	41.83	6.21	0.401	0.007	4.54	0.282	0.0	34.55	1.0240	0.381	2.96	1:7.7
9,800	20.11	28.07	5.26	0.525	0.011	4.36	0.544	0.036	33.84	1.0219	0.418	2.50	1:6.0
9,842	34.39	40.00	5.03	0.480	0.006	3.98	0.292	0.0	32.10	1.0197	0.357	1.94	1:7.6
9,821	34.74	40.99	5.04	0.412	0.004	4.40	0.404	0.0	32.40	1.0194	0.307	2.27	1:8.0
9,808	34.56	35.71	4.94	0.257	0.0	4.37	0.182	0.0	31.45	1.0192	0.299	2.59	1:8.6
9,823	33.86	48.18	5.38	0.394	0.007	4.21	0.226	0.006	33.60	1.0214	0.370	2.90	1:7.8
9,843	32.45	34.33	5.11	0.320	0.0	4.55	0.280	0.0	32.05	1.0193	0.420	2.08	1:5.0
9,815	38.69	36.12	6.14	0.47	0.0	4.62	0.350	0.0	35.00	1.0227	0.426	1.85	1:4.3
9.801	20.11	32.45	5.07	0.406	0.038	4.04	0.353	0.004	31.95	1.0193	0.405	1.87	1:4.6

GROUP II.

CAOOT II,													
Vo.		lids	CONT	PULPE				I	FILTERE	d Juice	•		
Laboratory No.	Net Weight.	Per Cent. Solids by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Acid-Sugar Ratio.
10,090	19.93	35.40	4.75	0.530	0.019	3.68	0.420	0.011	30.50	1.0184	0.400	1.80	1:4.5
10,068	33.86	40.10	4.56	0.440	0.030	3.71	0.350	0.031	30.90	1.0183	0.280	2.23	1:8.0
10,088	32.62	50.80	5.19	0.337	0.010	3.87	0.350	0.014	32.00	1.0188	0.330	2.18	1:6.6
9,850	33.69	47.64	5.20	0.310	0.020	3.72	0.333	0.023	31.20	1.0187	0.416	2.17	1:5.2
9,840	33.29	43.90	5.00	0.270	0.0	3.91	0.224	0.0	31.20	1.0188	0.405	2.40	1:5.9
10,093	18.51	44.76	4.98	0.415	0.015	3.88	0.430	0.003	30.20	1.0192	0.340	2.57	1:7.5
9,858	34.74	50.50	4.46	0.290	0.0	3.60	0.326	0.0	30.20	1.0183	0.330	2,12	1:6.6
8,976	19.75	35.71	4.91	0.359	0.0	3.72	0.404	0.0	31.30	1.0185	0.377	1.74	1:4.6
8,976	20.63	47.00	4.95	0.380	0.0	3.72	0.290	0.0	31.30	1.0184	0.341	2.09	1:6.1
8,977	34.92	30.00	4.66	0.402	0.010	3.79	0.310	0.002	31.55	1.0186	0.371	1.87	1:5.1
8,974	20.10	45.61	4.53	0.440	0.011	3.62	0.280	0.0	31.20	1.0187	0.350	1.90	1:5.4
8,974	20.63	38.44	4.97	0.450	0.002	3.89	0.280	0.0	31.58	1.0184	0.327	2.16	1:6.6
8,862	33.86	39.90	5.04	0.309	0.008	3.83	0.387	0.005	31.50	1.0189	0.258	2.54	1:9.8
9,859	33.86	43.73	4.72	0.419	0.006	3.70	0.308	0.003	31.60	1.0191	0.350	2.11	1:6.0
9,860	34.74	33.33	4.74	0.349	0.0	3.59	0.306	0.0	30.90	1.0184	0.405	2.08	1:5.1
10,063	32.45	36.23	4.55	0.570	0.210	3.77	0.380	0.110	31.15	1.0186	0.380	1.91	1:5.0
8,903	33.86	33.16	4.80	0.290	0.0	3.79	0.340	0.0	31.10	1.0189	0.341	2.12	1:6.2
9,847	33.33	34.39	4.88	0.384	0.008	3.72	0.412	0.011	31.15	1.0189	0.352	1.90	1:5.4
9,848	37.39	33.00	4.51	0.419	0.012	3.63	0.420	0.023	30.80	1.0181	0.349	2.00	1:5.7
9,891	20.11	35.09	4.86	0.470	0.013	3.52	0.409	0.012	30.88	1.0184	0.389	1.98	1:5.1
9,856	33.50	40.00	5.01	0.357	0.0	3.49	0.356	0.0	30.90	1.0181	0.360	2.05	1:5.7
10,070	32.27	37.36	4.54	0.430	0.027	3.66	0.340	0.006	31.35	1.0186	0.340	2.06	1:6.6
9,854	19.22	52.29	4.63	0.280	0.011	3.55	0.270	0.0	30.85	1.0184	0.280	2.36	1:8.2
8,873	32.45	33.33	3.98	0.320	0.010	3.66	0.200	0.0	30.85	1.0189	0.430	2.06	1:4.7
8,874	33.68	30.89	5.02	0.406	0.006	3.83	0.320	0.002	31.42	1.0184	0.410	2.34	1:5.6
8,874	32.09	36.95	5.21	0.370	0.002	3.90	0.174	0.0	30.70	1.0186	0.431	2.38	1:5.5

# REPORT OF THE

GROUP II—Continued.

Vo.	-	tids 19.		ENTS 01 PULPER				F	TILTERE	d Juice.				
Laboratory No.	Net Weight.	Per Cent. Solids by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Acid-Sugar Ratio.	
9,839	33.15	40.00	5.00	0.204	0.0	3.87	0.161	0.0	31.60	1.0193	0.324	2.22	1:6.8	
10,073	19.93	39.82	4.95	0.370	0.021	3.83	0.340	0.022	31.15	1.01855	0.402	2.16	1:5.0	
9,846	33.33	33.86	4.79	0.361	0.0	3.68	0.219	0.0	30.72	1.0182	0.357	2.28	1:6.4	
9,857	33.86	42.70	4.57	0.332	0.009	3.61	0.265	0.0	31.10	1.0194	0.447	2.08	1:4.6	
9,851	32.98	33.68	4.77	0.401	0.011	3.65	0.419	0.018	30.65	1.0183	0.338	2.12	1:6.3	
9,855	33.15	35.63	4.98	0.374	0.005	3.71	0.334	0.003	31.25	1.0187	0.420	2.16	1:5.0	
9,849	35.27	41.50	4.86	0.590	0.018	3.65	0.428	0.0	30.80	1.0186	0.350	1.93	1:5.4	
9,844	19.05	´8.33	4.67	0.450	0.033	3.69	0.460	0.058	31.00	1.0188	0.477	1.73	1:3.6	
9,852	32.62	31.01	4.95	0.501	0.005	3.83	0.403	0.003	31.90	1.0198	0.554	1.86	1:3.4	
8,978	104.72	42.91	5.01	0.410	0.006	3.93	0.220	0.0	31.20	1.0188	0.317	2.31	1:7.3	
8,978	105.53	46.16	4.74	0.320	0.006	3.85	0.330	0.002	30.98	1.0182	0.348	2.20	1:6.3	
8,897	33.86	33.26	5.19	0.237	0.0	3.89	0.240	0.003	31.60	1.0190	0.250	2.14	1:8.6	
10,079	33.47	45.78	5.08	0.392	0.031	3.78	0.380	0.040	30.55	1.0185	0.380	1.57	1:4.1	
8,894	20.45	37.04	4.97	0.38	0.0	3.78	0.21	0.006	30.41	1.0187	0.392	2.12	1:5.4	
9,845	33.33	34.39	4.95	0.355	0.0	3.72	0.251	0.0	30.75	1.0187	0.353	2.12	1:6.0	
9,237	20.28	37.39	4.76	0.200	0.003	3.85	0.310	0.002	31.75	1.0184	0.298	1.73	1:5.8	
10,098	33.33	38.62	4.82	0.321	0.005	3.71	0.316	0.008	30.90	1.0182	0.356	2.12	1:6.0	
10,096	33,15	40.42	4.87	0.361	0.0	3.81	0.397	0.003	30.85	1.0190	0.340	2.05	1:6.0	
9,853	34.74	36.36	4.82	0.438	0.010	3.43	0.300	0.0	30.55	1.01805	0.330	2.03	1:6.1	

GROUP III.

No.		lids 19.	CONT	ENTS OF	F CAN,			I	FILTERE	D JUICE			
Laboratory No.	Net Weight.	Per Cent. Solids by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Acid-Sugar Ratio.
9,867	19.40	50.90	4.84	0.323	0.007	3.43	0.225	0.009	29.48	1.0162	0.328	1.83	1:5.6
9,871	19.75	47.32	4.90	0.320	0.003	3.55	0.395	0.015	30.21	1.0176	0.367	2.00	1:5.4
9,884	32.27	45.05	4.28	0.440	0.006	3.55	0.340	0.003	30.05	1.0176	0.320	1.99	1:6.2
9,074	19.75	41.98	4.34	0.450	0.019	3.23	0.380	0.002	29.25	1.0172	0.410	1.29	1:3.1
9,875	19.75	37.50	4.45	0.310	0.0	3.34	0.330	0.0	29.45	1.0174	0.480	1.45	1:3.0
9,865	20.11	37.72	4.64	0.459	0.028	3.52	0.230	0.006	29.95	1.0166	0.325	2.17	1:6.7
9,881	32.45	16.13	5.08	0.520	0.025	3.30	0.478	0.023	29.75	1.0185	0.438	1.46	1:3.3
9,888	34.74	33.00	4.36	0.295	0.008	3.41	0.187	0.0	30.00	1.0177	0.336	2.16	1:6.4
9,882	34.39	38.46	4.26	0.338	0.0	3.24	0.284	0.0	28.50	1.0156	0.240	1.95	1:8.1
8,974	20.11	43.86	4.44	0.265	0.003	3.24	0.250	0.003	29.38	1.0165	0.332	1.63	1:4.9
8,977	33.86	27.46	4.35	0.240	0.004	3.45	0.408	0.017	29.55	1.0170	0.381	1.79	1:4.7
9,866	20.28	33.91	4.47	0.339	0.003	3.39	0.287	0.0	28.95	1.0166	0.339	1.88	1:5.5
9,880	33.33	37.56	4.24	0.308	0.01	3.42	0.313	0.015	29.40	1.0170	0.312	1.90	1:6.1
10,082	32.62	51.89	4.18	0.320	0.026	3.51	0.330	0.009	29.92	1.0171	0.314	1.98	1:6.3
9,879	34.56	31.12	4.37	0.297	0.0	3.44	0.385	0.0	29.90	1.0172	0.392	2.72	1:6.9
8,898	34.57	32.47	4.08	0.357	0.004	3.18	0.357	0.020	29.05	1.0164	0.433	1.57	1:3.6
8,898	34.56	35.71	4.10	0.350	0.0	3.16	0.271	0.0	28.40	1.0160	0.404	1.59	1:3.9
8,899	32.27	9.62	2.50	0.258	0.0	1.70	0.186	0.0	21.70	1.0095	0.213	0.79	1:3.7
9,885	34.74	35.35	4.30	0.450	0.009	3.33	0.370	0.0	29.55	1.0169	0.305	1.75	1:5.7
9,872	33.86	34.37	4.87	0.402	0.012	3.82	0.320	0.0	30.90	1.0174	0.393	2.22	1:5.6
8,902	32.28	28.41	4.32	0.444	0.024	3.36	0.460	0.006	29.85	1.0174	0.367	1.81	1:4.9
8,902	33.65	37.17	4.37	0.475	0.020	3.40	0.310	0.005	29.45	1.0171	0.351	1.83	1:5.2
8,901	33.86	33.33	4.50	0.264	0.0	3.36	0.370	0.0	29.85	1.0171	0.393	1.64	1:4.2
8,903	34.56	36.73	4.69	0.320	0.0	3.56	0.360	0.002	30.60	1.0181	0.350	2.02	1:5.7
9,864	19.05	33.33	4.52	0.502	0.051	3.48	0.389	0.017	29.88	1.0173	0.380	1.95	1:5.1

# REPORT OF THE

GROUP III—Continued.

													-
No.		lids 19.		ENTS OF		,		F	LILTERE	D JUICE.			
Laboratory No.	Net Weight.	Per Cent. Solids by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Satt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Acid-Sugar Ratio.
8,973	32.27	29.73	4.30	0.390	0.004	3.46	0.230	0.005	29.82	1.0175	0.350	1.80	1:5.2
8,973	32.27	39.56	3.66	0.247	0.0	2.83	0.307	0.0	26.92	1.0146	0.297	1.60	1:5.6
9,877	32.09	42.39	4.09	0.298	0.0	3.21	0.299	0.0	28.75	1.0159	0.272	1.82	1:6.7
9,890	32.27	35.13	4.26	0.377	0.006	3.44	0.292	0.0	29.82	1.0176	0.390	2.06	1:5.3
9,886	32.45	30.64	4.51	0.388	0.006	3.22	0.374	0.0	30.15	1.0172	0.334	0.26	
9,873	33.86	37.50	4.71	0.334	0.010	3.55	0.260	0.004	30.50	1.0175	0.338	2.20	1:5.5
9,878	31.92	7.18	4.01	0.366	0.0	3.40	0.355	0.0	29.42	1.0170	0.427	1.90	1:4.4
9,863	18.87	35.51	4.57	0.501	0.020	3.58	0.282	0.011	30.72	1.0168	0.406	2.01	1:5.0
9,889	34.56	43.87	4.10	0.236	0.002	3.26	0.275	0.003	29.95	1.0170	0.318	2.07	1:6.5
10,080	32.27	36.06	4.85	0.385	0.008	3.56	0.394	0.009	30.30	1.0180	0.333	2.56	1:7.6
9,861	20.11	31.58	4.23	0.436	0.013	3.39	0.305	0.009	29.75	1.0174	0.340	2.08	1:6.1
9,869	20.11	25.43	4.44	0.474	0.005	3.58	0.238	0.003	30.33	1.0168	0.462	1.60	1:3.5
8,895	20.28	43.26	4.12	0.164	0.0	3.16	0.139	0.0	28.75	1.0161	0.325	1.73	1:5.3
8,895	19.75	38.40	4.26	0.180	0.002	3.24	0.230	0.003	28.38	1.0169	0.255	1.79	1:7.0
8,896	17.28	1.02	1.39	1.17	1.20	$\overline{1.42}$	0.850	0.867	20.65	1.0111	0.0	0.0	0.0
10,091	19.93	46.90	4.65	0.447	0.019	3.63	0.410	0.017	30.15	1.0177	0.308	2.17	1:7.0
10,074	20.11	50.00	4.75	0.380	0.014	3.51	0.350	0.005	30.60	1.0180	0.376	2.04	1:5.4
9,876	34.21	44.39	5.32	0.273	0.0	3.56	0.270	0.0	30.25	1.0177	0.343	2.13	1:6.2
10,069	33.50	37.89	4.62	0.400	0.009	3.54	0.357	0.010	30.98	1.0173	0.478	1.92	1:4.0
8,979	41.62	1.27	1.06	0.160	0.0	0.99	0.120	0.0	18.45	1.0052	0.056	0.45	1:8.0
10,081	34.21	42.78	4.20	0.450	0.024	3.35	0.380	0.006	28.75	1.0165	0.350	1.89	1:5.1
10,094	10.40	55.93	4.59	0.416	0.013	3.44	0.380	0.014	29.40	1.0172	0.310	2.15	1:7.0
9,868	19.57	28.82	4.73	0.365	0.007	3.63	0.261	0.01	30.42	1.0175	0.403	1.83	1:4.5
9,870	19.22	14.60	4.34	0.374	0.004	3.31	0.421	0.057	29.28	1.0179	0.386	1.50	1:3.9
10,084	31.92	45.30	4.42	0.420	0.038	3.45	0.410	0.031	29.85	1.0169	0.360	1.82	1:5.0

# GROUP III—Continued.

70.		Solids ning.	Cont	rents o	F CAN, D.			F	ILTEREI	Juice.			
Laboratory No.	Not Weight.	Per Cent. Solid by Draining.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Per Cent. Solids.	Per Cent. Ash.	Per Cent. Salt.	Refracto- meter Reading.	Specific Gravity.	Per Cent. Acid.	Per Cent. Sugars.	Acid-Sugar Ratio.
9,887	33.50	36.84	4.48	0.280	0.0	3.27	0.190	0.0	30.35	1.0172	0.380	1.90	1:5.0
9,883	33.86	49.11	4.91	0.370	0.005	3.66	0.397	0.003	30.65	1.0179	0.420	2.04	1:4.9
8,893	32.28	28.95	3.54	0.294	0.0	2.72	0.336	0.011	26.40	1.0039	0.319	1.51	1:4.7
8,891	32.80	25.27	4.58	0.440	0.008	3.66	0.337	0.0	30.75	1.0179	0.423	1.53	1:3.6
8,893	31.39	23.60	3.58	0.266	0.002	2.89	0.240	0.0	26.95	1.0144	0.283	1.71	1:6.0
8,892	31.74	2.22	2.98	0.320	0.0	2.80	0.336	0.0	26.95	1.0144	0.355	1.46	1:4.1
9,238	20.11	14.36	4.54	0.360	0.003	3.60	0.280	0.002	29.75	1.0170	0.348	2.36	1:6.8
9,862	19.75	40.18	4.70	0.438	0.047	3.48	0.297	0.006	30.65	1.0176	0.330	2.13	1:6.4

. Respectfully submitted,

CHARLES CASPARI, JR., State Food and Drug Commissioner.

BALTIMORE, March 10, 1916.

# Report of Cannery Inspector.

Dr. Chas. Caspari, Jr.,
State Food and Drug Commissioner.

DEAR SIR:

I herewith beg to submit my report for the year ending December 31, 1915.

Five hundred and four (504) canneries were inspected, a large number of them being visited several times. In the spring months a considerable number of canners were conferred with in regard to proper changes for the betterment of the sanitary conditions of their establishments, and I am glad to say with very gratifying results, 28 canneries in the counties now being model plants. In the year 1914 there was but one cannery in the counties which complied with the requirements of the Sanitary Inspection Law, whereas in the past year, 1915, inspection showed 102 to be so equipped. The tabulation which follows shows the percentage of increase in the several different requirements.

The 44 canneries in Baltimore City have all been provided with the equipment required by law, but improvements are still necessary in the insanitary way in which some of them are conducted and the uncleanly condition of their help.

It was thought desirable to secure samples of freshly canned tomatoes for the purpose of chemical investigation, and with this end in view the following information was secured in connection with each sample taken:

The name and location of cannery; date when sample was taken; condition of tomatoes before packing; were cores and skins removed before packing; time between gathering and packing of the tomatoes; does the sample secured represent the tomatoes above indicated; how were the cans filled; how was the juice added, if any; how long were the canned tomatoes processed.

Later in the season a sample can was secured from each of the Baltimore canners, with the following information: Name and location of cannery; date when sample was taken; date when the tomatoes were packed; how were the cans filled; was juice added or not; was juice boiled or not, and if boiled, how, by dry steam, open steam coils or jacketed kettle; how long were the tomatoes processed.

A large quantity of early Jersey tomatoes were brought to Baltimore, of which a great many arrived green, sour or soft, the juice in some instances running from the baskets and the tomatoes covered with white mold. As such fruit was unfit for canning purposes, the canners were commanded to have them picked over and were allowed to use only the sound, ripe fruit, the balance being hauled to the dump.

Six concrete bases with sanitary drainage and sewer connections were built this year, on which to rack up corn and pea ensilage to form a silo, whereby the stench usually coming from the seepage was avoided.

The sanitary toilet, as designed by the Bureau of Sanitary Engineering, does not meet the requirements of the larger canneries, and it seems impossible for them to keep the cans from overflowing, thus producing a most insanitary condition. Where concrete, flyproof pits have been built, these have proved most satisfactory.

The following information, arranged in tabular form, will give a better insight into the present conditions and the improvements which have been made during the past year:

Total number of fruit and vegetable canneries in the State	504
Number of canneries operating in Baltimore City	44
Number of cameries in the counties	460-
In operation during 1914 and 1915	350
Operating in 1914, but not in 1915	76
Operating in 1915, but not in 1914	11
Not operating in 1914 and 1915	23
New canneries built during the past year	15
Canneries destroyed by fire	6
Canneries dismantled	8

TABLE SHOWING THE INCREASE IN THE NUMBER OF COUNTY CANNERIES COMPLYING WITH THE REQUIREMENTS OF THE SANITARY INSPECTION LAW IN 1915, AS COMPARED WITH THE NUMBER COMPLYING IN 1914.

	1914.	1915.	Increase.
Floors	65.0%	91.0%	40.0%
Sanitary washstands	32.9%	71.2%	116.0%
Sanitary toilets	5.5%	51.1%	830.0%
Living quarters	93.7%	97.1%	3.6%
Disposal of tomato skins	37.8%	73.3%	93.7%

# As per following recapitulation:

# FLOORS.

Old concrete floor	118
New concrete floor	53
Old tongue-and-groove floor	53
New tongue-and-groove floor	50
Old part concrete and part tongue-and-groove	17
New part concrete and part tongue-and-groove	6
Built over water	10
Concrete and over water	2
Tongue-and-groove and over water	2
Concrete and brick	1
Tongue-and-groove and brick	. 1
Caulked, new	1
Caulked, old	1
Part tongue-and-groove and part caulked	1
Plain	18
Old double	6
New double	1
Part concrete and part plain	.4
Part tongue-and-groove and part plain	9
Part tongue-and-groove and part double	3
Plain and over water	2
Plain and tarred	1
Concrete and plain and tongue-and-groove	1
WASHING FACILITIES.	

Stationary washstands	102
New stationary washstands	155
Ordinary washstands	96
No provision	8

### TOILETS.

Old flush toilets	3
New flush toilets	13
Sanitary toilets with concrete	6
Old sanitary toilets	11
New sanitary toilets 1	
Toilets not fly-proof	
Separate toilets 1	
No toilets	

# LIVING QUARTERS.

Local help	164
	177
New private living quarters	10
Living quarters not private	10
DISPOSAL OF TOMATO SKINS.	
	8
Elevated box over concrete base and concrete platform for wagon.	
New elevated box over concrete base and concrete platform for wagon	16
Old concrete box and concrete platform for wagon	8
New concrete box and concrete platform for wagon	62
Wooden box over concrete base and concrete platform for wagon	25
Old concrete platform for wagon	17
New concrete platform for wagon	42
Elevated box over concrete base	2
New concrete skin box	15
Skins loaded in wagon, no platform	15
Skins hauled in tight wagons	5
Skins hauled in barrels	14
Skins pumped to the creek	4
Skins pumped to the field	4
Skins dumped into river	39
Skins dumped on pile near cannery	11
Skins carried by conveyor	2
Skins put into wooden box, resting on wooden platform, for wagon.	2
Skins put into wooden box	30
Skins put in iron box	2
Skins removed in zinc-lined wagon	3
Drain under wooden skin box	3
Skins carried by shute to swamp	1
DRAINAGE.	
DRAINAGE.	11
DRAINAGE.	11
DRAINAGE.  Drains to sewer	4
DRAINAGE.  Drains to sewer	8
DRAINAGE.  Drains to sewer	4 8 14
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.	$\begin{array}{c} 4 \\ 8 \\ 14 \\ .29 \end{array}$
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.  Tile drain to pit.	$\begin{array}{c} 4 \\ 8 \\ 14 \\ .29 \\ 9 \end{array}$
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.	$\begin{array}{c} 4 \\ 8 \\ 14 \\ .29 \\ 9 \\ 2 \end{array}$
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river  Cement drain to creek.	$\begin{array}{c} 4 \\ 8 \\ 14 \\ .29 \\ 9 \\ 2 \\ 9 \end{array}$
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.  Cement drain to river.  Cement drain to creek.  Cement drain to creek.	$ \begin{array}{r} 4\\ 8\\ 14\\ .29\\ 9\\ 2\\ 65 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to ditch  Cement drain to ditch  Cement drain to ditch  Cement drain to ditch	$ \begin{array}{r}  4 \\  8 \\  14 \\  .29 \\  9 \\  2 \\  65 \\  5 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek  Cement drain to field.  Drains to river.	$ \begin{array}{r}     4 \\     8 \\     14 \\     .29 \\     9 \\     2 \\     9 \\     65 \\     \hline     66 \\   \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.	$ \begin{array}{r} 4\\ 8\\ 14\\ .29\\ 9\\ 65\\ 66\\ 47 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.	$ \begin{array}{r} 4\\ 8\\ 14\\ .29\\ 9\\ 2\\ 65\\ 66\\ 47\\ 44 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to ditch.  Drains to ditch.  Drains to river.	$ \begin{array}{r} 44\\ 88\\ 14\\ 29\\ 9\\ 2\\ 9\\ 65\\ 5\\ 66\\ 47\\ 44\\ 13 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to ditch.  Drains to river.  Drains to ditch.  Drains to railroad ditch.  Drains to field.	$ \begin{array}{r} 44\\ 88\\ 14\\ 29\\ 9\\ 65\\ 56\\ 66\\ 47\\ 44\\ 13\\ 8 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek  Cement drain to ditch.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to rilroad ditch.  Drains to field.  Drains to pit.	$ \begin{array}{r} 48\\ 8\\ 14\\ 29\\ 9\\ 2\\ 9\\ 65\\ 5\\ 66\\ 47\\ 44\\ 13\\ 8\\ 6 \end{array} $
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to ditch.  Drains to field.  Drains to railroad ditch.  Drains to field.  Drains to branch.	$ \begin{array}{r} 48\\ 8\\ 14\\ .29\\ 9\\ 2\\ 9\\ 65\\ 5\\ 66\\ 47\\ 44\\ 13\\ 8\\ 6\\ 5 \end{array} $
DRAINAGE.  Drains to sewer  New drains to sewer  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river  Cement drain to creek  Cement drain to field.  Drains to river.  Drains to creek  Drains to river.  Drains to branch  Drains to branch  Drains by pump.	44 88 144.29 9 22 9 65 5 66 47 44 13 8 6 5 8
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to ditch.  Drains to field.  Drains to railroad ditch.  Drains to field.  Drains to branch.	$ \begin{array}{r} 48\\ 8\\ 14\\ .29\\ 9\\ 2\\ 9\\ 65\\ 5\\ 66\\ 47\\ 44\\ 13\\ 8\\ 6\\ 5 \end{array} $
Drains to sewer New drains to sewer Tile drain to river Tile drain to ditch Tile drain to pit. Cement drain to river Cement drain to creek Cement drain to field Drains to river Drains to creek Drains to river Drains to river Drains to river Drains to river Drains to brain to ditch Drains to pit. Drains to pit. Drains to pit. Drains to pit. Drains to branch Drains by pump No drainage.	44 88 144.29 9 22 9 65 5 66 47 44 13 8 6 5 8
DRAINAGE.  Drains to sewer  New drains to sewer  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river  Cement drain to creek  Cement drain to field.  Drains to river.  Drains to creek  Drains to river.  Drains to branch  Drains to branch  Drains by pump.	44 88 144.29 9 22 9 65 5 66 47 44 13 8 6 5 8
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to ditch.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to railroad ditch.  Drains to field.  Drains to field.  Drains to field.  Drains to branch  Drains to branch  Drains to branch  Drains by pump.  No drainage.  MISCELLANEOUS.	44 88 144.29 9 22 9 65 5 66 47 44 13 8 6 5 8
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to ditch.  Drains to railroad ditch.  Drains to pit.  Drains to branch.  Drains to branch.  Drains by pump.  No drainage.  MISCELLANEOUS.	48 814 .29 9 29 65 56 47 44 13 8 65 58
Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to river.  Drains to ditch.  Drains to railroad ditch.  Drains to field.  Drains to pit.  Drains to branch  Drains to branch  Drains to branch  Drains by pump.  No drainage.  MISCELLANEOUS.  New screened syrup-rooms.  Syrup-rooms not screened.	48 814 1429 9 22 9 9 65 566 47 44 13 8 6 5 8 23
DRAINAGE.  Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to ditch.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to ditch.  Drains to ditch.  Drains to ditch.  Drains to branch  Drains to pit.  Drains to pit.  Drains to branch.  Drains by pump.  No drainage.  MISCELLANEOUS.  New screened syrup-rooms.  Syrup-rooms not screened.  Husking sheds.	48 814 1429 9 22 9 9 65 566 47 44 13 8 6 5 8 23
Drains to sewer.  New drains to sewer.  Tile drain to river.  Tile drain to creek.  Tile drain to ditch.  Tile drain to pit.  Cement drain to river.  Cement drain to creek.  Cement drain to field.  Drains to river.  Drains to creek.  Drains to river.  Drains to ditch.  Drains to railroad ditch.  Drains to field.  Drains to pit.  Drains to branch  Drains to branch  Drains to branch  Drains by pump.  No drainage.  MISCELLANEOUS.  New screened syrup-rooms.  Syrup-rooms not screened.	48 88 144 .29 9 29 65 56 66 47 444 13 8 6 5 8 23

15

13

20

9

Concrete base for pea ensilage, sewer connection.....

Concrete platform to load corncobs on.....

Canners suspected of adding water to tomatoes.....

Canneries found in a very insanitary condition	Э
NOTICES MAILED TO CANNERS ORDERING THEM TO CORRECT CONDITIONS SO AS TO COMPLY WITH THE SANITARY INSPECTION LAW.	г
TO COTTECT HOOFS	27
To correct washing racingtessining to contract with the same and the s	12
10 Collect tolicus	88
TO COLLECT HAIR dual corp	2
10 correct disposar of change	14
10 correct dramage	<b>L</b> 7
To screen syrup-room	6

A detailed list of names showing the conditions found upon inspection of various canning establishments during the present year is withheld for the present, in order to give those parties, who have promised to do so, a chance to make further improvements before opening of the next season. Such a list, however, will be added to my report for 1916.

To provide caps and aprons for help.....

It would seem very desirable to have specifications for building sanitary toilets at the canneries prepared by our Bureau of Sanitary Engineering, in order that these may be supplied to those willing to conform to our requirements but ignorant of the best manner of fulfilling the latter. In a few isolated cases it may be necessary to institute prosecution against parties who do not appear willing to comply with our requirements, and this can undoubtedly be best done after the necessary specifications have been supplied.

Respectfully submitted,

Alfred K. Beasley, Cannery Inspector.

Baltimore, March 1, 1915.

# Report of Veterinarian and Dairy Inspector.

Dr. Chas. Caspari, Jr., State Food and Drug Commissioner.

DEAR SIR:-

I beg to submit my report for the year ending December 31, 1915.

Inspections of dairies and slaughter-houses during 1915 were limited to the following counties: Anne Arundel, Baltimore, Carroll, Frederick, Harford, Howard, Kent, Prince George, Queen Anne, Somerset, Talbot, Washington and Wicomico.

Number of inspections of dairies	523
Number of cows examined	660
Number of cows found diseased	42
Number of gallons of milk sold daily	016
Number of dairies found in sanitary condition	114
Number of dairies found in insanitary condition	125
Number of milk samples brought in for analysis	322
Number of inspections of slaughter-houses	105
Number of slaughter-houses found in sanitary condition	33
Number of slaughter-houses found in insanitary condition	22
Special inspections	15
Total number of inspections	643

The number of dairies and slaughter-houses inspected does not correspond to the number in a sanitary and insanitary condition, since some were inspected several times.

In making inspections of dairies I might explain what is required of the average dairyman. Good, healthy cattle and kept clean and udders washed before milking, and the mlk immediately cooled. To have the stables clean and whitewashed, well ventilated and well drained; also the barnyard well drained and the manure so disposed of that the cows cannot wade through it. To have a fly-proof dairy house and wash all utensils thoroughly as soon as possible after milking, and keep them in a fly-proof place, well lighted and ventilated. To have a sanitary toilet on the premises and to see that the water supply is not polluted.

There has been a marked improvement along these requirements in nearly every part of the State visited, even in Frederick county, where the dairymen had organized a dairymen's association and had combined to fight against these requirements, the majority have complied. By way of explanation, I might say that the President of this Association had his herd tested for tuberculosis with the intention of selling his milk in Washington, D. C., where they require all animals supplying milk to the District of Columbia to be tuberculin tested, and after three animals had reacted he had the test stopped and said he would sell his milk in Baltimore. I notified the Baltimore City Health Department, with the result that his milk was excluded from sale in this city. Since the guiding element of this Association was willing to sell milk from his herd to the public after knowing that three of his cattle had tuberculosis, with the possibility that more were affected, it looks bad for the Association and their object for associating. In fact, they said it was not the expense of making these tests or the changes necessary to safeguard the public health that they objected to, but the fact that they were commanded to do so.

There seems to be still one defect in the minds of a great many dairymen, and that is that they can strain the germs and filth out when they get into the pail of milk.

The great majority have also taken steps to prevent the pollution of their water supply. A great many have put concrete for quite a distance below and above the level of the ground, and also a tight cover over their wells to prevent surface drainage from getting in.

I must confess I feel greatly elated over the improvements the past year. Some farmers who were angry at the first and second inspection I have since had them to ask me to come to their place to make an inspection or see some new changes they had made or intended to make and have me suggest a way to improve their sanitary condition. If I can get equally as good results in the year 1916, I will feel that I have accomplished much for the benefit of the public health.

In reference to slaughter-houses, there has been a marked improvement in the sanitary conditions of all visited. They have put in tight floors and made their places fly-tight and provided adequate drainage and separate rooms for their hides, all whitewashed or painted. One of the greatest defects seems to be the

failure to remove offal daily, and it seems that constant inspection is the only remedy to overcome this evil.

Animals in an advanced stage of tuberculosis are being utilized for food. One man in Frederick, Md., killed an animal in an advanced stage of tuberculosis (two large pulmonary abscesses with plural adhesions). After I condemned it he insisted that he be allowed to grind it up for bologna, saying that he frequently used that kind for bologna.

The great difficulty is to visit these places at the time when slaughtering is going on. The only remedy I see is to have incorporated towns pass local inspection laws similar to what I suggested in last year's report.

Respectfully submitted,

W. B. BILLINGSLEY, Veterinarian.

March 1, 1916.

# Report of the Bureau of Sanitary Engineering, 1915.

ROBERT B. MORSE, Chief Engineer

# GENERAL STATEMENT.

The year 1915 represents the first full year of operation of the Bureau under the new water supply and sewerage law, Chapter 810 of the Acts of 1914. A brief outline of the powers granted the Department by this law was given in the report of With the increased power and appropriation provided the activities of the Bureau have been greatly extended and the number of employes increased to meet the demands of the work. The establishment of a resident engineer in the Southern District, with headquarters at Hyattsville, completed the chain of engineering representatives in different parts of the State, the Northern, Eastern and Western residencies having been filled in 1914. The organization of the entire force was so changed as to provide greater facility for handling the increased amount of work in both office and field. Office and field divisions were established, each under the direction of a division engineer, to whom all office and field employes, respectively, are responsible. Special work of any description is to be done by a special assistant engineer, who reports also upon the examinations of private water supplies. The present arrangement of organization is shown graphically on Figure I.

The amount of work done by this Bureau in 1915 as compared with 1914 is made clear by the data in Table I. The great increase in the various forms of activity is at once apparent. Particular attention is called to the marked change in the amount of general correspondence, resulting from the increased number of requests for the approval of plans and for general investigations, and in the number of investigations demanding written reports. The experience of 1915 has illustrated that the work demanding the attention of this Bureau is limited only by the working force allowed and that proper control and supervision over water supply and sewerage systems and conditions throughout the State are tasks which are far greater than the present means at the disposal of the Bureau make it capable of performing as completely as would be desired.




FIGURE I .- Organization, Office of Chief Engineer.

TABLE I.

COMPARISON OF WORK DONE BY ENGINEERING DIVISION DURING 1914 AND 1915.

ıs ved.	1915.	21	989	49	94	43	2	83	18	38	28	35	99	1	563
$\frac{Plans}{Received.}$														}	437
Permits Issued.	1915.	11	က	16	13	11	17	18	15	14	12	16	0	1	152
														1	7.7
$Field \\ Examinations.$	1915.	32	65	109	91	132	181	129	111	134	123	129	146		1,445
														l	509
Letters Received.	1915.	298	259	223	247	422	417	554	397	381	513	517	208		4,733
Let Rece	1914.	109	96	81	57	126	151	198	174	204	190	291	243		1,920
Reports, Record Data, etc.—Pages.	1915.	774	633	574	826	1.019	862	1.287	1,190	1,450	1.326	1,083	1,092		12,116
															6,309
Circular Letters Written.	1914. 1915.	0	0	<u>2</u>	59	200	0	0	0	0	0	0	0	1	331
Cir. Lea Wri	1914.	0	0	0	0	0	0	0	0	0	0	0	0	1	•
$\frac{1}{1}$	1914. 1915.	446	311	209	477	530	899	795	578	1.032	448	529	620		6,943
. Lea Wri	1914.	119	91	97	62	228	294	386	324	331	583	457	307		2,985
	Month.	January	February	March	April	May	Inne	July	August .	September	October	November	December		Totals

# Investigations in 1915

A detailed summary is shown in Table II of the total number of investigations of various kinds made in the different counties during the past year. The regular examination of water supply and sewerage systems was continued, while considerable additional work was carried on in the investigation of ice supplies. Some of the examinations included systems that had been previously investigated, while others were entirely new. Samples were collected at intervals from practically all the water supplies, and in this way an activity was extended which had not been given regular nor requisite attention previously on account of the small and none-too-mobile working force. By the end of the year practically all of the important water supplies in the State had been investigated and reported upon.

In addition to the more or less routine examinations, a number of special investigations were carried on, chief among which are:—a complete study, report and estimate of cost for an increased water supply for Brunswick; a preliminary plan and estimate of cost for a system of water supply for Myersville; a series of float investigations of the currents in Chesapeake Bay off Tolchester Beach as a part of a study for improving sewerage conditions at that resort; and the preparation of complete construction plans for improved water supply and sewerage systems at the Maryland House of Correction.

Sewerage improvements at the Springfield State Hospital were completed during the year and studies for improving the water supply were begun.

Sanitary surveys were made of Elkridge, Broome's Island, Emmitsburg and Mt. Airy, and the Bureau participated in the sanitary survey of Salisbury, which was carried on under the supervision of the Deputy State Health Officer.

TABLE II. FIELD INVESTIGATIONS MADE DURING 1915.

County.  Allegany	. Water Supply.	26 Semenage.	Bottled Water	0 0 0 I Ice Supply.	2 c Miscellaneous.	114 255 19
Baltimore City	7	1	11		0	
Calvert	3	3	0	0	18	24
Caroline	20	40	0	$\frac{3}{1}$	0	63
Carroll	31	20	0		8	60
Cecil	35	2	0	1	4	42
Charles	11	5	0	0	9	25
Dorchester	25	14	0	$\begin{array}{c} 0 \\ 2 \\ 0 \end{array}$	1	40
Frederick	55	12	0	<b>2</b>	3	72
Garrett	9	8	0		0	17
Harford	26	$\frac{2}{2}$	-0	0.	0	28
Howard	6		0	1	1	10
Kent	33	44	0	3 1	7	87
Montgomery	46	64	0	1	17	128
Prince George's	52	65	0	ī	6	124
Queen Anne's	11	6	0	$\frac{1}{2}$	7	26
St. Mary's	7	1	0	1	15	24
Somerset	13	24	0	9	1	47
Talbot	11	14	0	3 1	8	36
Washington	28	5	0	1	1	35
Wicomico	25	28	0	$\frac{1}{2}$	5	59
Worcester	22	21	0	2	2	47
			_			
Totals	710	536	21	33	145	1,445

Investigations of sanitary conditions were made also at the principal bay and river-shore summer resorts—Tolchester, Betterton, Chesapeake Beach, Bay Shore and Riverview. In the case of the first three, sanitary improvements are already in progress or under advisement.

The sanitary arrangements at schoolhouses, with particular reference to those buildings proposed for future construction, were given careful attention, and an attempt was made to have modern water supply and sewerage conveniences installed in all new buildings of appreciable size. This endeavor has resulted already in the installation of complete equipment in the new high schools at Betterton, Kennedyville and Rock Hall, and it is expected that several other schools on the Eastern Shore will make similar additions.

Brief summaries of the more important water supply and sewerage investigations will be found in a later portion of this report.

Considerable time was spent in carrying out investigations of the oyster beds in the vicinity of Cambridge and Annapolis in co-operation with the United States Public Health Service. The descriptions of the work and the conclusions reached have been discussed in detail in the official reports of the latter, and therefore only a short resumé will be repeated later in this report.

PLANS OF WATER SUPPLY AND SEWERAGE SYSTEMS RECEIVED

During the year 563 plans were received and filed. Water supply and sewerage plans were received from the following places:

# WATER SUPPLY SYSTEMS-

Existing—Braddock Heights, Cambridge, Elkton, Frederick, Frostburg, Hagerstown, New Windsor, Roland Park, Salisbury, Sparrows Point, Takoma Park.

Proposed—Annapolis (Annapolis Water Co., improvements), Boonsboro (Boonsboro Water Co., improvements), Chestertown (extensions and improvements), Decatur Heights, Elkton (Maryland Water Co., improvements), Greensboro, Laurel (extensions and filtration plant), New Windsor (extensions), Westernport (extensions and improvements), Westminster (Consolidated Public Utilities Co., filtration plant).

# SEWERAGE SYSTEMS-

Existing—Frostburg, Guilford, Havre de Grace, Loch Lynn, Roland Park Co., Sparrows Point (Maryland Steel Co.), Takoma Park (extensions).

Proposed—Arlington and vicinity (including disposal plant), Betterton (including disposal plant and storm-water drains), Cambridge (including disposal plant), College Park (Maryland Agricultural College, disposal plant), Decatur Heights (including disposal plant), Easton (disposal plant), Grantsville, Hurlock (including disposal plant), Kensington, Laurel (including disposal plant), Loch Lynn (extensions), Millington, Ocean City (including disposal plant), Takoma Park (extensions), Towson (Maryland State Normal School, disposal plant), Westernport (extensions).

# Plans Prepared in Connection With Investigations

Fifty-six complete plans and 183 sketches were prepared in this office in connection with various investigations. The principal plans completed are as follows:

## WATER SUPPLY SYSTEMS-

Bridewell-Maryland House of Correction-Existing water mains. Bridewell-Construction plans for water supply and sewage disposal improvements.

Brunswick-Existing water mains,

Emmitsburg—Water mains and private sewers.

Frostburg—Proposed layout of water mains in Mechanic St.

Myersville-Profile of proposed water supply and distribution system. Anacostia River Valley-Proposed water supply.

# SEWERAGE SYSTEMS-

Bridewell—See Water Supply.

Easton—Plan of future sewers for southern extension.

Emmitsburg—See Water Supply.
Rock Hall—Changes in sewer layout at High School.

Sykesville—Springfield State Hospital—Stream relocation near sludge bed.

### MISCELLA NEOUS-

Elkridge-Map for sanitary survey. Havre de Grace-Map for float investigation. Maryland—Map of State. Mt. Airy—Map of sanitary survey. Miles River and Vicinity-Map. Salisbury—Map for sanitary survey.
Tolchester—Results of float investigations. Details of sewers. Organization chart of Engineering Department.

# PERMITS ISSUED

Seventy-seven water-supply permits were issued to various individuals, companies and municipalities. The more important are as follows:

Construction of New Water Works Systems-Decatur Heights, East New Market.

Extension of Existing Distribution Systems — Boonsboro Water Co., Braddock Heights Water Co., Cambridge (Dorchester Water Co.), Chestertown, Frederick, Hagerstown (Washington County Water Co.), Kensington, Laurel, Mortimer Heights (Suburban Water Co.), New Windsor, Pocomoke City, Princess Anne, Roland Park Water Co., Salisbury Water Co., Takoma Park, Westernport.

Driving Wells—Cambridge (Dorchester Water Co.), Centerville, Ellicott City Water Co., Elkton (Maryland Water Co.), Mortimer Heights (Suburban Water Co.), Rockville, Sparrows Point (Maryland Steel Co.),

Sherwood Forest Co.

Construction of Filtration Plants—Baltimore County Water and Electric Company, Laurel, Westminster (Consolidated Public Utilities Co.).

Improvements to Existing Systems—Annapolis Water Co., Boonsboro Water Co., Braddock Heights Water Co., Brunswick, Chestertown, Elkton (Maryland Water Co.), Laurel, Princess Anne, Rockville, Westernport.

Installation of Disinfecting Apparatus—Annapolis Water Co., Arlington and vicinity (Suburban Water Co.), Piedmont Water Works Co.

Raising Money for Improvements—Chestertown.

Sixty-seven sewerage permits were issued. The more important are as follows:

Construction of New Sewerage Systems and Disposal Plants—Arlington and vicinity, Baltimore City Water Department (Montebello Filtration Plant), Betterton, Decatur Heights, Hillsdale (Kernan Hospital), Hurlock, Laurel, Ocean City, College Park (Maryland Agricultural College), Easton, Tolchester Beach (The Tolchester Co.), Towson (Maryland State Normal School).

Extension of Existing Sewerage Systems—Brooklyn, Cambridge, Chestertown, Curtis Bay, Frostburg, Germantown, Kensington, Loch Lynn, Pocomoke City, Preston, Ridgely, Riverdale (Riverdale Park Co.), Roland Park Co., Salisbury, Takoma Park, Tolchester Beach (The Tolchester Co.), Westernport.

Raising Money for Improvements—Easton, Kensington, Ocean City. Expending Money for Improvements—Easton, Kensington.

Three permits were issued for the sale of ice, as follows: Deal Island Ice Company, Easton Ice Company, Robert Maddox, Princess Anne.

Five permits were issued for the sale of bottled water, as follows:

Brooklandwood Spring Water Company, Big Rock Water Company of Baltimore, Chattolanee Spring Water Company, H. N. Robertson, Rockville (2).

# Orders Issued

Under the provisions of the new law giving the State Board of Health supervision and control over the maintenance, alteration, extension and construction of water and ice supplies, sewerage systems, trades wastes and refuse disposal in the State, orders were issued during the year to various municipalities, corporations and individuals, the more important of which are as follows:

Avalon—Baltimore County Water and Electric Company: Cessation of work upon the Avalon filtration plant pending receipt of a permit.

Westernport—Mayor and Commissioners: Repair and operation of disinfecting plant.

Mt. Rainier—Mayor and Common Council: Installation of water supply and sewerage systems.

Ocean City—Mayor and City Council: Installation of partial sewerage system.

Easton — Mayor and City Council: Sewage-disposal plant improvements.

Roland Park—Roland Park Company: Cessation of discharge of untreated sewage into Stony Run.

Kensington—Mayor and City Council: Construction of outfall sewer to Rock Creek.

Silver Spring—Commissioners of Montgomery County: Installation of water supply and sewerage systems.

Myersville—Burgess and Council: Submission of plans and specifications for a water supply system.

Delmar—Commissioners: Submission of plans and specifications for sewerage system.

Crisfield—Mayor and City Council: Submission of plans and specifications for a sewerage system.

# WATER SUPPLY INVESTIGATIONS.

Summaries of the more important reports on water supply investigations made during the year are given below:

### ANNE ARUNDEL COUNTY

Bridewell—Maryland House of Correction—The extremely bad condition of the water supply of this institution was described in the 1914 report. After authorization by the Governor, surveys were started late in 1914 and complete construction plans and estimates of cost of a project for providing a pure water supply and an improved distribution system for proper fire protection have been prepared.

Under the new plans Dorsey Run is to be utilized as the source of supply, the intake to be located preferably above the railroad track on a farm, the purchase of which is contemplated for farming purposes by the institution authorities. The water will be conveyed by gravity through a wood pipe line to an abandoned pool near the present pumping station, which is to be rebuilt to provide several days' water storage, so that at times of muddy water in the stream it may be drawn from. Provision is made for applying chemicals to coagulate the water and cause subsidence of a large part of the sediment when necessary. The existing pumping station will be used to raise the water from the pool to a slow-sand filtration plant of 250,000 gallons per 24 hours capacity, situated to the east of the prison, from which the clear water will be drawn by the pumps in the pumping station at the power-house and delivered to a new distribution system designed to provide greater pressure. Liquid chlorine will be used to disinfect the water after passing through the filters. An elevated steel tank will be placed on the hill near the old distribution reservoir.

# CAROLINE COUNTY

Federalsburg—Population in 1910, 1950. Federalsburg is located in the extreme southern part of Caroline county on Marshyhope Creek, a branch of the Nanticoke River. The water system, which is privately owned, was installed in 1900. Water

is obtained from two artesian wells, each 265 feet deep, and is collected in a small basin, from which it is pumped into the distribution mains, surplus pumpage going to an elevated tank of 10,000 gallons capacity. There are approximately 2 miles of mains and 175 services. The average consumption is about 33,000 gallons per day. The water is of good quality, but the system is thoroughly inadequate.

# CARROLL COUNTY

New Windsor—Population in 1910, 446. New Windsor is located in the western part of Carroll county on the Western Maryland Railway. The water system is owned by the municipality and was installed in 1908. Water is obtained from Big Sink Spring, located about two and one-half miles southeast of the town, from which it flows to a collecting reservoir, with a capacity of 400,000 gallons, and thence to the distribution mains by gravity. There are approximately 3.5 miles of mains, 160 services and 29 fire hydrants. No data as to water consumption were available, although the yield of the spring was estimated at about 95,000 gallons per day.

### FREDERICK COUNTY

Braddock Heights-Braddock Heights, an unincorporated community situated at the top of Catoctin Mountain about midway between Frederick and Middletown, is one of the popular mountain summer resorts. It has an all-year population of about 150 and a summer population of over 2000. The water system, which was installed about 1906, is owned by the Braddock Heights Water Company. Water is obtained from three springs. One of these is the famous Braddock Spring, located along the National Pike on the east slope of the mountain. The water flows to a collecting basin, from which it is pumped to a main distributing reservoir, which has a capacity of about 69,000 gallons, and to a smaller reservoir holding about 9000 gallons. From these reservoirs it flows by gravity to the distribution mains, of which there are approximately 2.5 miles. There are 68 services and 29 fire hydrants on the system. The water consumption is estimated at about 35,000 gallons per day at the height of the season. This water supply has been subjected to pollution as a result of insanitary conditions existing

on properties situated above the springs. The water company has been advised to acquire as many of these properties as possible for the purpose of protecting the water from contamination.

Brunswick—Population in 1910, 3721. The water supply of Brunswick was reported upon fully in the 1913 report. The city has large railroad yards and the population is rapidly increasing. The present source of supply is entirely inadequate and serves only about half of the residents. At the request of the city authorities an exhaustive investigation, extending over a year, has been made of all the possible water supply sources in the surrounding country. Mountain springs and available streams have been examined and measured.

Of the many possibilities that were presented, the investigation seemed to indicate as most favorable the development of Yourtee Spring, a large spring on the west side of South Mountain, yielding about 250,000 gallons per day in the dry seasons. The water from this source would be conveyed to the city in a wood pipe line passing around the south end of the mountain, then part of the way up the east side in order to intercept several springs on that side of the mountain, and finally would enter the town at the northwest corner and connect to the existing elevated tank. This development would serve for about five years and, it was estimated, would cost \$35,500, exclusive of land and rights-of-way. It was recommended that, after the capacity of these sources has been reached, a dam be built on the north branch of Little Catoctin Creek just above the Knoxville road, which would impound ultimately 270,000,000 gallons of water and supply 1,500,000 gallons per day. Since this water would require filtration, a plant was to be located near the existing elevated tank, to which the water would flow by gravity. Under such an arrangement both the high and low portions of the town would be served by gravity by the spring and stream waters, respectively. In the final system the distribution system would be divided into three zones, the highest one of which would be served by pumping.

A number of alternative schemes were presented. One of these provided for securing water from the Potomac River, to be used after filtration. Another suggested the immediate development of Little Catoctin Creek, with the idea of serving water to the railroad and using the revenue thereby derived to pay for a large part of the improvement.

The town is to apply to the next Legislature for the right to issue bonds sufficient to pay for the first improvement.

Myersville-Population, 475. Myersville has at present no system of public water supply. Investigations have shown that most of the private wells are unsafe because of the congested character of the town's development. A study for a proposed water system was made at the request of the County Health Officer, Dr. Ralph Browning. After investigating several springs on private property about two miles north of the town, in the vicinity of Pine Knob, a practicable method of supplying water seemed to be possible. It was recommended, therefore, that the springs on the properties of Marshall Warrenfeldt and I. S. Smith be developed first, to be followed if necessary by the development of springs on the property of John Frey. Water would be conveyed through wood and terra-cotta pipe to a reservoir of 225,000 gallons capacity, to be situated on high ground about three-quarters of a mile northwest of the town. The distribution system would consist mostly of 8-inch pipe in order to provide ample fire protection. The total cost of this system, exclusive of engineering, land and rights-of-way, was estimated at \$10,500.

Maryland Tuberculosis Sanatorium—This institution is situated in the Blue Ridge just north of Sabillasville. Including patients, doctors, nurses and attendants, about 600 persons live in the institution. On May 14 two cases of typhoid fever developed among laborers and eventually 13 cases appeared.

The water supply is derived from springs and driven wells. The main source of supply, that of Bowman Springs, is situated a little over a mile southwest of the buildings. The water from five small springs and one main spring is conveyed by a single pipe to a pumping station just east of the Sabillasville road. The water is pumped thence to a covered concrete reservoir of 132,000 gallons capacity near the buildings. In addition to the above supply, there are two springs, known as Naylor Springs, and five driven wells, varying in depth from 150 to 175 feet. All are located in the vicinity of the pumping station. Naylor Springs, the flow of which is small, discharge by gravity to the pump well, while water from the wells is obtained by air-lift. The flow from Bowman Springs supplies the de-

mand of the institution at all times except during the dry seasons of the year. The investigation of this water supply showed that each one of the sources, with the possible exception of the driven wells, was subject to pollution. At Bowman Springs a dwelling house, owned by the institution, is situated just above the sources, and the Naylor Springs appear to have insufficient protection.

As a precautionary measure a hypochlorite plant was installed to treat all the water supplied to the institution. Later a permanent liquid chlorine plant was put in.

#### MONTGOMERY COUNTY

Bradley Hills—Bradley Hills is a residential development located northwest of the District of Columbia. The water system is owned by the real estate company which is developing the property. Water is obtained from a well 263 feet deep and pumped directly into an underground pressure tank of about 24,000 gallons capacity, from which it is forced into the distribution mains. The total length of mains is small, and there are at present but six services and two fire hydrants on the system. No data are available as to the water consumption. The water is of good quality.

Friendship Heights—Friendship Heights is a small unincorporated suburb of about 200 population situated northwest of and adjacent to the District of Columbia. In 1910 a water system, privately owned, was installed. Water is obtained from three wells—two dug wells, 52 and 62 feet deep, and a driven well 115 feet deep. The water is pumped from the wells to an elevated tank of 7000 gallons capacity, from which it flows to the distribution mains. There are about 500 feet of mains and 16 services. The water consumption is estimated at 2500 gallons per day.

#### PRINCE GEORGE'S COUNTY

Hyattsville—Population in 1910, 1917. Hyattsville is located about one and one-half miles northeast of the District of Columbia between the Northeast and Northwest branches of the Anacostia River. The source of water supply comprises six driven wells, ranging in depth from 112 feet to 230 feet, situated near the easterly limit of the town. The water is delivered into a reservoir of 113,000 gallons capacity, from which it is pumped

into the distribution mains, the surplus pumpage passing to an elevated tank of 100,000 gallons capacity. There are on the system approximately 5 miles of mains, 460 services and 75 fire hydrants. The estimated average consumption is about 130,000 gallons per day. Although the water contains a large amount of iron, it is of safe hygienic quality.

#### WORCESTER COUNTY

Berlin—Population in 1910, 1317. Berlin is located in the northeastern part of Worcester County at the junction of the Delaware, Maryland and Virginia and the Baltimore, Chesapeake and Atlantic Railroads. It is served by a municipally owned public water supply system installed in 1909. Water is obtained from four driven wells about 100 feet in depth and is pumped directly into the distribution mains, surplus pumpage passing to a standpipe of 106,000 gallons capacity. There are approximately 3 miles of distribution mains, 205 services and 35 fire hydrants on the system. The average water consumption is about 55,000 gallons per day.

Pocomoke City—Population in 1910, 2369. Pocomoke City is located in the southwestern part of Worcester County on the south bank of the Pocomoke River. It is served by a municipally owned public water supply system installed in 1898. Water is derived from six wells about 130 feet in depth and is pumped directly into the distribution system, the surplus going to a standpipe of 86,000 gallons capacity. There are approximately 4.5 miles of distribution mains, 200 services and 43 fire hydrants on the system. The average consumption is 50,000 gallons per day. The water is of good quality.

Snow Hill—Population in 1910, 1844. Snow Hill, the county seat of Worcester County, is located on the south bank of the Pocomoke River, in the eastern part of the county. A municipally owned public water supply secures water from two driven wells 150 and 265 feet in depth. The water is pumped from the wells directly into the distribution system, the surplus passing to a standpipe of 38,000 gallons capacity. There are about 3.5 miles of distribution mains, 209 services and 40 fire hydrants on the system. The average consumption is about 55,000 gallons per day. The water is of good quality.

# SEWERAGE INVESTIGATIONS.

Summaries of reports on the more important sewerage investigations carried on during 1915 are given below.

#### ANNE ARUNDEL COUNTY

Bridewell-Maryland House of Correction-This Bureau was authorized by the Governor to prepare complete construction plans and estimates of cost for sewerage improvements at the institution along with those of water supply. A new sewage disposal plant of 200,000 gallons per 24 hours' capacity is contemplated under these plans, to be situated on the opposite side of the road from the existing disposal works, and whatever new sewers are necessary in the vicinity of the buildings. posal plant will consist of a screen chamber, a primary Imhoff tank, a sprinkling filter, a final Imhoff tank and sludge beds. The effluent will discharge into Dorsey Run. As this stream and the Little Patuxent River, into which it discharges, are not used as sources of water supply and flow through farming country, it has been deemed necessary to provide for only sufficient treatment of the sewage to prevent a nuisance in the stream, and a plant of this type will give such results.

#### BALTIMORE COUNTY

Arlington and Govans—On October 23, 1914, an order was served on the Commissioners of Baltimore County directing the construction of complete sewerage systems in the districts of Arlington and Govans, to be completed by January 1, 1916. This order was contested by the Commissioners and the case was tried in the Circuit Court of Baltimore County. A large amount of preparation was made for this case, the two areas in question were investigated thoroughly and much evidence collected to show the necessity of sewerage systems. The decision rendered was adverse to the State Board of Health, and an appeal was carried to the Court of Appeals. No decision had been given at the end of the year.

#### KENT COUNTY

Tolchester Beach—Tolchester Beach, a one-day summer resort owned by the Tolchester Company of Baltimore, is located on Chesapeake Bay 28 miles east of Baltimore City. Sewerage conditions at Tolchester attracted the attention of this Department early in the summer season, when it was noted that sewage discharging from the main sewer outlet was being carried by the tide directly over the bathing beach. As a temporary expedient, until experiments could be made to indicate a suitable location for an outlet, it was advised that the sewer be extended about 200 feet further, to discharge under the steamboat wharf. This was done.

During the latter part of the year float investigations were carried on to determine the effect of the discharge of sewage, from the existing sewerage system at the beach, upon the area used for bathing purposes. An effort was made to determine the direction of the currents in Chesapeake Bay under varying conditions of wind and tide and the influence of such currents upon particles floating on the surface of the water or suspended at various depths. A large number of floats were set out, at different stages of the tide and under varying wind conditions, from the present sewer outlet and from a point near the end of the steamboat landing, about 600 feet from shore, and their paths carefully followed. The majority of the floats passed directly over the bathing beach or grounded on it. It was apparent, therefore, that either of the above locations for a sewer outlet would menace seriously the purity of the water on the bathing beach. In the light of these experiments the Tolchester Company was prevailed upon to employ an engineer to prepare plans for rearranging the sewerage system and for a complete disposal plant. These plans had not been completed at the end of the year.

#### OYSTER-BED INVESTIGATIONS.

A brief discussion of the investigations made in co-operation with the United States Public Health Service is given below.

#### ANNE ARUNDEL COUNTY

Annapolia—Fresh raw sewage is discharged from many sewers along the shore of Annapolis in immediate proximity to a number of oyster beds. The dilution in the harbor and in the neighboring waters has been shown to be insufficient for the adequate disposal of the volume of sewage discharged from the town and from the United States Naval Academy. These conditions have resulted in the recommendation by this Bureau of a project for sewage disposal which would eliminate the danger of infectious pollution of oysters, but this project has not yet been favorably acted upon. Since any adequate system of sewage disposal should include the sewage from the Naval Academy, an effort was made to obtain the co-operation of the authorities of this institution, but little progress has been made so far.

The conditions existing in Spa Creek and the vicinity indicated great danger of infection from oysters kept there, and an order was promulgated, therefore, that the taking of shellfish for consumption from the waters between a line drawn from Greenbury Point to the mouth of Lake Ogleton to the upper railroad bridge was unlawful.

#### DORCHESTER COUNTY

Cambridge—Previous to the joint investigation of Cambridge Harbor by the Public Health Service and this Bureau, a detailed study of the pollution of this section of the Choptank River had been made by this Department. The result of both this and the later co-operative study indicate a polluted condition of the harbor which is a serious menace to the oyster bars in the Choptank River.

Inasmuch as sewerage conditions in this town are responsible directly for the intermittent recurrence of putrefaction in the

harbor, preliminary plans for the eradication of the objectionable conditions were prepared in 1914. A design for a complete sewerage system, including a disposal plant, designed along the lines laid down in the early report, was later prepared by a firm of consulting engineers. The construction of the disposal plant has never been started, but certain changes in the sewer lines have been made and the gradual construction of the entire system may be expected in the future. The elimination of the sources of pollution will accompany the development of improved sewerage arrangements.

# DISINFECTION OF PUBLIC WATER SUPPLIES IN THE STATE

(Extracted from Journal of the New England Water Works Association, Vol. XXX, No. 3. "Progress of Water Disinfection in Maryland," by Morse and Hall.)

There are 83 public water supplies in Maryland, serving about \$42,500 people—62.1 per cent. of the estimated total population of 1,357,374. Several community systems delivering water to small groups of houses, and also institutional supplies, are not included in this number.

In 1912, when the engineering investigations were begun, it was found that, in general, knowledge of the condition of the public water supplies in the State was meager. Analyses of the water from many of the supplies had been made occasionally, but no systematic method of examination had been followed. The Baltimore system was looked after by the city water and health departments, and consequently received little or no attention from the State Board of Health.

The State's investigations commenced to show that many of the water systems were badly in need of attention. The surface supplies were derived from streams fed by watersheds containing habitations, in many cases providing numerous sources of contamination, and, in general, protection was not afforded by long storage, filtration or other treatment. Water was usually taken from the stream at an intake dam and passed into a distributing reservoir of insufficient capacity to provide a safe period of storage. Moreover, water from some of the tubular well systems was found to be polluted, because the wells either were of too shallow depth and located in the midst of thickly settled districts, where sources of pollution were numerous, or were driven in rock containing fissures. Some supplies previously considered to be beyond suspicion were found, upon close investigation, to be in a bad condition and a menace to the public health.

In a large number of the instances where supplies were polluted the water was satisfactory from a physical standpoint, there being an absence of color, turbidity and other objection-

able characteristics. This was naturally true in the case of most well and spring waters. The water from streams fed by mountain watersheds and from watersheds containing much wooded or uncultivated land was also generally in good physical condition.

Where water, otherwise acceptable, was found to be in an unsafe sanitary condition, filtration did not appear to be necessary or even advisable for assuring the proper degree of purity. Expense and the improbability of proper operation are two factors which would make filtration plants undesirable for many of the small systems which abound in Maryland. Consequently, disinfection was resorted to wherever it would serve the purpose. There were, however, a considerable number of surface supplies derived from streams, flowing through cultivated country and furnishing a water occasionally or persistently turbid. Such supplies as these naturally require filtration. Disinfection in connection with filtration is general and is required in all new installations. While the Department has been able to obtain good results from several filter plants without the addition of disinfectants, the character of attention which small works ordinarily receive makes their use essential.

Prior to the establishment of the Engineering Division of the State Board of Health, in 1912, there were only four disinfecting plants in operation on water supplies in Maryland. Two of these were located on the Gunpowder and Jones Falls systems of Baltimore, and two, owned by the Baltimore County Water and Electric Company, were at the Avalon and Herring Run plants, which serve large sections in Baltimore County near the city. At Avalon disinfection was carried on in conjunction with filtration. All these installations used hypochlorite of calcium, and at Herring Run ozone was employed at times.

Since 1912 many towns in other parts of Maryland have been provided with purification plants, and it is a striking fact that, of the 842,565 persons served by public water supplies at the present time, 745,149, or 88.4 per cent. (approximately 54.9 per cent. of the population of the State), use water which is treated by either filtration or disinfection, or both.

The number of public systems in the State using raw, filtered, or disinfected ground and surface water, and the total populations of the communities served are shown by Table III.

There are 83 public water supplies in Maryland, 61 of which

derive their water from underground sources. The population thus served amounts to 102,099. The number of consumers is small in comparison with the number of systems, because these supplies, in general, are located in the smaller towns. The remaining 22 systems take water from surface streams and serve a population of 740,466. The large population using surface water is accounted for by the fact that Baltimore and most of the other places of considerable size are thus supplied. Ground waters are used by 12.1 per cent. of the people served by public systems, and by 7.5 per cent. of the total population of the State. The corresponding figures for surface waters are 87.9 and 54.6 per cent.

TABLE 111.

CLASSIFICATION OF PUBLIC WATER SUPPLIES IN MARYLAND.

TREATED.

UNTREATED.

El Zapplies.

	Per Cent. of Mondation Substitution Public Gupplies.	1.9	85.8 0.7 86.5 10.3 76.7	87.0 1.4 88.4	
	$\begin{vmatrix} Per \ Cent. \ of Total \\ Total \\ Population \\ Oo \ of State. \end{vmatrix}$	1.2	53.3 0.4 0.4 53.7 6.5 47.6	54.0 54.0	842,565 357,374
	Population of Communities (Served.	16,284 76,916 645,760	722,676 6,189 728,865 87,200 645,760	732,960 12,189 745,149	84 1,357
	to not $uv$ is sufficient.	9 6 +		18 21 21 21 21 21 21 21 21 21 21 21 21 21	
	Form of Treatment, Ground-Water Supplies, Disinfected only	Total treated	Total disinfected  Total filtered only  Total treated  All Supplies.  Disinfected only  Filtered and disinfected	Total disinfected Total filtered only	Total population served by public water supplies
	Per Cent. of Population Population Supplies.	1.4	11.6		served by post of State
7	Per Gent. of Tololl $G$ Tololl	6.9	7.2		population
	go Population of go Communities governed.	11,601	97,416		Total Total

1-

65

Of the ground-water supplies six are treated, while of the surface supplies only seven deliver untreated water, and these to only 11,601 people. The six plants for purifying ground water serve 16,284 persons, and communities with a total population of 728,865, supplied by fifteen systems, are using treated surface water. There are eighteen public supplies with disinfecting plants and three in which the water is filtered but not disinfected. A population of 732,960 is served with disinfected water, 12,189 with water filtered only, and 97,416 use raw water. Thus 54.1 per cent. of the total population of Maryland drink disinfected, 54.9 per cent. treated, and only 7.2 per cent. raw water. Of the total number of people served by public water systems, 87.0 per cent. use disinfected, 88.4 per cent. treated, and 11.6 per cent. raw water.

Not only is the influence of water purification felt by the public water supplies, but it has reached to State and private institutions as well. There are, as shown by Table IV, six institutions, embracing a population of 2352, where the water receives treatment of some kind, and in all but one of these it is disinfected. Included in this number are a penal institution, a tuberculosis sanatorium and four educational institutions.

TABLE IV.

CLASSIFICATION OF TREATED INSTITUTIONAL WATER SUPPLIES IN MARYLAND.

	MARYLAND.		
Class.	Form of Treatment.	No. of Supplies.	Population Served.
Ground water	Disinfected only	2 -	652
	Disinfected only	1	$1{,}100$ $350$ $250$
All supplies treated Disinfected supplies		 6 5	2,352 2,102

The location and nature of the public water systems and treated institutional supplies are shown in Fig. II. The universal use of ground-water sources, and the consequent scarcity of purification works, in the section of the State east of Chesapeake Bay is notable.

Disinfection by means of hypochlorite of calcium or liquid chlorine is naturally the most common method practiced, although one small mill town uses distilled water, which is the condensed exhaust from the steam turbines in the mill, and there is an ozone plant which is operated in conjunction with one of the hypochlorite installations. There is a possibility that ultra-violet ray disinfection will shortly be introduced, as its use is under consideration for a mechanical filter plant now being constructed.

The type of apparatus suitable for hypochlorite or liquid chlorine treatment in any particular case depends upon the method by which water is supplied. Where the flow of water to be treated is at most times substantially uniform, as in the case of certain pumping plants or in gravity lines discharging into reservoirs, a manually controlled apparatus is sufficient. Where fluctuations occur in the rate of flow, automatic control devices are essential for proper results. Liquid chlorine plants, so-called, introduce chlorine into the water either as a gas or in solution. The gas feed, usually called "direct feed," is well adapted for use where no water under pressure, for dissolving the gas, can be easily obtained. There are no direct-feed plants in Maryland at the present time, and only one, at Perryville, is regulated automatically, although automatic equipment is about to be furnished at Frederick.

The early disinfecting plants, as stated previously, all made use of hypochlorite of calcium, and with effective results. Those on the Avalon and Herring Run supplies of the Baltimore County Water and Electric Company were the first to be installed in the State. The former was constructed in 1908 and the latter during the year following. Late in 1910 the Jones Falls supply for Baltimore was first disinfected, and in June, 1911, the Gunpowder River plant was put into use. Until 1913 these were the only installations of disinfecting apparatus used for water purification in Maryland. A filtration plant had been previously built by the Washington County Water Company to supply Hagerstown from Antietam Creek during dry periods and hypochlorite dosing apparatus had been provided, but the disinfectant was not used until the summer of 1913. About the same time the State Board of Health constructed a plant for the Ellicott City system. Cumberland next followed, during the fall of the same year, by putting into operation a hypochlorite plant in conjunction with the mechanical filters on its new supply. In December the mountain supply of Hagerstown was furnished with hypochlorite apparatus. A typhoid fever epidemic at Rockville during February, 1914, made necessary the construction of an emergency hypochlorite plant by the State Board of Health for treating the supply derived from tubular wells driven in rock and proven to be the cause of the disease. The first liquid chlorine apparatus was put into operation at Union Bridge in January, 1914, for treating about 120,000 gallons of water derived from tubular wells driven in limestone rock. An ozone plant had been installed in 1909 at the Herring Run supply of the Baltimore County Water and Electric Company, but at first it was used, so the writers are informed, only experimentally.

After the powers of the State Board of Health were broadened, in 1914, and an increased appropriation was provided for sanitary engineering work, water purification plants were naturally introduced at a more rapid rate than previously. At the present time there are eleven liquid chlorine installations treating about 5,600,000 gallons of water daily and eleven hypochlorite plants treating approximately 90,600,000 gallons daily. Of the latter amount 80,000,000 gallons are delivered to Baltimore City.

Data concerning the disinfecting plants in Maryland are shown in Table V.

Changes are taking place continually in the water supply situation in this State, and it is expected that the number of disinfecting plants soon will be increased and that alterations will be made in some of the water supplies, which will involve plants now in operation. At Annapolis the existing liquid chlorine plant probably soon will be used in conjunction with a mechanical filtration plant, the construction of which is contemplated, and at the Maryland House of Correction the liquid chlorine apparatus already installed will supplement purification by slow-sand filters which were recently designed by the writers and which will soon be built. A filtration plant is being constructed at Westminster and one is being designed for the Springfield State Hospital. Liquid chlorine probably will be used for both of these installations. Liquid chlorine will soon replace hypochlorite at Westernport, and the same change is being considered for the Baltimore filtration plant, for which

TABLE V.

DATA CONCERNING WATER SUPPLY DISINFECTING PLANTS IN MARYLAND.

	Character of Attention.		Good. Indifferent. Not yet in continuous	Good.  Neglected—used only	Indifferent.	Good.	Good.	150,000 Water is condensed exhaust from steam tur- bines in mill of W. Va. Pulp and Paper Co.	Not yet in full opera-	Intermittent — much	Good Cood
	Where Chemical Is Applied.		Pump suction Pump suction Supply main	Pump suction Pump suction	Supply main	unuer pressure Before entering	Pump well	r is condensed ex s in mill of W. V	Pump suction	Pump suction	Pump suction
	Chlorine Parts per Million.		0.32	0.23	0.36	0.44	0.40	Water		09.0	0.89
	Average Amount Treated Daily (Gallons).	on Only.	$\begin{array}{c c} 1,224,000 & 0.32 \\ *72,000 & 0.30 \\ 15,000 & \dots \end{array}$	170,000 0.23 70,000	1,000,000 0.36	2,500,000 0.44	2,800,000 0.40	150,000	30,000	200,000 0.60	90,000 0.89
	Date.	infecti	1915 1914 1915	1915	1915	1913	1909	1912	1916	1915	1915
	Character of Supply.	Plants for Disinfection Only.	Surface Ground Surface	Ground	Surface	Surface	Surface	Surface	Surface	Surface	Ground
0	Type of Plant.	Pla	Liquid chlorine Liquid chlorine Liquid chlorine	Hypochlorite Hypochlorite	Liquid chlorine	Hypochloriter	Hypochlorite	Ozone Distillation	Hypochlorite	Liquid chlorine	Hypochlorite† Liquid chlorine
	Population (Estimated 1916).		8,656 5,250 372	2,730	11,056	25,500	25,493	950	300	800	009
	. Place.		Annapolis Arlington Bloomington	Chestertown	Frederick	Hagerstown	Herring Run	Luke	Maryland Agricultural	Maryland House of Correction	Maryland Tuberculosis Sanatorium

Neglected—used only occasionally. Indifferent. Good. Good. Good. Neglected at first; at present good.		Good. Good. Good.	Not yet in continuous operation.	Good.
9,000 0.83 Storage tank 300,000 0.50 Supply main 30,000 0.22 Collect. res. 4,000 1.50 Storage tank 120,000 0.35 Pump suction 450,000 0.40 Intake at dam		Surface         1908         1,500,000         0.80         Outlet of coagu-food         Good           Surface         1915         80,000,000         0.30‡         Filtered water         Good           Surface         1913         5,500,000         0.20         Inlet to coagu-food         Good	Filtered water	30,000 0.25   Pump suction
9,000 0.83   0,000 0.50 0.22 0.000 1.50 0.000 0.35 0.000 0.40	ion.	0.80 0.30‡ 0.20	:	0.52
9,000 0.83 300,000 0.20 30,000 0.22 4,000 1.50 120,000 0.35 450,000 0.40	: Disinfect	1,500,000 80,000,000 5,500,000	70,000	30,000
1915 1915 1916 1916 1914 1914	n and	1908 1915 1913	1915	1915
Ground   1915   Surface   1915   Ground   1915   Ground   1915   Ground   1914   Surface   1914   Surface	Plants for Filtration and Disinfection	Surface Surface Surface	Surface	Ground
Hypochlorite Liquid chlorine Liquid chlorine Hypochlorite Liquid chlorine Hypochlorite	Plants f	Hypochlorite Hypochlorite Hypochlorite	Hypochlorite	350 Liquid chlorine   Ground   1915
200 632 1,221 52 883 3,109		30,197 587,112 25,843	2,608	350
Perryville		Avalon 30,197 (Baltimore County). Baltimore City587,112 Cumberland	Laurel	Notre Dame

\*This amount is mixed with raw water from other sources.

‡As high as 3.0 parts per million of available chlorine have been used in treating the raw water before the filtration †Now kept for emergency use. plant was installed. hypochlorite was recently quoted at \$290 a ton, on the basis of a yearly contract. There is a marked tendency to use liquid chlorine instead of hypochlorite of calcium in connection with old as well as new installations, and the high price which now has to be paid for the latter will stimulate this movement.

The amount of chemical used at the disinfecting plants in Maryland has varied widely according to local conditions. It has ranged from 0.2 to 3.0 parts per million of available chlorine in the case of hypochlorite treatment, and from 0.25 to 0.67 parts where liquid chlorine is used. Where the disinfectant is applied to a spring or well water, or to the effluent from a well-operated filtration plant, the amount required is usually near the minimum mentioned above. In the case of turbid waters, larger amounts have been found necessary for proper treatment, and at times of extreme turbidity excessive quantities are required. In the treatment of the Baltimore City water as high as 3 parts per million of available chlorine, or 75 pounds of hypochlorite to the million gallons, have been used on raw water. The amount necessary in many instances depends also upon the character of attention given the apparatus.

It is by no means a fact that the disinfecting plants in Maryland have been operated with complete satisfaction, for most of them have had trouble of some kind. The difficulties encountered could generally have been avoided with more attention to proper operation. With hypochlorite plants the principal annoyance has been due to the formation of coatings which clog the pipes; and, where the solution was particularly strong, the materials of which the piping was made have been attacked, necessitating the replacement of parts. These are not serious troubles, however, and can be eliminated by frequent inspection. In the case of liquid chlorine plants, it has been found that occasionally a part becomes corroded and requires replacement, or a valve gets out of adjustment, but the gradual perfection of apparatus is making such occurrences less frequent.

Where faithful and intelligent attention has been given to operation, and the amount of disinfectant used has not been excessive, few complaints have been made regarding tastes or odors in the water. In one instance, where a liquid chlorine plant of the manual control type was disinfecting water the flow of which fluctuated considerably, there was always an overdose at night, which caused complaint due to tastes noticeable during the early hours of the day. This, of course, was the

fault of the method employed and not of the treatment itself. In general, fewer complaints have been registered concerning disagreeable tastes where liquid chlorine was used than in the case of hypochlorite.

There has been a marked decrease in the prevalence of typhoid fever in some of the communities where disinfecting plants have been installed. In most cases, however, the period of operation has been so short, and, on account of the small size of some of the places, the appearance of typhoid so irregular, that records do not indicate clearly the effect produced. Moreover, much of the typhoid fever in Maryland is attributable to causes other than impure water supplies. Nevertheless, there are certain well-defined instances where a decrease in typhoid fever morbidity and mortality can be traced directly to purification of a water supply. In Baltimore the reduction has been very pronounced. After the disinfecting plants were built the prevalence of typhoid immediately fell off, and the im-In 1915 both the morbidity provement has since continued. and mortality rates from this disease, although higher than in many of the larger cities of the country, were approximately half as great as in 1910. Following the introduction into Cumberland of filtered and disinfected water from Evitts Creek, in place of the former raw supply from the Potomac River, the occurrence of typhoid fever was so reduced that, in 1915, there were scarcely more than one-sixth as many cases of, and onefifth as many deaths from, this disease as in 1913, in spite of the fact that the corporate limits had been enlarged in the meantime. However, the change in source and the long storage period provided in the reservoir would undoubtedly have caused a large proportion of this improvement without the presence of a purification plant. One of the most striking instances of the effect upon typhoid occurrence, caused by disinfecting a water supply in Maryland, was that of the Rockville epidemic, which, after 28 cases had occurred, was checked suddenly by the installation of an emergency hypochlorite plant.

In order that prompt remedial measures may be instituted when a water supply is at any time found to be in an unsafe condition, or when an outbreak of typhoid fever casts suspicion on a water previously untreated, the engineers of the State Board of Health are all instructed in the setting up of an emergency hypochlorite dosing apparatus, making use of ordinary and easily obtainable materials. A sketch showing the general arrangement of the device used, including a bill of material, has been prepared, and is usually sent ahead, so that the necessary material may be on hand when the engineer arrives at his destination. This apparatus, shown on Fig. III, has proven its value on many occasions, and several of the installations are still in operation.

# Clerical Report and Financial Statement.

# WALTER M. KIRKMAN, Chief Clerk

# CLERICAL REPORT.

Number of letters written	24,471
Number of letters received	16,775
Number of multigraphed sheets (196 electros, 485 forms)	819,229
Number of pages folded on folding machine	73,000
Number of pages typewritten for reports, etc	16,346
Number of birth certificates filed	20,458
Number of death certificates filed	13,532
Number of second-class packages sent out	5,530
Number of laboratory and miscellaneous reports	12,366
Number of reports sent to physicians	6,023
Number of disinfection sheets mailed	435
Number of certificates of qualification issued	109
Number of certificates of registration issued	75
Number of certified copies of death issued	500
Number of certified copies of births issued	32
Number of searches made for copies of births and deaths	877
Number of circulars and circular letters sent out	10,446
Number of receipts for duplicate portion of transit permits	1,349
Number of cards written for cross index	79,940
Number of registration cards written	10,399
Number of postal cards to obtain given name of children written	3,383
Number of cards sent out	11,356
Number of Anne Arundel Co. Survey cards punched and num-	
bered	12,953
Number of given names entered on certificates of birth	652
Number of cards resent, including miscellaneous	154
Number of cards unclaimed and sent to local registrars	181
Number of vouchers sent out	1,111

# DISTRIBUTION OF TUBERCULOSIS PROPHYLACTIC SUPPLIES.

$\it Materials.$	To Substations.	Individuals.	To Institutions, Baltimore City Health Dept., etc.	Total.
Sputum Cup Fillers	125,300	26,337	447,790	599,427
Sputum Cup Holders	1,347	223	2,408	3,978
Napkins	125,300	23,300	1,211,000	1,359,600
Disinfectant (½ pt. bottles)	1,347	346	12,813	14,506
Disinfectant (gallons)			50	50
Waterproof Pockets	1,566	200	500	2,266
Books of Information	3,783	400		4,183
Pocket Flasks	11.750	2.000	1.300	15,050

With respect to the clerical report, every item shows a very decided increase in the amount of work performed, thus illustrating the growing activities of the Department.

The following is a statement of the receipts and expenditures of the Department for the year ending December 31, 1915. The receipts are classified according to the names of the appropriations, and the disbursements by bureaus according to the objects of expenditure.

The balance on account of appropriations as of January 1, 1915, of \$75,236.12, does not represent unexpended balances, since the fiscal year of the Department does not coincide with the calendar year, because the due dates of the several appropriations fall between April 1 to March 31 as the fiscal year for the Department. To the balance as of January 1 (\$75,236.12) has been added the amounts due on the several appropriations—namely, \$142,500—and also the amounts received from sundry sources, which latter items total \$1,482.23, giving total receipts for the year \$219,218.35.

The disbursements follow, being arranged according to bureaus and objects of expenditure. The footings of the columns at the bottom of the sheet show the amounts spent by the various bureaus, and the total column on the right shows the total amount spent for each item by all the bureaus. The total amount of expenditures was \$140,188.19.

The next division represents the amounts which reverted to the State Treasury on account of unexpended balances of appropriations. It will be noted that on account of Sanitary Districts appropriation an amount of \$18,102.33 was returned to the State Treasury. This balance existed because the field organization of the Deputy State Health Officers was not complete, there being six Deputy State Health Officers in the field and one unassigned, instead of ten as required by law. Therefore, the money which would have been spent in equipping and maintaining these offices was not spent, and the balance, therefore, reverted to the State Treasury. This was an unusual situation and would not have existed had the field organization of the Deputy State Health Officers been complete. It will also be noted that \$1280 reverted to the State Treasury on account of Pasteur Treatment Appropriation. This represented an unexpended balance in an appropriation which was provided solely for the administration of Pasteur treatment to indigent

persons. Deducting the total amount of reversions to the State Treasury, \$19,495.90, from the amount obtained by deducting the total expenditures from the total receipts—namely, \$79,030.16—leaves a balance of \$59,534.26 as of December 31, 1915, to carry to due dates of the several appropriations during the year 1916.

For the first three months of the year the accounts were audited monthly by the Executive Committee of the State Board of Health. Commencing April 1, the Board employed the Baltimore Audit Company to audit the accounts of the Department quarterly. Reports of the auditor are hereto attached.

Baltimore, Md., October 1, 1915.

The State Board of Health, 16 West Saratoga Street, Baltimore, Md.

## Gentlemen:

In accordance with your request we have made an audit of the books and accounts of your Board for quarter ending June 30, 1915, and found everything in good order for the period under review.

We verified additions, checked postings and found all disbursements supported by proper vouchers.

Very truly yours,

BALTIMORE AUDIT COMPANY,

(Signed) Ernest E. Wooden, Certified Public Accountant, Secretary and Treasurer.

Baltimore, Md., November 3, 1915.

State Board of Health, 16 West Saratoga Street, Baltimore, Md.

#### Gentlemen:

We have completed the audit of the books and accounts of your Board for quarter ending September 30, 1915, and found everything in good order for the period under review. We verified addition, checked postings, examined cancelled checks, and found all disbursements supported by proper vouchers.

Very truly yours,

## BALTIMORE AUDIT COMPANY,

(Signed)

ERNEST E. WOODEN, Certified Public Accountant, Secretary and Treasurer.

Baltimore, Md., March 7, 1916.

State Board of Health, Baltimore, Md.

## Gentlemen:

We have completed the audit of the books and accounts of your Board for quarter ending December 31, 1915, and found them in good order for the period under review. We verified additions, checked postings, examined cancelled checks and found all disbursements supported by proper vouchers. We also proved the cash expense funds and found same to be correct.

Very truly yours,

# BALTIMORE AUDIT COMPANY,

(Signed)

ERNEST E. WOODEN, Certified Public Accountant, Secretary and Treasures


# INDEX OF TABLES AND CHARTS

			PAGE
Table	I.	Estimated Population in Maryland for the	6
m - 1-1 -	T 4	Year 1915 Estimated Population for the Year 1915	7
Table		Birth Rates, Maryland, 1905-1915	8
Table Table		Birth Rates, Death Rates and Rate of in-	J
		crease, 1915	9
Table	III-A.	Births, Deaths and Increase of Population,	10
/II-1-1-	TXY	1915	11
Table Table		Births, Male and Female, by Months—	11
		Maryland, 1915	15
Table		Stillbirths—Maryand, 1915	
Table		Illegitimate Births—Maryland, 1915	19
Table	VIII.	Percentage of Native and Foreign Born Parents of Children Born in Maryland	
		During 1915	20
Table	IX.	Table of Fecundity for White and Colored Mothers by Age Periods—Rural Mary-	
		land, 1915	22
Table	IX-A.	Table of Fecundity for White and Colored Mothers by Age Periods—Baltimore City,	00
m - 1-1 -	37	Table of Marinary Franchista Nambar 6	23
Table	Λ,	Table of Maximum Fecundity—Number of	
		Children in Excess of Tenth Born in Maryland, Exclusive of Baltimore City,	97
Table	VI	1915	27
rabte	Χ1,	and Mothers of Children Born in Mary-	00.00
Table	VII	land, 1915	8-30
Table	711'	Ages of Parents of Children Born in Mary-	91
Table	VIII	land During 1915	31
Table	X111.	Average Age of Parents of Children Born in Rural Maryland, 1906-1915	32
Table	XIV.	Proportion of Births Attended by Physi-	
		cians and Midwives, White and Colored,	
_		Maryland, 1915	33
Table	XV.	Percentage of Births_Attended by Physi-	
		cians and Midwives, White and Colored,	
	****	Maryland, 1915	34
Table	XVI.	Births by Sex and Color, Estimated Popu-	
		lation and Birth Rates per 1000 of Cities	
		and Towns in Maryland with Population	
mahl.	XXXXXX	of 2000 and Over, 1915	35
Table	XVII.	Deaths by Sex and Color, Estimated Popu-	
		lation and Death Rates per 1000 of Cities	
		and Towns in Maryland with Population	0-
Table	VIIII	of 2000 and Over, 1915	37
ranie	XVIII.	Deaths for the Year 1915 by Ages, Showing	
		Percentages of Total Mortality in the	
		Several Age Periods of Life in Rural	
		Maryland, Baltimore City and the State	20
Table	XIX-A.	of Maryland Population and Deaths per Thousand at	38
Table	**141_U'	the Several Age Periods, Maryland, 1915	42
			Tá
		235	1944

	`		PAGI
Table	XIX-B.	Estimated White Population and Deaths	
		per Thousand at the Age Periods for the	
		Year 1915—Maryland's Estimated White	43
Tabla	XIX-C.	Population, 1,120,769 Estimated Colored Population and Deaths	
Table	A1A-0,	per Thousand at the Age Periods for the	
	× .	Year 1915—Maryland's Estimated Col-	
		ored Population, 231,171	44
Table	XX.	A Classification of Causes of Death, With	
20010		the Number Occurring and their Ratios	
		to the Mortality (Maryland, 1915)	47-48
Table	XXI.	Twenty Principal Causes of Death, Mary-	
		land, 1915	51
Table	XXII.	Pulmonary Tuberculosis — 1915 — Giving	
		Deaths by Ages, Deaths per 10,000 of	
		Population at Age Periods, Per Cent. in	٠.
	******	Each Age Period, White and Colored	54
Table	XXIII.	Deaths per 10,000 of Population from Pul-	
		monary Tuberculosis—White and Colored	57
Table	XXIV.	—Maryland—1915	57
Table	AAIV.	Maryland, 1915, Showing Number and	
		Percentage of Deaths in Each of Three	
		Age Periods	60-62
Table	XXV.	Age Distribution of Mortality from Twelve	00-02
111010		Diseases by Percentages, Rural Mary-	
		land, 1915	• 64
Table	XXVI.	Deaths from Five Principal Respiratory	
		Diseases in Rural Maryland, 1907-1915,	
		Inclusive	71
Table	XXVII.	Pulmonary and Laryngeal Tuberculosis.	
		1908-1915	72
Table	XXVIII.	Deaths by Months for 1915, 1914, 1913,	
		1912, 1911 and 1910	76
		Eight Prominent Causes of Death by	78
Tabla	XXIX.	Months—Rural Maryland, 1915	~18
Lable	AAIA.	Parasitic Diseases, Showing the Organ Affected	80
Table	XXX.	Special Causes of Death	
	XXXI.	Infant Mortality Rate, White and Colored,	00-01
		Maryland	89
Table	XXXII-A.	Presumptive Expectancy and Duration of	
		Life—Maryland, 1915	94-95
Table	XXXII-B.	Presumptive Expectancy and Duration of	
		Life—White, 1915	96-97
Table	XXXII-C.	Presumptive Expectancy and Duration of	
rn 1 1 -	X7X7X7XX	Life—Colored, 1915	
	XXXIII.	Mean Age at Death by Counties, 1915	104
	XXXIV. XXXV-A.	Mean Age at Death by Months, 1915	105
Table	AAAV-A,	Survivorship in Maryland, 1915, Total	105
Tabla	XXXV-B.	Population	107
Labic	TEXEST A.D.	Maryland. 1915, White and Colored	107
Table	XXXVI-A.	Estimated Population of Maryland	111
	XXXVI-B.	Deaths by Ages	111
Table	XXXVI-C.	Death Rate per Thousand at Age Periods	112
	XXXVII.	Survivorship by Ages—1910, 1911, 1912	
		1913, 1914, 1915	112
	XXXVIII.	Marriages for Maryland, 1915	114
Table	XXXXX	Divorces for Maryland 1915	115

INDEX. 237

			PAGE
Chart	1.	Births by Months, Male, Female and Total Mary-	
02		land, 1915	14
Chart	2.	Chart Showing Fecundity for White and Colored	25
		Mothers by Age Periods, 1915	39
Chart	3.	State of Maryland—Deaths by Ages, 1915 Deaths by Ages—Rural Maryland and Baltimore	90
Chart	4.	City. 1915	40
Chart	5.	Distribution of the Population and Deaths by Ages,	
Спагс	Ð.	Maryland, 1915	45
Chart	6.	Maryland Classification of Causes of Death, 1915	49
Chart	7.	Twenty Principal Causes of Death, 1915	52
Chart	8.	Tuberculosis, 1915 — Comparative Chart Giving	
		Deaths per 10,000 at Various Age Periods, Mary-	
		land. White and Colored	55
Chart	9.	Tuberculosis, 1915—Comparative Chart Demonstrat-	
		ing the Percentage of Deaths at Various Age	
		Periods from Tuberculosis, White and Colored,	=0
Q1 i	10	Maryland	56
Chart	10.	Principal Causes of Death, Classified According to Their Importance in the Middle Period of Life	59
Chart	11	Age Distribution of General Mortality, 1915	65
Chart		Age Distribution of Mortality from Malignant Neo-	09
Chart	1	plasms, Diphtheria, Cerebral Hemorrhage	66
Chart	13.	Age Distribution of Mortality from Typhoid Fever,	
-		Whooping Cough, Tuberculosis	67
Chart	14.	Age Distribution of Mortality from Paralysis,	
		Bright's Disease, Accidental Violence	68
Chart	15.	Age Distribution of Mortality from Pneumonia,	
O7:	10	Broncho-Pneumonia, Organic Heart Disease	69
Chart	16.	Mortality from Five Respiratory Diseases, 1907-	=0
Chart	17	1915	70
Спать	11.	Rural Maryland	74
Chart	18.	Eight Principal Causes of Death, by Months—Rural	12
		Maryland, 1915	77
Chart	19.	Comparative Susceptibility of the Several Organs	
		to Parasitic Infections—Maryland Mortality, 1915	81
Chart	20-A.	Presumptive Expectation and Duration of Life-	
Ob - wt	00 <b>D</b>	Total Maryland, 1915	100
Chart	20-В.	Presumptive Expectation and Duration of Life-	
Chart	20 C	Total Maryland, 1915	101
Спаті	<b>20-0.</b>	Presumptive Expectation and Duration of Life— Total Maryland, 1915	100
Chart	21.	Maryland Biometer. 1915—Number of Survivors of	102
Chart		10.000 Persons Born in Maryland in 1915 at Suc-	
		ceeding Quinquennial Periods, Assuming the	
		Death Rate of that Year to be Constant	109
Chart	22.	Maryland Biometer, 1915-Number of Survivors of	
		10,000 Persons, White and Colored, Born in Mary-	
		land in 1915 at Succeeding Quinquennial Periods,	
		Assuming the Death Rate of that Year to be Con-	
		stant	110



# ALPHABETICAL INDEX

Accidental Violence—PAGE
Age Distribution of Deaths from, by Percentages, Chart 14 68
Deaths from 1906-1915, Table XXX
Age Periods—
And Deaths Per Thousand at the Table XXXVI-C
Population of Maryland by Table XXXVI-A 111
Population and Deaths Per Thousand at the Several, Table
XIX-A, B. C, Chart 542-45
Table of Fecundity for White and Colored Mothers, Tables IX
and IX-A, Chart 222-26
Age at Death—
Mean, by Counties, Table XXXIII
Mean, by Months, Table XXXIV
Age Distribution—
Of Deaths from Accidental Violence by Percentages, Chart 14 68
Of Deaths from Bright's Disease by Percentages, Chart 14 68
Of Deaths from Broncho-Pneumonia by Percentages, Chart 15. 69
Of Deaths from Cerebral Hemorrhage by Percentages, Chart 12. 66
Of Deaths from Diphtheria by Percentages, Chart 12 66
Of Deaths from Malignant Neoplasms by Percentages, Chart 12. 66
Of Deaths from Organic Heart Disease, Chart 15
Of Deaths from Paralysis by Percentages, Chart 14 68
Of Deaths from Pneumonia by Percentages, Chart 15 69
Of Deaths from Tuberculosis by Percentages, Chart 13 67
Of Deaths from Typhoid Fever by Percentages, Chart 13 67
Of Deaths from Whooping Cough by Percentages, Chart 13 67
Of Mortality from Twelve Diseases by Percentages, Rural Dis-
tricts, Table XXV, Chart 1164-65
Age of Parents of Children Born in Rural Maryland During 1914, Table XII
Ages—
Average of Parents
Deaths by, Table XXXVI-B
Deaths by, from Pulmonary Tuberculosis, Deaths Per Ten Thou-
sand of Population at Age Periods, Per Cent. of Deaths in
Each Age Period, White and Colored, Table XXII, Charts
8 and 9
Deaths by, in Maryland Rural Districts and Baltimore City.
Table XVII. Charts 3 and 4
Maximum and Minimum of Fathers and Mothers in Counties of
Maryland, Table XI
Of Parents, Mean, of Children Born in Rural Maryland, Table
XIII
Survivorship by, 1910-1915, Table XXXVII
Alcoholism, Deaths from, 1906-1915, Table XXX
Allegany County, Area, Population and Reported Cases of Notifiable
Diseases
Annapolis, Oyster Bed, Investigation of
Anne Arundel County, Area, Population and Reported Cases of No-
tifiable Diseases
Anti-Typhoid Vaccine, Distribution of
Area, Population and Reported Cases of Notifiable Diseases—
Allegany County
Anne Arundel County
Baltimore County
Calvert County

	PAGE
Caroline County	138-139
· Carroll County	
Cecil County	
Charles County	
Dorchester County	
Frederick County	144-145
Garrett County	145-146
Harford County	
Howard County.	
Kent County	148
Montgomery County	149-150
Prince George's County	150-151
Queen Anne's County	151-152
St. Mary's County	152-153
Somerset County	159 154
To bot Country	100-104
Talbot County	154-155
Washington County	
Wicomico County	157
Worcester County	158-159
Arlington, Sewerage, Investigation of	215
Arteries, Deaths from Infections of, Table XXI, Chart 7	51.59
Average Age at Death, Method Used for Obtaining	01.00
A verage Age of Pearly Method Osed for Obtaining	91-92
Average Ages of Parents	31-32
Average Ages of Parents of Children Born in Rural Maryland, Tal	
XIII	.:. 32
Apparatus for Disinfecting Water	224
Bacteriological Examinations—	
For the Detection of Diseases and Insanitary Conditions, 7	Гя~
ble I	164
Of Foods, Number of	164
Of Milk	104
Of Overtone	164
Of Water Name of	164
Of Water, Number of	164
Baltimore City—	
Birth Rates, 1905-1915, Table II	8
Deaths by Months, 1910-1915, Table XXVIII	76
Deaths in, by Ages, Chart 4	40
Deaths Per Ten Thousand Population from Pulmonary Tub	er-
culosis, White and Colored, Table XXIII	57
Estimated Population of, in 1915, Tables I and I-A	6-7
Number of Pinthe Deported in 1015	6-7
Number of Births Reported in 1915	8
Number of Deaths Reported in 1915.	36
Baltimore County, Area, Population and Reported Cases of Notifial	hle
Diseases	135-137
Beasley, A. K., Report of	192-196
Beitler, Dr. Frederic V. Report of	5-118
Berlin, Investigation of the Water Supply of	91.1
Belated Returns of Typhoid Fever from Frederick City Hospital	414
Billingsley, Dr. W. B., Report of	130
Diameter Mandard	197-199
Biometer, Maryland	106-113
Birth Rate—	
By Sex and Color, Rural Districts, Baltimore City and Mar	ry-
land, Table IV	11
Computed on Scientific Groups of Population, Table IV	11
In Garrett County	11
In Towns in Maryland, Table XVI	35
Birth Rates—	100
Baltimore City, 1905-1915, Table II	8
Counties of Maryland 1915 Table III	0

	PAGE
Death Rates and Rates of Increase, Counties of Maryland, Ta-	
ble III	8
Maryland, 1905-1915, Table II	8
Rural Districts, 1905-1915, Table II	. 8
Birth—	00.05
Registration	32-39
Returns, Counties of Maryland Having Defective	11
Returns, Counties of Maryland Having Good	11
Births—	. 10
By Counties, and Increase of Population, Table III-A	
Table V	15
By Sex and Color in Towns of Maryland, Table XVI	
By Sex and Color, Rural Districts, Baltimore City and Mary	-
land, Table IV	. 11
Colored, Reported in 1915	. 8
Deaths_and Increase of Population, Counties of Maryland, Ta	-
ble III-A	
Deaths and Rates, Maryland (1914) Summary, Table IV	. 11
Female, by Months, Chart 1, Table V	14-15
Female, Reported in 1915	. 8
Illegitimate, by Counties, and Color, Table VII	. 19
In Maryland	. 8-35
Male, by Months, Chart 1, Table V	14-15
Male, Reported in 1915	. 8
Number of, Attended by Physicians and Midwives in Counties o	. 33
Maryland, Table XIV	
Number of, Recorded in Maryland in 1915	
Number of, Recorded in Rural Districts in 1915	
Percentage of, Attended by Physicians and Midwives, Table XV	
Percentage of, Returned in Maryland	
Reported in Baltimore City in 1915	. 5
Reported in Maryland in 1915	. 5
Reported in Rural Districts, 1915	. 5
White, Reported in 1915	8
Bottled Water, Permits Issued for Sale of	. 207
Bottled Water, Supply Investigations	. 204
Braddock Heights, Investigation of the Water Supply of	
Bradley Hills, Investigation of the Water Supply of	. 213
Investigation of Water Supply at	209
Sewerage Investigation at	. 215
Bright's Disease—	
Age, Distribution of Deaths from, by Percentages, Chart 14	. 68
Chronic, Deaths from, Table XXI, Chart 7	51-52
Deaths by Months from, Chart 18	77-78
Bronchitis (Acute), Deaths from, 1907-1915, Chart 16, Table XXVI	.70-71
Broncho-Pneumonia, Age, Distribution of Deaths from, by Percen	t-
ages, Chart 15	
Brunswick, Investigation of the Water Supply of	. ZII
Bureau of Bacteriology, Report of the Chief of	7 <del>1-</del> 100 36-160
Bureau of Communicable Diseases—	70-109
Prosecutions by	. 163
Report of the Acting Chief of	19-163
Bureau of Sanitary Engineering—	
Organization of	00-201
Report of the Chief of	00-230
Bureau of Vital Statistics Report of the Chief of	.5-118

n.o.
PAGE
Calvert County, Area, Population and Reported Cases of Notifiable
Diseases
Cambridge, Oyster Bed Investigations at
Canned Goods, Chemical Analysis of
Canned Goods, Chemical Examination of
Canned Tomatoes, Results of Examinations of182-191
Canneries—
Character of Floors in
Conditions Found in
County, Percentage of Increase of, Complying with the Law 194
Disposal of Tomato Skins at
Drainage at
Kind of Toilets in
Number of Inspected. 192
Operating in 1915
Ordered to Correct Insanitary Conditions
Washing Facilities in
Nashing Facilities III
Cannery Inspector, Report of
Caroline County, Area, Population and Reported Cases of Notifiable
Diseases
Carroll County, Area, Population and Reported Cases of Notifiable
Diseases
Cases Examined for Diagnosis—
By Counties
By Months and Diseases
Cases of Sickness from Notifiable Diseases in the Counties of Mary-
land, 1911-1915 120
Caspari, Charles, Jr., Report of
Causes of Death—
Classification of, with the Number Occurring and Their Ratios
to Mortality, Table XX, Chart 6
Fifteen Principal, Number and Percentage of, in Each of Three
Age Periods, Table XXIV
Principal
Principal, Classified According to Their Importance in the Mid-
allo Dovid of Life Chart 10
dle Period of Life, Chart 10
Cecil County, Area, Population and Reported Cases of Notifiable Dis-
eases
Cerebral Congestion and Hemorrhage, Deaths from, Table XXI,
Chart 7
Cerebral Hemorrhage—
Age Distribution of Deaths from, by Percentages, Chart 12 66
Deaths by Months from, Chart 18
Charles County, Area, Population and Reported Cases of Notifiable
Diseases
Chemical Analysis—
Of Dairy Products 167
Of Oils and Fats 167
Of Canned Goods 167
Of Soft Drinks
Of Vinegars
Of Extracts
0f Drugs
Of Miscellaneous
Of Waters
Of Sewage
Chemical Laboratory, Investigation Made by
Chief Chemist, Report of
Chief Cloub Deposit of

	PAGE
Age of Parents of, Table XII	31
Born in Rural Maryland, Average Ages of Parents of, Table	
XIII	32
Born of Native and Foreign-Born Parents, Table VIII	20
Classification of Causes of Death, with the Number of Occurring and	
Their Ratios to the Mortality, Table XX, Chart 64	7-49
Classification—	
Of Public Water Supplies in Maryland, Table III	222
Of Treated Institutional Water Supplies, Table IV	
Clerical Report	
Colored Births Reported in 1915	8
Colored Population, Higher Death Rate in, Chart 22	110
	35
Color, Births by, in Towns of Maryland, Table XVI	99
Comparative Susceptibility of the Several Organs to Parasitic Infec-	04
tions—Maryland Mortality, 1915, Chart 19	81
Comparative Table of Survivorship in Maryland, 1915, White and	
Colored, Table XXXV-B, Chart 22107	
Comparison of Fecundity Rates Between White and Colored Races	26
Comparison of Work Done by Engineering Division During 1914 and	
1915	202
Condemnation of Food Products	
Conditions Found in Canneries	
Congenital Debility, Deaths from, Table XXI, Chart 75	1-52
Congenital Diseases and Malformations, Deaths from, Table XX,	
Chart 64	7.40
Constitutional Dyscrasia, Deaths from, Table XX, Chart 64	7.40
Convulsions (Infantile), Deaths from, Table XXI, Chart 7	1 50
Co-operative Bacteriological Laboratory, Report of	169
Counties—	
Births, Deaths and Increase of Population in, Table III-A	10
Cases Examined for Diagnosis by	
And Months, Reported Cases of Notifiable Diseases by	
Having Good Registration of Notifiable Diseases	122
Having Poor Registration of Notifiable Diseases	122
Illegitimate Births in, by Color, Table VII	19
Maximum and Minimum Ages of Fathers and Mothers in, White	
and Colored, Table XI2	8-30
Mean Age at Death by, Table XXXIII	104
Number of Births Attended by Midwives and Physicians in,	
Table XIV.	33
Percentage of Native and Foreign Rorn Parents of Children	66
Percentage of Native and Foreign-Born Parents of Children Born in the, Table VIII	20
Public Health Conditions by	
Reports of Divorces by, Table XXXIX	-109
	114
Counties of Maryland—	
Birth Rates, Death Rates and Rate of Increase in, Table III	· 8
Deaths Per Ten Thousand Population from Pulmonary Tuber-	
culosis, White and Colored, Table XXIII	57
Having Defective Birth Returns	11
Percentage of Births Attended by Physicians and Midwives in	
the, Table XV	34
Reports of Notifiable Diseases from	121
Status of Notification of Infectious Diseases in the	122
Table of Maximum Fecundity in, Table X	27
County Canneries, Percentage of Increase of, Complying with the	
Law	194
Crownsville State Hospital, Investigation of Mumps at	197
Dairies Inspected, Number of	197

PAGE
Dairy Inspector, Report of
Dairy Products, Chemical Analysis of
Data Concerning Water Supply Disinfecting Plants, Table V226-227
Death—
Average Age at. Method Used for Obtaining91-92
Classification of the Causes of, with the Number Occurring and
Ratios to Mortality, Table XX, Chart 647-49
Mean Age at, by Counties, Table XXXIII
Mean Age at, by Months, Table XXXIV
Principal Causes of
Principal Causes of, Classified According to Their Importance in
the Middle Deviced of Life Chant 10
the Middle Period of Life, Chart 10
Special Causes of, 1906-1915, Table XXX82-88
Twenty Principal Causes of, Table XXI, Chart 751-52
Deaths—
By Ages, Table XXXVI-B 111
By Ages, Maryland Rural Districts and Baltimore City, Table
XVIII, Charts 3 and 4
By Counties, Table III-A
By Months, from Bright's Disease, Chart 18
By Months, from Cerebral Hemorrhage, Chart 1877-78
By Months, from Diarrhoea, Chart 18
By Months, from Heart Disease, Chart 18
By Months, from Influenza, Chart 18
Dy Months, from Thursday, Charl 10.
By Months, from Pneumonia, Chart 18
By Months, from Tuberculosis, Chart 18
By Months, from Typhoid Fever, Chart 1877-78
By Months, Rural Districts and Baltimore City, 1910-1915,
Table XXVIII 76
By Sex and Color, Estimated Population and Death Rate in
Towns of Maryland, Table XVIII
De See and Calon Devel Districts De Visco City 1 1 16
By Sex and Color, Rural Districts, Baltimore City and Mary-
land, Table IV
By Suicide, 1905-1915, Table XXX
From Accidental Violence, Table XXI, Chart 751-52
From Accidental Violence, 1906-1915, Table XXX
From Accidental Violence, Age Distribution of, by Percentages,
- 101 Precidental Florence, rige Distribution of, by Leftenburges,
Chart 14

	PAGE
From Drowning, 1906-1915, Table XXX	84
From Fifteen Principal Causes, Rural Districts, Number and	
Percentage of Death in Each of Three Age Periods, Table	
XXIV	0-62
From Gastric Diseases, Table XXI, Chart 7 5 From Influenza, Table XXI, Chart 7	61-52
From Influence Table XXI Chart 7	1-52
From Influenza, 1907-1915, Chart 16, Table XXVI	0-71
From Intestinal Parasites, 1906-1915, Table XXX	84
From Malignant Neoplasms, Table XX, Chart 64	17.49
From Malignant Neoplasms, Age Distribution of, by Percent-	. 10
ages, Chart 12	66
From Meningitis, Table XXI, Chart 7	31.59
From Occupational Poisonings, 1906-1915, Table XXX	83
From Organic Heart Disease, Table XXI, Chart 7	CI 59
From Organic Heart Disease, Age Distribution of, by Percent-	11-02
From Organic Heart Disease, Age Distribution of, by Tercent-	69
ages. Chart 15	
From Panalysis, Table AAI, Chart I	68
From Paralysis, Age Distribution of, by Percentages, Chart 14	80
From Parasitic Diseases, Table XXIX	7 40
From Parasitic Diseases, Table XX, Chart 6	11-48 10-71
From Preumonia, 1907-1915, Chart 10, Table AAVI	(4 EU
From Pneumonia, Table XXI, Chart 7	2G-1(
From Pheumonia, Age Distribution of, by Percentages, Chart 19.	ยย 40
From Poisonings and Intoxications, Table XX, Chart 6 4 From Pregnancy, 1906-1915, Table XXX	61-39
From Pregnancy, 1906-1915, Table AAA	0 40
From Pregnancy, Table XX, Chart 6	13-49
From Pulmonary Tuberculosis—Maryland, 1908-1915, by Sex,	72
Table XXVII. From Rabies, 1906-1915, Table XXX.	
From Rables, 1906-1915, Table AAA	83
From Senile Debility, Table XXI, Chart 7	01-02
From Smanpox, 1906-1915, Table XXX	85
From Tetanus, 1906-1915, Table XXX	70.71
From Tuberculosis, 1907-1915, Chart 16, Table XXVI	10-71
From Tuberculosis, Age Distribution of, by Percentages, Chart	
13	67
From Tuberculosis, by Ages, Deaths Per Ten Thousand of Pop-	
ulation at Age Periods, Per Cent. of Deaths in Each Age	
Period, White and Colored, Table XXII, Charts 8 and 9	) <del>4-</del> 96
From Tuberculosis, Pulmonary and Laryngeal, Table XXI,	
Chart 7	01-02
From Typhoid Fever, Table XXI, Chart 7	)1-52
From Typhoid Fever, Age Distribution of, by Percentages,	0.7
Chart 13	67
From Violence Table XX Chart 6	86
From Violence, Table XX, Chart 6	18-49
From Whooping Cough, Table XXI, Chart 7	01-92
From Whooping Cough, Age Distribution of, by Percentages,	
Chart 13	67
In Maryland	601-6
Number of Benefit in Beltimer City is 1015	30
Number of, Reported in Baltimore City in 1915	36
Number of Reported in Rural Districts in 1915	36
Number of, Showing Comparative Susceptibility of the Several	0.4
Organs to Parasitic Infections, Chart 19	81
Number Reported in Maryland in 1915	36
Per Thousand Population at the Several Age Periods, Table	0.4-
XIX-A, B, C, Chart 5	140
Per Thousand at the Age Periods, Table XXXVI-C	112
Per Ten Thousand Population from Pulmonary Tuberculosis,	

	AGE
Table XXIII	57
Registered in Baltimore City, 1915	5 5
Registered in Rural Districts, 1915.	5
Death Rate—	J
By Sex and Color, Rural Districts, Baltimore City and Mary-	
land, Table IV	11
Computed on Specific Groups of Population, Table IV	11
	110
In Garrett County	11
Death Rates—	
Counties of Maryland, Table III	8
Degenerations, Deaths from, Table XX, Chart 64	7-49
Diagnosis, Cases Examined for→	
By Counties	163
By Months and Diseases.	162
Diarrhoea and Enteritis, Deaths from, Table XXI, Chart 75	
Diarrhoea, Deaths by, Months from, Chart 18	
Difference Between Birth Rate and Death Rate, Table IV Diphtheria, Age Distribution of Deaths from, by Percentages, Chart	12
12	66
Diphtheria—	.00
At Hancock, Investigation of	128
	127
Diseases—	
	164
Cases Examined for Diagnosis by	162
Occupational, Reported, by Months	161
Parasitic, Deaths from, Table XXIX	80
Disinfection of Public Water Supplies in the State219-	
•	195
Distribution—	
	164
Of the Population and Deaths by Ages, Chart 5	$\begin{array}{c} 45 \\ 231 \end{array}$
Divorces—	201
	115
	115
Dorchester County, Area, Population and Reported Cases of Notifi-	110
able Diseases	143
	195
Drowning, Deaths by, 1906-1915, Table XXX	84
Drugs—	
Chemical Analyses of	
Not Passed, List of	181
Number of Samples Analyzed	
Duration of Life	109
XXXII-A, B, C, Charts 20-A, 20-B, 20-C94-	102
Elkridge, Investigation of Scarlet Fever at	128
Engineering Division, Comparison of Work Done by, in 1914-1915	
Estimated Population—	
And Death Rate in Towns of Maryland, Table XVII	37
In Maryland, 1915, Tables I and I-A	6-7
Of Baltimore City, 1915, Tables I and I-A	6-7
Of Maryland by Age Periods, Table XXXVI-A	
Of Rural Districts, 1915	2
Tables I and I-A	6-7

Examination—	PAGE
Of Canned Tomatoes, Result of	82-191
Of Oysters1	.71-173
Excess of Male Over Female Births, Chart 1	14
Expectancy, Presumptive, and Duration of Life, Rural District	ts,
Table XXXII-A, B, C, Charts 20-A, 20-B, 20-C	94-103
Extracts. Chemical Analyses of	168
Fecundity Rates, Comparison of, Between White and Colored Race	s. 26
Fecundity—	
Table of, for White and Colored Mothers by Age Periods, Table	.es
IX and IX-A, Chart 2	.22-26
Table of, Maximum, Counties of Maryland, 1915, Table X	17
Federalsburg, Investigation of the Water Supply of	209-210
Female Births by Months, Chart 1, Table V	.14-15
Female Births Reported in 1915	8
Female Births Reported in 1915	204
Financial Report	232-234
Floors, Character of, in Canneries	194
Food and Drug Inspectors—	101
List of Inspections Made by	179
Towns Visited by	178-179
Food and Drug Law, Prosecutions Under	176-177
Food and Drug Prosecution Cases, Settlement of	76-177
Food Products, Condemnation of	120-111
	100-101
Foods— Not Passed, List of	101
Not Passed, List of	171
Number of Samples Analyzed	
Frederick County, Area, Population and Reported Cases of Notifial	
Diseases	
Frederick City-Hospital, Belated Returns of Typhoid Fever from	130
Friendship Heights, Investigation of the Water Supply of	213
Garrett County—	
Area, Population and Reported Cases of Notifiable Diseases	
Birth and Death Rates in	
Gastric Diseases, Deaths from, Table XXI, Chart 7	51-52
Govans, Sewerage Investigation at	215
Harford County, Area, Population and Reported Cases of Notifial	ole
Diseases	146-147
Havre de Grace, Investigation of Scarlet Fever at	127
Heart Disease, Deaths by Months from Chart 18	77-78
Howard County, Area, Population and Reported Cases of Notifial	ole
Diseases	147
Hyattsville, Investigation of the Water Supply of	213-214
Ice, Chemical Analyses of	169
Ice Supply Investigations	204
Illegitimate Births by Counties and Color, Table VII	19
Illegitimate Births, 1915, Table VII	19
Increase in the Number of County Canneries Complying with t	
Law	194
Infant Mortality Rate—White and Colored, Rural Maryland, Tal	
XXXI	89
Influenza— Dootha by Months from Chart 18	77.70
Deaths by Months from, Chart 18  Deaths from, 1907-1915, Chart 16, Table XXVI	70.71
Deaths from Table VVI Chart 7	51 E0
Deaths from, Table XXI, Chart 7	01-02
Insanitary Conditions—  Bacteriological Examinations for the Detection of	164
Canneries Ordered to Correct	
Investigation of, by Bureau of Sanitary Engineering	
investigation of, by Dureau of Samuary Engineering	200

Inspection— PAGE
Of Canneries
Of Dairies
By Food and Drug Inspectors in Towns of Maryland178-179
List of, Made by Food and Drug Inspectors
Inspectors—Food and Drug, Towns Visited by
Intestinal Parasites, Deaths from, 1906-1915, Table XXX
Intoxications, Deaths from, Table XX, Chart 647-49
Investigations and Sanitary Surveys
Investigations Made by Bureau of Sanitary Engineering 203
Investigations Made by Chemical Laboratory
Investigations of Insanitary Conditions by Bureau of Sanitary En-
gineering
Jones, Dr. C. Hampson, Report of
Kent County, Area, Population and Reported Cases of Notifiable
Diseases
Kirkman, Walter N., Report of
Life—
Duration of91-105
Presumptive Expectation and Duration of, Rural Districts, Charts 20-A, 20-B, 20-C, Table XXXII-A, B, C94-103
Charts 20-A 20-B 20-C Table XXXII-A B C 94-103
List—
Of Foods and Drugs Not Passed
Of Inspections Made by Food and Drug Inspectors
Of Prosecutions Under Food and Drug Law
Of Samples of Foods and Drugs Analyzed
Male Births—
By Months, Chart 1, Table V14-15
Reported in 1915
Malignant Neoplasms—
Age Distribution of Deaths from, by Percentages, Chart 12 66
Doothy from Table VV Chart C
Deaths from, Table XX, Chart 6
Marriages, 1919, Table XXXVIII
Maryland—
Age of Parents of Children Born in the Rural Districts, Table
XII
Biometer
Births by Sex and Color in Towns of, Table XVI
Births in
Comparative Table of Survivorship in, 1915, White and Colored,
Table XXXV-B, Chart 22
Deaths from Tuberculosis in, by Sex, 1908-1915, Table XXVII 72
Deaths in
Deaths in, by Ages, Chart 3
Estimated Population of, by Age Periods, Table XXXVI-A 111
Estimated Population of, in 1915, Tables I and I-A
Number of Births Recorded in 1915
Number of Deaths Reported in 1915
Stillbirths in, by Months, Sex and Color, Table VI
Manyland Tuberculogic Sepitation, Chart 21, Table XXXV-A. 107-109
Maryland Tuberculosis Sanitarium, Investigation of the Water Sup-
Ply of
Maximum and Minimum Ages of Fathers and Mothers, White and
Colored, in Counties, Table XI
Maximum recundity, Table of, Counties of Maryland, Table X 27
Mean Age at Death by Counties, Table XXXIV
Most Lyguester Berent of
Meat Inspector, Report of

INDEX. 249

Meningitis, Deaths from, Table XXI, Chart 75	AGE
	1-52
Methods Used for Obtaining the Average Age at Death9	1-92
Midwives—	
Number of Births Attended by, in Counties of Maryland, Table	
	33
YIV  Percentage of Births Attended by, in Counties of Maryland,	
Table XV	34
	164
Minimum Ages of Fathers and Mothers, White and Colored, in Coun-	101
Minimum Ages of Fathers and Mothers, white and Colored, in Coun-	2 20
ties of Maryland, Table XI	1.00
Miscellaneous Chemical Examinations of Food Products	109
Miscellaneous Investigations by Bureau of Sanitary Engineering	204
Mitten, C. N., Report of	182
Montgomery County, Area, Population and Reported Cases of Noti-	
fiable Diseases149-	150
Months—	
	162
Deaths by, in Rural Districts and Baltimore City, 1910-1915,	
Table XXVIII	76
Mean Age at Death by, Table XXXIV	105
Reported Cases of Notifiable Diseases by	
Reports of Divorces by, Table XXXIX	
Reports of Occupational Diseases by	161
Returns of Marriages by, Table XXXVIII	114
Morse, Robert B., Report of	230
Morbidity—	200
Tuberculosis, by Color and Age Periods	195
Tuberculosis, by Counties, Sex and Color	197
Tuberculosis, by Months	104
Tuberculosis, by Sex and Color	124
Mortality—	
Age Distribution of, from Twelve Diseases by Percentages, Ru-	
ral Districts, Table XXV, Chart 116	66-4
Comparative Susceptibility of the Several Organs to Parasitic	
Infections, Chart 19	
infections, onary 100 and 100	81
From Five Respiratory Diseases, 1907-1915, Chart 16	81
From Five Respiratory Diseases, 1907-1915, Chart 16 From Five Respiratory Diseases, 1907-1915, Chart 16, Table	81 . 70
From Five Respiratory Diseases, 1907-1915, Chart 16  From Five Respiratory Diseases, 1907-1915, Chart 16, Table XXVI	. 70 . 72
From Five Respiratory Diseases, 1907-1915, Chart 16  From Five Respiratory Diseases, 1907-1915, Chart 16, Table XXVI	. 70 . 70 )-72 89
From Five Respiratory Diseases, 1907-1915, Chart 16	81 . 70 )-72
From Five Respiratory Diseases, 1907-1915, Chart 16	.70 .70 )-72 .89 .5
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 )-72 89 5
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 .72 .89 .5 8-40 3-78 74
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74 127
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74 127 212
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74 127 212
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74 127 212
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74 127 212 83 210 .127
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 3-78 74 74 127 212 83 210 .127
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 )-72 89 5 8-40 3-78 74 127 212 83 210 .127
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 )-72 89 5 8-40 3-78 74 127 212 83 210 .127
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 8-40 74 74 127 212 83 210 127 121
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 8-40 74 74 127 212 83 210 127 121
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 0-72 89 5 8-40 8-40 74 74 127 212 83 210 -127 121 120
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 .70 .72 .89 .5 .8-40 .3-78 .74 .127 .212 .83 .210 .127 .121 .120 .132 .134
From Five Respiratory Diseases, 1907-1915, Chart 16	81 .70 .70 .72 .89 .5 .8-40 .3-78 .74 .74 .127 .212 .83 .210 .127 .121 .121 .120 .132 .134 .137
From Five Respiratory Diseases, 1907-1915, Chart 16.  From Five Respiratory Diseases, 1907-1915, Chart 16, Table XXVI	81 .70 .70 .72 .89 .5 .74 .74 .127 .210 .127 .121 .120 .132 .134 .137 .138

PAGE
In Cecil County
In Charles County
In Dorchester County
In Frederick County144-145
In Garrett County145-146
In Harford County146-147
In Howard County
In Kent County
In Montgomery County
In Prince George's County
In Queen Anne's County
In St. Mary's County
In Somerset County
In Talbot County
In Washington County
In Washington County
In Wicomico County
In Worcester County
Notifiable Diseases, Reported Cases of, by Counties and Months 160
Notifiable Diseases, Reported Cases of, by Months and Counties 130
Notifiable Diseases, Reported Cases of, by Months 123
Number of Births Attended by Physicians and Midwives-White and
Colored—Rural Maryland, 1915, Table XIV
Number of Canneries Inspected
Number of Dairies Inspected
Occupational Diseases Reported by Months
Occupational Poisonings, Deaths from, 1906-1915, Table XXX 83
Oils and Fats, Chemical Examination of
Orders Issued for Construction of Sewerage and Water Supply Sys-
tems
Organic Heart Disease, Deaths from, Table XXI, Chart 751-52
Organization of the Bureau of Sanitary Engineering200-201
Oysters—
Bacteriological Examination of
Examination of171-173
Oyster Bed Investigations217-218
Paralysis—
Age Distribution of Deaths from, by Percentages, Chart 14 68
Deaths from, Table XXI, Chart 751-52
Parasitic Diseases—
Deaths from, Table XX, Chart 6
Deaths from, Table XXIX
Parents—
Average Ages of31-32
Native and Foreign-Born of Children Born in the Counties of
Maryland, Table VIII
XII
XII
Penniman, Dr. W. B. D., Report of
Permits Issued—
For Construction of Sewerage Systems
For Construction of Water Supply Systems
For Sale of Bottled Water
Percentage of—
Births Attended by Physicians and Midwives, Table XV 34
Births Attended by Physicians and Midwives, White and Col-
ored, Counties of Maryland, Table XV
Deaths from Pulmonary Tuberculosis at Various Age Periods,
Table XXII, Charts 8 and 954-56

	PAGE
Native and Foreign-Born Parents of Children Born in the Co	
ties of Maryland During 1915, Table VIII	20
Percentages of Total Mortality in Age Periods, Maryland, Rural I tricts and Baltimore City, Table XVIII, Charts 3 and 4	JIS- 29 40
Physicians—	00-40
Number of Births Attended by, in Counties of Maryland, Ta	ible
XIV	33
Percentage of Births Attended by, in the Counties of Maryla	nd,
Table XV	34
Plans for—	202 200
Sewerage Systems Received for Approval	
Water Supplies Prepared in Connection with Investigations.	
Water Supply Systems Received for Approval	.205-206
	60
Age Distribution of Deaths from, by Percentages, Chart 15 Broncho, Deaths from, Chart 16, Table XXVI	70-71
Deaths by Months from, Chart 18	77-78
Deaths from, Table XXI, Chart 7	51-52
Deaths from, 1907-1915, Chart 16, Table XXVI	70-71
Pocomoke City, Investigation of Smallpox at	127
Pocomoke City, Investigation of Water Supply at	214
Poisonings and Intoxications, Deaths from, Table XX, Chart 6	47-49
Population and Deaths per Thousand at the Several Age Perio	ods,
Table XIX—A, B, C, Chart 5	42-40
Estimated, Deaths by Sex and Color and Death Rate in Toy	arne
of Maryland, Table XVII	37
Estimated, of Baltimore City, 1915, Table I and I-A	6-7
Estimated, of Maryland in 1916, Table I and I-A	6-7
Estimated, of Rural Districts, 1915, Table I and I-A	
Pregnancy—	
Deaths from, Table XX, Chart 6	48-49
Deaths from, 1906-1915, Table XXX	84
Presumptive Expectation and Duration of Life, Rural Distri	ots,
Table XXXII—A, B, C, Charts 20-A, 20-B, 20-C Prince George's County, Area Population and Reported Cases of	No-
tifiable Diseases	150-151
Principal Causes of Death	50-88
Principal Causes of Death Classified According to Their Importa	nce
in the Middle Period of Life. Chart 10	59
Prosecutions by Bureau of Communicable Diseases	163
Prosecutions by Bureau of Vital Statistics	.116-118
Prosecution Cases, Food and Drug, Settlement of	176-177
Prosecutions Under Food and Drug Law  Public Health Conditions by Counties	190-111
Queen Anne's County, Area, Population and Reported Cases of N	oti-
fiable Diseases	.151-152
Rabies, Deaths from, 1906-1915, Table XXX	83
Report of—	
A. K. Beasley	192-196
Dr. Frederick V. Beitler	5-118
Dr. W. B. Billingsley	197-199
The Cannery Inspector	170 200
Dr. Chas. Caspari, Jr	164-165
The Chief Chemist	.166-169
The Chief of the Bureau of Communicable Diseases	.119-163
The Chief of the Bureau of Sanitary Engineering	.200-230
The Chief of the Bureau of Vital Statistics	5-118
The Chief Clerk	.231-234
The Co-operative Rectariological Laboratory	165

252 INDEX.

	PAGE
Dr. C. Hampson Jones	
Walter N. Kirkman	
The Meat Inspector	
C. N. Mitten	102 200 220
Dr. W. B. D. Penniman	166 160
Dr. W. R. Stokes.	164-165
The State Food and Drug Commissioner	170-200
The Veterinarian and Dairy Inspector	197-199
The Veterinarian and Dairy Inspector	123
Reported Cases of Notifiable Diseases, Rural Maryland, by Coun	ties
and Diseases	160
Reported Cases of Notifiable Diseases, Rural Maryland, by Mon	nths
and Counties	130
Reported Cases of Tuberculosis	121
Reports of—	
Divorces by Months and Counties, Table XXXIX	
Notifiable Diseases	.119 - 127
Occupational Diseases by Months	161
Respiratory Diseases, Mortality from Five, 1907-1915, Chart	16,
Table XXVI	70-72
Results of the Examinations of Canned Tomatoes	.182-191
Returns of Marriages by Months and Counties, Table XXXVIII.	114
Rural Districts—	41
Age Distribution of Mortality from Twelve Diseases in Table XXV, Chart 11	tne,
Birth Rates, 1905-1915, Table II	<del>04-</del> 65
Deaths from Pulmonary Tuberculosis in, 1908-1915, by S	· · · · · · · · · · · · · · · · · · ·
Table XXVII	$\frac{563}{1}$
Deaths in, by Ages Chart 4.	40
Deaths in, by Ages, Chart 4  Deaths by Months in, 1910-1915, Table XXVIII	$\frac{10}{76}$
Estimated Population of in 1915. Table 1 and I-A	-6-7
Fifteen Principal Causes of Death in the, Showing Number	and
Percentage in Each of Three Age Periods, Table XXIV	60-62
Infant Mortality Rate in Table XXXI	90
Mortality from Five Respiratory Diseases in the, Chart	16,
Table XXVI	70-72
Number of Births Reported in 1915	8
Number of Deaths Reported in 1915.	36
Samples of Foods and Drugs Analyzed, List of	174
Sanitary Engineering, Investigations Made by the Bureau of	203
Sanitary Surveys and Investigations	127
Elkridge, Investigation of	190
Havre de Grace, Investigation of	197.198
St. Inigoes, Investigation of	128
St. Mary's Industrial School, Investigation of	127
Western Maryland College, Investigation of	128
Seasonal Incidence of Mortality	73-78
Seasonal Incidence of Mortality	74
Seasonal Incidence of Mortality in the Rural Districts Chart 17	74
Senile Debility, Deaths from, Table XXI, Chart 7	51-52
Sewage, Chemical Analyses of	169
Sewerage Investigations	
Sewerage Investigations	204
Sewerage Investigations at— `	04-
Bridewell	215
Arlington	215
Govans Tolchester Beach	215
Sewerage Systems, Orders Issued for Construction of	210
orders assued for Construction Of	200

INDEX. 253

PAGE
Sewerage Systems, Permits Issued for Construction of 207
Sewerage Systems, Plans of, Prepared in Connection with Investigations
Sewerage Systems, Plans Received for Approval
Sex, Births by, in Towns of Maryland, Table XVI. 35 Smallpox, Deaths from, 1906-1915, Table XXX. 83
Smallpox, Investigation of at Pocomoke City 127
Snow Hill, Investigation of Water Supply of
Soft Drinks, Chemical Examinations of
able Diseases
Special Causes of Death, 1906-1915, Table XXX
St. Mary's County, Area, Population and Reported Cases of Notifiable Diseases
State Food and Drug Commissioner, Report of
Statement of Receipts and Disbursements232-234
Status of Notification of Infectious Disease, Counties of Maryland. 122
Status of the Counties of Maryland, Notifiable Disease Report 121 Stillbirths—
By Months, Sex and Color, Table VI
Maryland, 1915, Table VI
Stokes, Dr. W. R., Report of.
Survivorship by Ages, 1910-1915, Table XXXVII
Survivorship in Maryland, 1915, Comparative Table of, White and
Colored, Table XXXV-B, Chart 22
Colored, Table XXXV-B, Chart 22
Susceptibility, Comparative, of the Several Organs to Parasitic In-
fections, Chart 19
Of Fecundity for White and Colored Mothers by Age Periods
Tables IX and IX-A, Chart 2
Of Maximum Fecundity, Counties of Maryland, 1915, Table A 27 Showing Maximum and Minimum Ages of Fathers and Mothers
Table A128-50
Talbot County, Area, Population and Reported Cases of Notifiable
Diseases
Toilets, Kind of, in Canneries
Tolchester Beach, Sewerage Investigation at
Tomatoes, Canned, Results of Examinations of182-191 Towns of Maryland, Deaths in by Sex and Color, Estimated Popula-
tion and Death Rate, Table XVII
Towns Visited by Food and Drug Inspectors
Tuberculosis—
Age Distribution of Deaths from, by Percentages, Chart 13 67 Deaths by Months from, Chart 18
Deaths from, 1907-1915, Chart 16, Table XXVI
Morbidity by Sex and Color
Morbidity by Color and Age Periods
Morbidity by Months. 127 Prophylactic Supplies, Distribution of 231
Prophylactic Supplies, Distribution of
Pulmonary and Laryngeal, Deaths from, Table XXI, Chart 751-52 Pulmonary, Deaths by Ages, Deaths per Ten Thousand of Popu-
lation at Age Periods, and Per Cent. of Deaths in Each Age
Period, White and Colored, Table XXII, Charts 8 and 954-56 Pulmonary, Deaths from, 1908-1915, by Sex, Table XXVII73-74
Pulmonary, Deaths from, 1908-1915, by Sex, Table XXVII73-74 Pulmonary, Deaths per Ten Thousand Population from, White
and Colored, Counties of Maryland and Baltimore City, Table
XXIII

Reported Cases of	ŧΕ
	1
Twenty Principal Causes of Death, Table XXI, Chart 751-5	2
Typhoid Fever—	
Age Distribution of Deaths from, by Percentages, Chart 13 6	37
At Belair, Investigation of	8
At Chase, Investigation of	28
At Hagerstown, Investigation of	8
At Maryland Tuberculosis Sanitarium, Investigation of 12	8
At The Rocks, Investigation of	8
Deaths by Months from, Chart 18	Q
Deaths from, Table XXI, Chart 7	9
Decrease Council by Division With a Western Western	2
Decrease Caused by Disinfecting Water 22	9
Unspecified or Ill-Defined Causes, Deaths from, 1906-1915, Table	
Table XXX 8	
Veterinarian, Report of197-19	9
Vinegar, Chemical Analyses of	8
Violence, Deaths from, Table XX, Chart 648-4	9
Vital Statistics	
Washing Facilities, Character of, in Canneries	1
	4
Washington County, Area, Population and Reported Cases of Noti-	
fiable Diseases	6
Water—	
Apparatus for Disinfection of	4
Number of Bacteriological Examinations 16	4
Water Supplies—	
Classification of Treated Institutional, Table IV 223	3
In Maryland, Classification of, Table III	2
In the State, Disinfection of	U
Plans of, Prepared in Connection with Investigation 200	6
Water Supply Investigation of—	
Bridewell	
Bradley Hills	3
Berlin	
Braddock Heights210-213	4
	1
Brunswick	$\frac{1}{2}$
Brunswick	$\frac{1}{2}$
Brunswick	1 2 3 4
Brunswick	1 2 3 4
Brunswick       211-212         Friendship Heights       216         Hyattsville       213-214         Federalsburg       209-216         Maryland Tuberculosis Sanitarium       212-215	1 2 3 4 0 3
Brunswick	1 2 3 4 0 3
Brunswick       211-212         Friendship Heights       216         Hyattsville       213-214         Federalsburg       209-216         Maryland Tuberculosis Sanitarium       212-215	1 2 3 4 0 3 2
Brunswick       211-212         Friendship Heights       218         Hyattsville       213-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-218         Myersville       212         New Windsor       216	1 2 3 4 0 3 2
Brunswick       211-212         Friendship Heights       218         Hyattsville       213-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-218         Myersville       215         New Windsor       216         Pocomoke City       214	1 2 3 4 0 3 2 0 4
Brunswick       211-212         Friendship Heights       215         215       115         Hyattsville       209-216         Maryland Tuberculosis Sanitarium       212-215         Myersville       215         New Windsor       216         Pocomoke City       214         Snow Hill       214	1 2 3 4 0 3 2 0 4 4
Brunswick       211-212         Friendship Heights       215         215       115         Hyattsville       209-216         Maryland Tuberculosis Sanitarium       212-215         Myersville       215         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227	1 2 3 4 0 3 2 0 4 4 7
Brunswick       211-212         Friendship Heights       215         Hyattsville       213-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-215         Myersville       212         New Windsor       210         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       204	1 2 3 4 0 3 2 0 4 4 7 4
Brunswick       211-212         Friendship Heights       215         Hyattsville       213-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-215         Myersville       212         New Windsor       210         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       204         Water Supply Investigations       209-214	1 2 3 4 0 3 2 0 4 4 7 4
Brunswick       211-212         Friendship Heights       218         Hyattsville       213-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-218         Myersville       212         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water-Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       204         Water Supply Investigations       209-214         Water Supply Systems—	1 2 3 4 0 3 2 0 4 4 7 4 4
Brunswick       211-212         Friendship Heights       218         Hyattsville       213-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-218         Myersville       212         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water-Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       204         Water Supply Investigations       209-214         Water Supply Systems—       0rders Issued for Construction of       208	1 2 3 4 0 3 2 0 4 4 7 4 4 8
Brunswick       211-212         Friendship Heights       215         215       115         Hyattsville       209-216         Federalsburg       209-216         Maryland Tuberculosis Sanitarium       212-215         Myersville       212         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       204         Water Supply Investigations       209-214         Water Supply Systems—       0rders Issued for Construction of       208         Permits Issued for Construction of       206-207	1 2 3 4 0 3 2 0 4 4 7 4 4 8 7
Brunswick       211-212         Friendship Heights       215         215       115         Hyattsville       209-214         Federalsburg       209-216         Maryland Tuberculosis Sanitarium       212-215         Myersville       212         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       204         Water Supply Investigations       209-214         Water Supply Systems—       070-216         Orders Issued for Construction of       208         Permits Issued for Construction of       206-207         Plans of, Received       205-206	1 2 3 4 0 3 2 0 4 4 7 4 4 8 7 6
Brunswick         211-212           Friendship Heights         215           Hyattsville         203-214           Federalsburg         209-210           Maryland Tuberculosis Sanitarium         212-215           Myersville         212           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         0rders Issued for Construction of         208           Permits Issued for Construction of         206-207           Plans of, Received         205-206           Waters, Chemical Analyses of         106	1 2 3 4 0 3 2 0 4 4 7 4 4 8 7 6
Brunswick         211-212           Friendship Heights         215           Hyattsville         203-214           Federalsburg         209-210           Maryland Tuberculosis Sanitarium         212-215           Myersville         212           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         0rders Issued for Construction of         208           Permits Issued for Construction of         206-207           Plans of, Received         205-206           Waters, Chemical Analyses of         106	1 2 3 4 0 3 2 0 4 4 7 4 4 8 7 6 9
Brunswick       211-212         Friendship Heights       215         Hyattsville       203-214         Federalsburg       209-210         Maryland Tuberculosis Sanitarium       212-215         Myersville       212         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       209-214         Water Supply Investigations       209-214         Water Supply Systems—       0rders Issued for Construction of       208         Permits Issued for Construction of       206-207         Plans of, Received       205-206         Waters, Chemical Analyses of       106         White Births Reported in 1915       58	1 2 3 4 0 3 2 0 4 4 7 4 4 8 7 6 9
Brunswick       211-212         Friendship Heights       215         215       115         Hyattsville       209-216         Maryland Tuberculosis Sanitarium       212-215         Myersville       215         New Windsor       216         Pocomoke City       214         Snow Hill       214         Water Supply Disinfecting Plants, Data Concerning       226-227         Water Supply Investigations       209-214         Water Supply Investigations       209-214         Water Supply Systems—       0rders Issued for Construction of       208         Permits Issued for Construction of       206-207         Plans of, Received       205-206         Waters, Chemical Analyses of       106         White Births Reported in 1915       8         Whooping Cough—       8	1234032044744 87698
Brunswick         211-212           Friendship Heights         215           Hyattsville         209-214           Federalsburg         209-215           Maryland Tuberculosis Sanitarium         212-215           Myersville         212           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         0rders Issued for Construction of         208           Permits Issued for Construction of         206-207           Plans of, Received         205-206           Waters, Chemical Analyses of         109           White Births Reported in 1915         8           Whooping Cough—         Age Distribution of Deaths from, by Percentages, Chart 13         67	1234032044744 87698 7
Brunswick         211-212           Friendship Heights         215           Hyattsville         213-214           Federalsburg         209-214           Maryland Tuberculosis Sanitarium         212-215           Myersville         212           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         0rders Issued for Construction of         208           Permits Issued for Construction of         208-207           Plans of, Received         205-206           Waters, Chemical Analyses of         109           White Births Reported in 1915         8           Whooping Cough—         Age Distribution of Deaths from, by Percentages, Chart 13         67           Deaths from, Table XXI, Chart 7         51-52	1234032044744 87698 7
Brunswick         211-212           Friendship Heights         215           11 Hyattsville         213-214           Federalsburg         209-210           Maryland Tuberculosis Sanitarium         212-215           Myersville         212           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         0rders Issued for Construction of         208           Permits Issued for Construction of         206-207           Plans of, Received         205-206           Waters, Chemical Analyses of         106           White Births Reported in 1915         8           Whooping Cough—         Age Distribution of Deaths from, by Percentages, Chart 13         67           Deaths from, Table XXI, Chart 7         51-52           Wicomico County, Area, Population and Reported Cases of Notifiable	1234032044744 87698 72
Brunswick         211-212           Friendship Heights         215           Hyattsville         209-216           Federalsburg         209-217           Maryland Tuberculosis Sanitarium         212-215           Myersville         215           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         00 rders Issued for Construction of         208           Permits Issued for Construction of         206-207           Plans of, Received         205-206           Waters, Chemical Analyses of         106           White Births Reported in 1915         8           Whooping Cough—         Age Distribution of Deaths from, by Percentages, Chart 13         67           Deaths from, Table XXI, Chart 7         51-52           Wicomico County, Area, Population and Reported Cases of Notifiable         Diseases         157	1234032044744 87698 72
Brunswick         211-212           Friendship Heights         215           11 Hyattsville         213-214           Federalsburg         209-210           Maryland Tuberculosis Sanitarium         212-215           Myersville         212           New Windsor         216           Pocomoke City         214           Snow Hill         214           Water Supply Disinfecting Plants, Data Concerning         226-227           Water Supply Investigations         209-214           Water Supply Investigations         209-214           Water Supply Systems—         0rders Issued for Construction of         208           Permits Issued for Construction of         206-207           Plans of, Received         205-206           Waters, Chemical Analyses of         106           White Births Reported in 1915         8           Whooping Cough—         Age Distribution of Deaths from, by Percentages, Chart 13         67           Deaths from, Table XXI, Chart 7         51-52           Wicomico County, Area, Population and Reported Cases of Notifiable	1234032044744 87698 72 7






TABLE A.

DEATHS IN BALTIMORE CITY, 1915, BY AGE, SEX, COLOR, MONTH AND CAUSE.

Jacousty.  March. May. May. Jude. September. October. November. Devember. Pennie. Pennie. Pennie. Pennie.		2 3	2 to 3.	3 to 4.	5 to 10	10 to 15, 15 to 20.	5	25 to 30.	\$	40 to 45. 45 to 50.	2	55 to 60. 60 to 65.	12	2	75 to 80. 80 and over.	Unknown.	Toral.
8 6 5 6 5 9 13 16 20 11 16 7 71 51 85 37 122	1. Typhoid fever		1		1 9	6 1	6 17	21 5	e  e	11	9 5	3	1 1	2	1		122
8 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2. Typhus fever																
	3. Relapsing fever 4. Malaria		:: :::				:   : : :		:   : : :		::::::	i	:: :::				
	5. Smallpox,																
1 2 1 1 1 1 1 1 3 7 9 10 15 4 19 2 1 1 1 2 4 13 10 20 3 23	6. Measles. 7. Scarlet fever	7 1	2 2	1	2 5	3	:   : : :	· i   · · i									19 23
5 2 2 3 2 2 1 3 3 10 11 13 8 21	S. Whooping cough	12	6 1								:: :::						21
14 5 3 4 2 2 2 4 6 5 5 12 20 35 61 3 64			17 10	11	5 11		1 3	3 3	1 1	11	7			1::1			64 97
10 8 22 28 10 5 1 3 10 40 57 71 26 97	11. Miliary fever				: *								12 9	11		1:::	91
	12. Aslatic cholera				:   : : :												
	14. Dysentery						i	2			::   :::		2 1	1:::1	2 3	1:::1	12
	15. Plagne																
	17. Leprosy.				:   : : :						:: :::1						
2 3 3 1 2 1 1 1 1 1 9 7 16 16	18. Erysipelas. 19. Other epidemic diseases.	6 .	] ]						1 1	1	1		2 2		1	(	16
	20 Purulcut infection and septichemia		i		· · · · i	· · i	2	2	1 2		3 4		3				29
	21. Glanders																
	22. Anthrax. 23. Rabies.				: ::::			::: ::			:: :::	::: :	:: :::	: :::		}	
1 1 1 1 1 3 1 2 7 4 8 3 11	24. Tetaous.		1			2	1 1	1									11
	25. Mycoses. 26. Pellagra.				. 2	1:::1::		::: ::		:::			i				4
	27. Beribert. 28. Tuberculosis of the lungs					12 7											
	29. Acute miliary tuberculosis		3 3	2	5 12		3 94	111 9	4 3		03 49	2 .			4 1	:::	821 50
10 6 7 2 6 4 6 4 7 2 1 3 30 28 45 13 58	30. Tuberculosis meningitis	11	11 3		4 7		4 4		2 1		1		1	1			58
	31. Abdominal tuberculosis		1		2 2		1	6	4 4	3 .	3 3		2		1	:::	43 8
1 1 1 1 1 3 2 2 4	33. White swelling	1 .							1			1 .		. 1			4
	34. Tuberculosis of other organs		1	i	. 1		1 2	``i	1 2		i	1 .	i	11			9
1 1 1 1 3 2 1 4 5	36. Rickets	3	1		1				1								5
	37. Syphills		5 3		. 1		2 5	4	6 1		3 3	4	2 1				102
1 2 2 2 4 2 2 4 1 2 19 3 21 1 22	39. Cancer and other malignant tumors of the									2	3 3	3	5 1				22
17 19 21 19 22 16 17 21 18 17 20 19 103 123 208 118 220	40. Cancer and other malignant tumors of the								1 1		-	-		1 1	3	***	22
	stomacb, liver						. 1	2	6 3	14 5	21 27	27	46 27	7 21	17 14		226
11 5 5 3 0 4 8 6 3 7 10 6 34 40 67 7 74	41. Cancer and other malignaut tumors of the peritoneum, rectum, intestines							2	1 7	4	3 9	10	8 1	5 9	5 1		74
3 6 14 9 8 3 8 16 9 3 4 5 88 78 10 88	42. Cancer and other malignant tumors of the						1 3				15 10	17			0 -		00
4 7 8 2 4 6 1 4 7 3 7 53 48 5 53	female genital organs						1 3	1	3 6		15 10	17	, ,	5 5	2 5		88
	breast							2	2 5	3	8 11	4	3 2	2 3	4 6		53
2 1 1 4 2 3 2 1 11 5 16 16	44. Caucer and other malignant tumors of the skin,		1							1 .	3	1	2	3	2 3		16
11 8 18 8 12 11 9 8 13 8 9 6 82 39 113 8 121	45. Cancer and other malignant tumors of						5 1	3	3 6	- 11	14 13	- 1	17 15				
	other organs, and of organs not specified		1	1			- 1		3 6	12 :	19 18	14	17 12	12	6 1	• • •	121
	tal organs excepted)						. 1						1		1		3
	47. Acute articular rheumatism							3					2 3		2 2	:::	40 17
	49. Seurvy													1			

\*1 Japanese. †1 Chinese.



# TABLE A-Continued.

DEATHS IN BALTIMORE CITY, 1910, BY AGE, SEX, COLOR, MONTH AND CAUSE.

										_					_	_					1 1		1	-				_										
January.	February.	March.	April.	May.	June.	July.	August	September.	October,	Novembor	November.	December.	Male.	Female.	White.	Colored.	Total		1	0 to 1.	1 to 2.	2 to 3.	4 to 5.	5 to 10.	2	2	20 to 25.	2	35 to 40.	3	2	50 to 55.	60 to 65.	65 to 70.	2 .	75 to 80.	knov	Total
13	5	12		11	7	3	5	1				12	42	63	103		10		50. Diabetes.							2	2	2	3		7	14		5 15	9		5	105
			2 .								3 .		1	5	5 3				51. Exophthalmic goitre							1		1 :			1	1	2		:::			6
	2					4			1 2	2	1 .		9	4	13			13	53. Leuchæmia		1	1				1	1 .		2 1	2		1 .		2				13
3	2	1	4	1 .				1		1		1	7	9	14 2		1		54. Anæmia, chlorosis	1						1	1 .		1		1	::: .		2 1		1 .		16
	1	6		2	:::	2	2		: :	3	1	3	20	5	20	5	2	25	56. Alcoholism (acute or chronic)									1 :	2 3	6	8		3	i ''i	1			25
									:	2		1	3		3				57. Chronic lead poisoning			1										1 .						3
		· ;	::: :		:::	···i					:: ::			2	1	····i	1	2	58. Other chronic occupation polsonings 59. Other chronic poisonings																	::: :	::   :::	2
1							1		j :	1			2	1	3			3	60. Encephalitis											1								3
-1	2 .		5	2		5		3			i	3	18	22	34				61. Meningitis	12	9	4						:: ::		2		9	2	1			:: :::	40 10
2	1	1 .		2	1	5	1			4 .		3	9	11	16			20	(3. Other diseases of the spinal cord	1	1	2		2		1	1 .		2		2	1 66		1 3	1		2	20
70	67	57	56	54	39	41		40		1	56	65	207	323	476 5				64. Cerebral hamorrhage, apoplexy 65. Softening of the brain		1				1	1	1	2	8 13	22	43	66	77 7		82	80	66	620
8	6	10	5	5	2	1	10		5	2	4	5	24	39	49	14	6	63	66. Paralysis without specified cause							1		::	. 1	1	4	5	11	5 7	11	5	12	63
1	2	5		2 .		3							11.	6	15 3		1		67. General paralysis of the insane															4 2	2	1	1	17
1		1	1			2	1			3	1	2	5	9	9		1		69. Epilepsy	1	1				2		1	1	2 1	. 1		1	1		1	1.		14
				1 .		2	····i	٧٠٠٠		3			18	12	1 22			30	70. Convulsions (nonpuerperal)	95																		30
6			1.						١,,,					1			ď		72. Chorea	20		1			1					1				. 11.5.		::: :		1
2		2 .											16	3 12	26			28	73. Neuralgia and neuritis						1		···i	2		1	1		1	i				4
6	1	1 .		*						°		2	10	12	26		1 2		74. Other diseases of the nervous system 75. Diseases of the eyes and their annexa																1:::1	::: :		28
	3	1	1	1			1	3				2	6	7	10 5		1	13	76. Diseases of the ears	1	2	1		. 2		1	1 .		1	2		1 .		. 1			.,	13
1 9	2	10	10	3 .	7	9	3			1	7	7	45	41	65		8	86	77. Pericarditis					1 1	4	"i	5 .	8	5 12	6	15	11	10	2	1	i	2	9 86
100	62	100	76	77	50	53	61			1	83	82	410	467	679				79. Organic diseases of the heart					1 8	14	11	14	24 2	3   35	51	59	69	77 11	3 106	101		87	877
10	1 8	6	5 14	8	4 7	2 14	2 15			7	17	3 25	28 74	15 71	36 120			43 45	80. Angina pectoris										1 2	1	-1	5	8	7 4	5	3	3	43
- 1	- 1									.	4				17				rysm, etc										1 1					1 14			40	145
3	1		1 .		:::	1						2	12	8	17		2		82. Embolism and thrombosis						1 1				1 2	2	2	1	4	. 2	3	2	1	20
					- 1		1						2			١.			rholds, phlebitis, etc.)										. 1	1								2
		,	1,.	• • •	-1				4	.	**		2	1	2	1		3	84. Diseases of the lymphatic system (lymphangitis, etc.)		1		1		1						1							3
				1 .		1	1					1	1	3	2	2		4	85. Hæmorrhage; other diseases of the circu-																			
										. l									latory system	1																		4
		1 .					1						6		6				87. Diseases of the larynx	2			1						1		1				1			6
10	10	ii	5	1 .	5	5	····i			· · ·	7	7	30	3 42	2 54		7	72	S8. Diseases of the thyreoid body	30								i				5	2	1				3 72
11	12	6	8	4	3	6	2	1 1	1	3	5	3	21	33	42	12	5	54	90, Chronic bronchitis	1					1 1			1	1	1	2	6	2	6 5	7		14	54
46	69 53	61 96	68	54	29	- 22 15	27 15	25 27	2:		33 60 :	48 84	243 350	261 255	340 419				91. Bronchopneumonia	216 61		30 10		2 8		5			9 8 6 40			11 52		2 24 6 46	18	24	18	504 605
4	2		114			2		2	:	1	2	3	10	8	11	7	1	18	92. Pneumonia			1				2	1 .		, 3	2	4	1	1	1	2			18
7	3	6	3 .	···i	1 9			2		1	3	4	13	17	28		3	30	94. Pulmonary congestion, pulmonary apoplexy 95. Gangrene of the lung.								1	1	1		1			3 3		2	7	30
3		3 .		î.				1		1	i	1	8	6	. 9		1		96. Asthma.				: ::	: :::	1			::  **	i	1	1	2		3 3	1	1	i	14
	1			``i .			····i	1		i			1	1	1 5	. 1		2	97. Pulmonary emphysema																1			2
1		1	1	1 .			,		1.	1	1	1	3	o	9	4		B	98. Other diseases of the respiratory system (tuberculosis excepted)	1		1				1				1	1	1	1	. 1	1			9
1		1 .	i :						:::		2	1 2	2 5	3	9	1			99. Diseases of the mouth and annexa	3	1																1	5
1																1::::			100. Diseases of the pharynx																			1
*1 Chir	iese.			_													-	-						_	- 1			-					-		-			

\*1 Chinese.



## TABLE A-Continued.

DEATHS IN BALTIMORE CITY, 1915, BY AGE, SEX, COLOR, MONTH AND CAUSE.

January.	February.	March.	Aprili.	May.	June.	July.	August.	September.	October,	November.	December.	Male.	Female.	White.	Colored.	TOTAL.			0 to 1.	1 to 2.		5 to 4.	2	10 to 15,		2 .	35 to 40.	2	45 to 50.	3 2	60 to 65.	2	75 to 80.	Uoknown.	Torat.
3 .		1	3	2 .	[	2	1	2	4	3	2	15		18	5	23	102. Ulcer of the stom	ach						1	2	1	4	5 1	1	5 2	1	. 1			23
11	3	7	6		7	8	5	- 1	5		4	32		45	24		cepted)	the stomach (concer ex	. 11	1		1	1	2	1 1	3	1 :	5 4	2	2 3	6	7 5	10	3	69
11 5	15	11	15	10	28	106	107	84	59 6	17				399 60	80 23		105. Diarrhoa and ente	eritis (under 2 years) eritis (2 years and over)		86	11	4	1 6	1	1 2	2	2	3 7	4	6 5	7	3 6	4	8	479 83
																		,\s																	
6	4	7 8	6	4 12	9	13	6	6	3	2	4			60 67	10 18	70	108. Appendicitis and	typhlitis					7	7 7	8 8	5 5	5 5	8 5	4	6 3 8 5	3	5 2			70 85
5 1 .				1 .		1 .				2				4	1	5	110. Diseases of the in	testines								1	1			1		1		1	5
												1					112. Hydatid tumor of	phy of the liver the liver												1					4
9	2	7	2	8 2 .	9	6	2	9		1	6	52	13	61 16	9	17	114. Biliary calculi	ver						:   : : :   :			1	5 7		13 11 5 4	2		2	1	70 17
1	1	1	5	2	2	3	1				1	8	14	19	3			the liver						: ::: : :				1		1 1				3	22
1 .		1	1	1	1 .		4		i	2		7	4 2	10	1 2			(nonpuerperal)  f the digestive system	. 1				1	. 1	1	1	1			2	1				11
			11	11	8	12	6	2	11		. 9	60		76	42		(cancer and tub	erculosis excepted)				i .	9		4 :			0 15	100	1	12	7 5	[] ·		3 118
17 75	80	79	88	85	57	59	66	56	76	72	70	116	417	676	*187	863	120. Bright's disease				1	1 .		. 1	1 10	21	22 4	0 57	79 1	07 108	119 1	11 70	66	49	863
1 .		1 .		2	3	2	5		3	1	2	8	12	17	8		122. Other diseases of	the kidneys and annexa.	.   2		1	1 .		:	1	. 1		2 2	1	2 1	1 .	1	2	i	20
··; :			1	1 .		1	1	2	2	2	1	5	7	2 11	1	12	124. Diseases of the b	nary passages	. 1	3						1		. 2				2	1	1	2 12
	2		"i :		8		i	1		1	1 2		- 1	26				ethra, urinary abscess, et rostate														2 10		7	3 29
											1	1		1			127. Nonvenereal disen	ses of the male genita	1							1									1
(),																	128. Uterine hæmorrha	ge (nonpuerperal)																	
		1 .		2	1	1	1	2	1	1	1		11		2	13	130. Other diseases of	oncancerous)the uterus						. 1	1 :	1 2	4	1		1					11 11
1		i		· · ·	1		1	2		····i	- i			11	6		132, Salpingitis and o	umors of the ovary ther diseases of the fe	-					. ]							2 .				4
																	male genital org	ansass of the breast (can							1 :	1 4	2	2 5	2 .						17
1	1	2	9	2	1			1	- 1		1		12	a	2	15	cer excepted)	nancy									·· 6 ··				:::   <u>:</u>				12
1 .	[	2	1	1 .		1	1	1	1				9	9			135. Puerperal hæmorr	hage							:	2 2	3	1 1							9
	1	3	3	2 .		3 .		3	1	2	1		22	17	6	28	137. Puerperal septicha	f lahoremia							3 8	2	5	5						::: :::	23
1 :			1			1 2								20 5	3		139. Puerperal phiegma	uria and convulsions isla, alha dolens, embolus								2					-				23
																		rth (not otherwise de			1				1 :	1		2 1							6
																	fined),	of the breast										: : : : :							
1 .		3	1	2	2			1		1			6 2	8	6	14	142. Gangrene		. 1	1					1			1 1		2	1 .	3	3	1	14
1 .			1 .		2				1		· · · · i	4	2 3	3	3 2		144. Acute abscess			2						1	1		1	1 1					6
i,				4 .		1	1	3	2	4				21	2		146. Diseases of the	the skin and annexa bones (tuberculosis ex																1	02
			1 .		1								2	2		2	147. Diseases of the ;	joints (tuberculosis and	1	1	2 .			1 2			- 1	1						.,	23
																	rheumatism exce	pted)												1		: :::			2
		hinese						•••••									140. Other diseases of	the organs of locomotion																	



## TABLE A-Concluded.

DEATHS IN BALTIMORE CITY, 1915, BY AGE, SEX, COLOR, MONTH AND CAUSE.

	1		1	1					7		Ī	1							Τ	Ī		T	T		1	TI			T					T	.1 1	
.   5							ber.	١.,	1	. je	j j		,												25.	88	35.	j 10	50.	55.	.09		12		over wn.	
January. February.	March.	į į	l in	9	l is	August.	tem	October	Vomomb	line.		Male.	Female.	White.	Colored.	TOTAL		to 1.	2,	60	to 4.	50 50	0 1	2 2	to 2	2	2   3	2 2	to 5			to 6		5	30 and Unknov	14.
Jan	Ma	April.	May.	June.	July.	Au,	Sel	Oct	12		5	Ma	Fe	¥	Co	To		0 0	7	12	80	4 1	2 3	15	ន	25	8 8	8 8	5 5	25	55	8 8			8 g	Η̈́
4	3	4 7		5   10	1	31 4	1	4 :	7	3	4	35	23	47	11	58	58 150. Coogenital malformations (stillbirths not																			
05 0		47 46	5	5 43	4	0 25	, ,	1 9.	4	97	32	972	210	381	102	489	laciuded)	483			1		1 .													58 483
10	9	21 8	1	1 12	1	2 14	i	5 10	0	17	14	94	59	114	39	153	53 152. Other diseases peculiar to early infancy	153																		153
10 1	2	1 I 10 11				7 5		5 6	1 · ·	4	12	33	61	80 .	5 14		6 153. Lack of care						:: :	:: ::			.:: :	:: :		: 'i		::: -	4 17		50	91
2	3	2		3 5		1 4		4 3	3	1	1	14	15	27 24	2		29 155. Suicide by poison					:::			3 6	3 4	3	5	3	3 1	2	]				29 24
1		2 1					٠				2	3	3	6			6 157. Suicide by hanging or strangulation					,			. 1		1 .		1							6
1 9	1	5 2				i		5			2	30	1	29	2	31	3 158. Suicide by drowning.									6	1	1 .		3 4	2					31
1	ī					. 2				'		5					5 160. Suicide by cutting or pierciog instruments 3 161. Suicide by jumping from high places									1	2 .		'	. 1		1 .				5
																	162. Suicide by crushing									.   ,					11	! .				
	; ···		1			i	100		i ::		i	1	3				163. Other suicides						il:	:: ::	: ::::		::: :	:: :			1	1			i	
1		1 1	1 :	1 2		. 1				2		7	2	9			9 165. Other acute poisonlogs. 5 166 Conflagration.	1	. 2	i	2				. 1	l		1 .		. 1	1				··i	9
10	6	3 9	· "·	4 2			3	4 10	0	9.	14	35	48	60	23	88	83 167. Burns (conflagration excepted)	2	8	9	7	9	7	5	2	3	2	4	1 .	5 5	2	2	· 8 · · :		i :::	83
2		1 1		. 2	2	1 1		. :	1		1	8	2	8	2		10 168. Absorption of deleterious gases (conflagra- tion excepted)	- 4	1	٠	1								1	1	9	1 .				10
2	3	2 2	2 3	3 10	1	1 11		4 :	5	4	1	57	1	42 4	16	58	58 169, Accidental drowning 7 170. Transatism by firearms			1:::			7	2	2 (	7	6	8	2 1	0 2	3	2 .	:	i	÷::	58
1						1					::	1	1	1	1		2 171. Tranmatism by cutting or piercing instru-				1 1		1	1	,	1			1						1	7
15	6	7 2		3 4	ı ı	3 7		6 9	9	8	10	50	41	80	11	91	ments.  1 172 Třaumatism by fail.									i	3			. 4			9 6		10	91
	1		1									1		1		1	1 173. Tranmatism in mioes and quarries									1										1
4 1	4 (	11 7		6 6	3	$\frac{1}{5}$ $\frac{2}{11}$	i	j 1	1	13	10	97	12	101	8		10 174. Traumatism by machines	1			1		1			2 2	1	1 .				2 .				10
			l	١,		. 1	Ú.,.					3		2	1	5	railroad, landslides, etc.)			. 3			9		5 5	8	13	8	5		11	4	6	1		109
																	. 177. Starvation																			
1		::   : : : :				7 2						6	6			19	1 178. Excessive cold	3	''i	i :::				i			i i		::  ''	i	2		::  '';	1 1	i	1 12
	i								٠.,								1 180. Lightning. 1 181. Electricity (lightning excepted)																			····i
		2 4				1 2		1 :	2	2	2	13	7	12	8	20	20 182. Homicide by firearms							1	1 4	1 6	2	2 .		. 3	1					20
2	1	2 5	5	$\begin{vmatrix} 1\\2 \end{vmatrix} = \begin{vmatrix} 2\\2 \end{vmatrix}$		i		i '':	2	1		5	3 14	2 16	3		8 183. Homicide by cutting or plercing instruments 19 184. Homicide by other means					:::				3 3	i	4		2						8
	1	3				. ' ' i			3	3		18		16	1	1	1 IS5. Fractures (cause not specified)										i			. 1		.			· i	
				1									1		1	1	1 187. Ili-defined organic disease										1 .									1
· · · i		1 2		2 1	•••	2 5		3	1	1	3	8	17	2 14	11	25	2 188. Sudden death	6		3						i   :::									'i :::	2 25
849 73	9 98	52 899	76	4 632	70	4 740	68	0 72	1 7	755 8	31 4	1835	1492	7149	2178	9.327		-		119						870	371 4					-		482		9,327
	- 0	000	'0			, 10	30		,			-	-102		-*10	0,021	TOTAL	1024	200	110	10	20				,,,,,		40		1020	300	9		702		0,021




TABLE B

DEATHS IN COUNTIES OF MARYLAND, 1915, BY AGE, SEX, MONTH AND CAUSE.

January. Rebrusty. March. April. May. June. August. Suptember. October.	November. December. Male. Female. Torax.		1002. 2004. 2004. 2006.
		1. Typhoid fever	2 3 3 2 2 15 18 34 27 21 11 16 12 10 7 8 3 3 3 1 1 2
7 12 8 6 11 10 15 36 29 28	19 21 115 87 125 77 202	2. Typhus fever	
		3. Relapsing fever 4. Malaria	
1 2 2 1 3 1		5. Smallpox,	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 1 5 9 10 4 14 7 7 29 20 46 3 49	6. Measles. 7. Scarlet fever	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
8 10 7 2 10 7 2 9 4 3	4 14 36 44 46 34 80	8. Whooping cough	40 19 4 5 2 6 3
9 10 13 1 5 4 1 5 7 16	24 22 58 60 104 14 118 1 17 62 66 94 34 128	9. Diphtheria and croup	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
		11. Miliary fever	
	1 2 3 3	12. Asiatic cholera 13. Cholera nostras	
1 1	3 1 20 31 33 18 51	14. Dysentery.	9 2 3 3 1 1 2 1 1 1 3 2 5 4 7 5 1
		15. Plague,	
		16, Yellow fever	
3 1 2 3 3 4 1 3 3	2 1 15 11 23 3 26 1 2 1 2 3	18, Erysipelas.	2 1 1 1 1 1 2 2 1 1 1 1 2 4 1 4 1
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3 3 27 7 28 6 34	19. Other epidemic diseases	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
		21. Glanders	
		22. Anthrax. 23. Rabies	
2 1 3 1 2 1 5 2 1	3 2 15 8 12 11 23	24. Tetanns.	7 1 2 2 2 2 3 1 1 1 1
	1 5 6 5 6 11	25. Mycoses. 26. Pellagra.	
		27. Beriberi	
130 124 138 150 130 108 134 116 85 103 2 3 7 2 7 3 4 8 3 4	84 110 737 675 876 *536 1412 4 2 27 22 21 28 49	28. Tuberculosis of the lungs	10 5 · 3 1 1 13 36 112 199 173 163 137 130 95 81 75 65 48 32 20 6 7 14 1 1 4 3 11 5 5 5 5 1 2 1 3 3 1 1 1
2 4 3 6 7 6 4 5 2 3	4 1 26 21 33 14 47	30. Tuberculosis meningitis	8 18 2 1 4 2 5 1 1 1 1 1 2 1
9 5 1 4 5 5 4 2 6 1	1 3 21 25 25 21 46 2 2 6 5 10 1 11	31. Abdominal tuberculosis	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
1	1 1 1	33. White swelling	1
1 2 1 1 1		34. Tuberculosis of other organs	1 1 1 1 2 3 1 1 1 1
2 1 1 1 1 4	4 1 8 7 6 9 15	30. Rickets.	11 2 1 1
3 10 5 5 3 3 10 7 5 5	8 10 47 27 27 47 74 2 1 1 2	37. Syphilis. 38. Gonococcus infection	15 1 1 2 6 7 8 9 6 2 6 3 5 3
1 1 1 1 3 2 1 4 2 4	1 1 13 9 17 5 22	39. Cancer and other malignant tumors of the	
12 14 21 20 20 17 18 24 22 23	21 16 109 119 198 30 228	40. Cancer and other malignant tumors of the	1 2 2 1 3 3 5 2
		stomach, liver	
8 5 10 4 6 6 4 11 10 5	5 3 31 46 63 14 77	41. Cancer and other malignant tumors of the peritoneum, rectum, intestines	
10 8 6 3 9 8 3 12 5 7	4 9 84 72 12 84	42. Cancer and other mallgaant tumors of the	
1 6 9 2 2 5 0 5 9 5	5 8 1 66 58 9 67	female geuital organs	4 3 12 11 10 11 8 7 9 4 5
, , , , , , , , , , , , , , , , , , , ,		43. Cancer and other malignant tumors of the breast	1 2 3 4 10 6 14 8 4 7 6 2
2 5 7 2 3 1 4 3	2 2 23 8 29 2 31	44. Cancer and other malignant tumors of the	
12 9 6 5 12 7 7 8 5 12	7 9 66 33 92 7 99	45. Cancer and other malignant tumors of	
	1 1	other organs, and of organs not specified	1 2 2 2 2 2 1 2 3 2 5 3 3 14 20 13 10 5 8 1
	1 1 1	46. Other tumors (tumors of the female geni- tal organs excepted)	1
1 2 1 1 2 1 2 3 3 5 7 4 4 4 1 1 1	5 6 16 11 18 9 27 15 11 18 8 26	47. Acute articular rheumatism	1 1 3 1 2 1 4 2 1 4 1 2 2 2
1 1 2 1		48. Chronic rheumatism and gout.	4 3 3 1 3 1 2 2 3 2 6 3
*One Indian.			



# TABLE B-Continued.

DEATHS IN COUNTIES OF MARYLAND, 1915, BY AGE, SEX, MONTH AND CAUSE.

January.	February.	March.	April.	May.	June.	July.	August.	September.	October,	November.	December.	Male.	Female.	White.	Colored.	TOTAL		0 to 1.	1 to 2.		3 to 4.	3 2	10 to 15,	15 to 20.	2	25 to 30.	35 to 40.	2	3 .	50 to 55.	60 to 65.	2	70 to 75.	3 8	Unknown.	Torat.
8	3	7	6	7	7	7	9	8	4				48			8	50. Diabetes				::::			1	2	2	4 1	5	7	9 1	5 12 1 1	15	3	3 3		81 8
					;	1			1			1 2	····.	10	3	1	2 52. Addison's disease 53. Leuchæmia.				:::::::::::::::::::::::::::::::::::::::									1	i			1		2 13
	1		3	2	3	4	3	ĩ	î			1.	10	· 21	3	2	4 54. Anæmia, chlorosis					1	l			4	1 1	1	1	2	6 2	1	3 .			24
	1		1 0		-1	1 2		1		1 2	I	48		32			6 55. Other general diseases, 56. Alcoholism (acute or chronic)	3			1   .			i	1	2	8 8		4		1 · · · 2	1		1		6 46
						1						1		1			1 57, Chrohic lead poisoning												1 .		!					1
								::::				1	7	9			58. Other chronic occupation poisonings 59. Other chronic poisonings		1								2						1 .			
27	2	1		2 .		1	2	1		2	1	10	4 31	11 42	3		4 60. Encephalitis. ,	3	1	1						1	1 2		2	1	i					14
7   2	10		1	2	5	4 7	10 2	4			3						61. Meniogitis. 62. Locomotor ataxia											2	1	4 :	3 2	· · ;	1 -	2		57 13
3	-1	5	10	5	2	4	3	5	1		3 2			36	11	4	63. Other diseases of the spinal cord	3	4	4	1 .	3	3 1	4			. 2	-22	2	2 51 6	3 2 5 3			3 , 3		47
56	62	64	64	64	60	57	44	50			i i	-	7	565 10		71			1:::		::: :		: :::					10	33	51 6	6 87		121	98 99	5	712 11
12	16	18	19	24 12	11	13		20 6				98		160 68		20° 8°	66. Paralysis without specified cause				:::		. 1	1		1		+	6	6	7 27			37 52 6 4		207 89
5 8	6	2	12	3	3	3	3	6	3	2	2 5	21	26	43	4	4	68. Other forms of mental allenation				:				2	1	5 5	8	5		3 2	2	4	5 1		47
2	7	4	5	7	3	5	-1	1	. 5	1 2	2 8	32	21	35	18	5	89. Epilepsy	1			:::1:		1 5	7	1	8	5 3			5	2 4	2	2	1 1	1	53 3
7	11	7	6	8	9	7	8	11	3		111	58		51		9.	70. Convulsions (nonpuerperal)	83	7	1		1										1				92
	2	1				i						3	1 3	3			3 72 Chorea				:::1:											···i				3 5
	i	2	2	3	1	2	4			1 8	3 2	16	13			2	73. Neuralgia and neuritis				1 .	:								1			ĩ .	2		29
1									i			1	5	7	3	· · · i	75. Diseases of the eyes and their annexa				::: :					i ::								· ; · · ·		10
2	···i	1	2				3	1		1	1 2	8	6	10	4	1	77. Pericarditis.					:	2 1			1			1	1	1 1	3	1 .	1		14
86	87	92	83	102	74	2 87	2 88	69	66							97	78. Acute endocarditis	- 1			:::		9	10	5	10 1	3 5 1 93		47		6 2 5 101		157 1	1 47 124	ii	37 978
6	7	7	3	4	2	7	2	7	5	7	3	38	21	56	4	6	80. Angina pectoris										1 2		3		9 7		16	6 7		60
26	15	28	21	7	7	10	8	14	20	8	3 11	105	70	137	38	17								1:1			1	1	3	6	5 13	16	40	33 57		175
4	3	4	2	2	6	1	1	2				13	22		11	3.								1			i	2	2	4	4 3	3				35
	1				3	1			1				- 4	-1			83. Diseases of the veins (varices, hæmor- rhoids, phlebitis, etc.)										11					1	1	1		4
							1					1		1			1 84. Diseases of the lymphatic system (lym-																			
			1	.			1					1	1	1	1		phangitis, etc.)	1																		1
																	latory system								1									1		2
			1									1111	3	6			86. Diseases of the pasal fossæ		2	1	1 .			1			. 1			1				:: :::	1:::	6
13	10	27				2		1				58	53				88. Diseases of the thyreoid body	****						1			. 2		1		1 1			ii 13		5 111
6	10	15	9	4	í		6	3	-1		3	20	37	50	16	11				-4	1 .			1:::		::			3		3 3	7	6	9 24		66
53 77	63 66	76 110	94 102	32 32	22 17	16 12	17	28 11		30						50	91. Bronchopneumonia	216	55	14	7	9 '	7 2	2	1	3			7 15		7 19 3 47			35 50 34 46		506 571
3	1	1	1	1	3	ĩ							8	14		57. 1		72		6		7 10		16	10	29 2	27				1 1		1	1		17
2	6	8	3	2	2	2	3				. 4	21	15	34	2 2	3	94. Pulmonary congestion, pulmonary apoplexy	8		1		:	1		1			1			. 2		6		:::	36 3
3	3	2	2	· i	i	· · · i	1	4	1		5 5	14	15	24 24		2	95. Gangrene of the lung 96. Asthma.														2 5	6	4	4 6		29
		2	5	***	3	1 3		1	3					15	1		97. Pulmonary emphysema													2	1	1	1	2		8
	- 1	- 1	"			3			,	1 '	. 2	19	8	13)	1	2	98. Other diseases of the respiratory system (tuberculosis excepted)	1	1					14.		1	2 1	2	3	3			3	5		22
					1			1	3			5	3	4	7		1 99. Diseases of the mouth and annexa	11																		11 17
	1											1	8	10		1	7 100. Diseases of the pharynx	1		1:::				:::			1					1:::		2		i
		*One	Chin	ese.				-			-						, and the adopting and the transfer					-		-				_			_	-	-			



#### TABLE B-Continued.

DEATHS IN COUNTIES OF MARYLAND, 1915, BY AGE, SEX, MONTH AND CAUSE.

				T	Т	-	1			_										1	[		T	T	1	T	1	T 1	_	1			. 1	
							Der.		er.	er.		:		,								10	9 4		35.	. t		15	٠,	نہ ا		ایا	over.	
ary.	February	ਰੂਂ :	i .	ته		August.	Septembe	October,	Novembe	December	le,	nale	White.	Colored.	TOTAL		to 1.	to 2.	to 3.	to 4.						to 40.				to 70.		to 80.	30 and over	ij
Janu	Febr	March	April.	June.	July.	Aug	Sep	Oct	No	Pe	Male.	Fer	W	Col	To	_	0 t	7.		2 4	10	10 to	2 2		8	335		50	55	8 8			88 15	Tor
1		3	3	3 2					4	3	9	14	18	5		102. Ulcer of the stomach		}				1	1	ı	2	2	5	. 1	1	2	3 2	2	]	23
9	5	11	6	9 10					5	-4	30	38	45			103. Other diseases of the stomach (cancer excepted).	3		1 .		. 4	1	3	3 2	4	3	2	2 7	1	6	4 9	6	7	68
6	9	9	17	9 28 7 5	122			118	11	10			464 136	212 31	167	104. Diarrhea and enteritis (under 2 years) 105. Diarrhea and enteritis (2 years and over).			34	7	5 8	1	1	2 1		···i	4	i	3	15 1	1 19	19	22	676 167
												1			1	106. Ankylostomiasis,		1						: :::		::: :	:: ::	:						····i
3 6	4 6	7 7	7 5	5 4 6 3	5 9	5	10	3	7	7	29 39		36 52	13 22	74	108. Appendicitis and typhlitis		2	1	1	. 3	3	4 2	8 3	5 6	2 2	2	7 1 3	7	3	2		1 1	49 74
1	2	2						::::			5 2	4 1	8 2	1		110. Diseases of the intestines	1			:: ::	:			. 2	2	::: :		1 1		1	i i		1	9
		7								5	24	31	· · · · ·	11	50	112. Hydatid tumor of the liver						1:::	i	i		3	4	5 6		5	6 7			55
3 .		i				3	2	····	1 2	2	11	8 8	11 16	3		114. Biliary calculi										1		2 1	-	2	. i	4	î	12 19
					2	3					13		18	9	2	116. Diseases of the spleea					.   1					1 .					1 3			2 27
		1		i							2		3	1		118. Other diseases of the digestive system (cancer and tuberculosis excepted)																	1 1	
11 82	16 63		15 06 7	7 8 9 65	14 73			11 75				70 419	*97 696	53 234	150 930	119. Acute nephritis 120. Bright's disease	9	5	5	1	5 5	4	5	7 5 7 10	5	5	12	7 12 0 71	14	11 1	1 10	7	5	150 930
													12	2														i i						14
															1	123. Calculi of the urinary passages																	5	1 19
			٠ ا										16			125. Diseases of the urethra, urinary abscess, etc. 126. Diseases of the prostate	2												1					
				.						1				1		127. Nonvenereal diseases of the male genital organs.										- 1				1				,
			1										3			128. Uterine hamorrhage (nonpuerperal) 129. Uterine tumor (noncaocerous)						1							1					
	2		1							1		4 8			4	130. Other diseases of the uterus							1	1		1	1	i 2						4
i	i i			2	3			1	1			10	5		10	132. Salpingitis and other diseases of the fe-					1										. ,	1		10
		1										1	1		1	male genital organs															1			10
	1			4	2			2				11 17	10 14	1	11	cer excepted)								1 3 3 3 2	1	2	2							11
2	1 .	1	1	1 2		1 2			 2 2			11	6 29	5	11	135. Pnerperal hemorrhage				'				1 5 9 6	1	4 .								17 11
8	4	5 2	3	3 3	1		2		3	3		37 28	21	8	28	137. Puerperal septichæmia								7 5				i			: :::			37 28
						1						3	1	2		139. Puerperal phlegmasia, alba dolens, embolus, sudden denth		١							2		1							3
												4	2	2		140. Following childbirth (not otherwise defined).						. 9.		1	1	1	1							4
	5	6	4	3	2	3		1	1				28	3	31	141. Puerperal diseases of the breast				1									1	1	1 4	8	13 1	31
1 .	1	1 .		1				1	1		1 2	3	4	····i	1	143. Furuncle. 144. Acute abscess					. 1			1				2			1			5
2 .			i							2	7		3 8	3		145. Other diseases of the skin and annexa 146. Diseases of the hones (tuherculosis ex-	3			- F					)	1								4
																cepted)	2			1								. 2					1	11
																rheumatism excepted)														}				
			1	-		1				or unit	2		2		2	149. Other diseases of the organs of locomotion															. 1	1		2

\*Oue color unknown, †One Chinese.

• ٠.

TABLE B-Concluded.

DEATHS IN COUNTIES OF MARYLAND, 1915, BY AGE, SEX, MONTH AND CAUSE.

January. February.	March.	April.	May.	June.	July.	August	September.	October.	November.	December.	Male,	Female.	White.	Colored.	TOTAL		0 to 1.	1 to 2.	2 to 3.		# to 3.	10 to 15.	15 to 20.	2	25 to 30.	35 to 40.	40 to 45.	2 .	50 to 55. 55 to 60.	60 tn 65.	2	70 to 75.	lkn an	Toral.
2 2 1 1 1 8	12 12 13 14 12 13 14 15 15 16 16 16 16 16 16 16 16 16 16 16 16 16	9 9 1 1 1 2 2 2 3 3 1 1 2 2 6 6 2 2 3 2 2 4 4	6 53 6 19 2 2 1 1 3 13 13 3 3 3 3 3 3 3 3	55 51 9 2 2 11 11 1 1 1 2 2 2 1 3 3	14 49 10 10 1 10 1 10 1 1 1 1 1 1 1 1 1 2 2 1 1 1 1	8 8 57 111 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	5 64 13 3 3 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 6 70 2 2 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	8 60 7 177	9 488 122 1 1 244 4 4 2 2 4 4 1 1 1 1 1 1 1 1	522 333 76 84 99 22 60 300 23 11 31 15 23 31 129 15 2 46 67 10 134 9 68 84 21 88 55 20 29	388 363 388 7 120 120 22 23 34 4 22 23 37 30 12 11 11 17 7 7 6 7 7 5 3 6 6 7 7 5 3 6	766 453 82 2 2 139 14 4 4 4 5 7 7 166 6 6 16 1 7 9 131 7 4 6 9 4 20 10 11 18	183 32 163 63 63 63 63 63 63 63 63 63 63 63 63 6	966 6366 114 155 204 14 14 14 14 15 19 9 17 7 7 60 6 6 141 16 9 9 17 7 7 9 9 4 28 28 25 25 25	150. Congenital malformations (stillbirths not included).  151. Congenital debility, icterus, and selerema.  152. Other diseases peculiar to early infancy.  153. Lack of circ.  155. Suicide by polyon.  155. Suicide by paphyxia.  156. Suicide by paphyxia.  157. Suicide by paphyxia.  158. Suicide by paphyxia.  159. Suicide by paphyxia.  150. Suicide by paphyxia.  150. Suicide by purpuing from high places.  151. Suicide by travening.  152. Suicide by travening.  153. Suicide by travening.  154. Suicide by travening.  155. Suicide by travening.  156. Suicide by travening.  157. Suicide by crashing.  158. Other acute poisonings.  159. Other acute poisonings.  150. Other acute poisonings.  150. Configuration.  151. Taramation of deleterious guess (configuration).  155. Absorption of deleterious guess (configuration).  156. Acute of the configuration of the configuration.  157. Traumation by free free free free free free free fre	\$88 6336114411515114115151515151515151515151515	1 2 1 1 1 1 1 1 1	3 3	1 1 1 5 5	1	1 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3			3	3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 1 1 5 2 2 1 4 4 1 1 1 1		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1	1 2  1 1 1 1 1 1 1	23 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	. 686 . 114 . 114 . 120 . 14 . 4 . 4 . 4 . 9 . 3 . 3 . 3 . 1 . 1 . 1 . 1 . 1 . 1 . 60 . 6 . 6 . 141 . 1 . 26 . 7 . 7 . 60 . 9 . 7 . 7 . 60 . 9 . 7 . 7 . 7 . 60 . 9 . 7 . 7 . 7 . 7 . 9 . 9 . 1 . 1 . 1 . 22 . 22 . 32 . 33 . 34 . 33 . 34 . 33 . 34 . 33 . 34 . 34
974 948		-	952		23 1023	1067	962	981		-	6380	_	*152 8618	173 3465	_	189. Cause of death not specified or ill-defined	2238	380	145	84	_	-	323	_				-1-			-		983 8	


\*Sex and color unknown.

.

TABLE C.

DEATHS IN MARYLAND, 1915, BY CAUSES, SEX, COLOR AND COUNTY.

					DEATH		III ZX ZX			,					COLC																
Male,	Female.	White.	Colored.	Total, Rural Dis- tricts.	Causes of Death.	Baltimore City.	Allegany.	Anne Arundel.	Baltimore.	Calvert	Caroline.	Carroll.	Cecil.	Charles.	Dorchester.	Frederick.	Garrett	Harford.	Howard,	Kent.	Montgomery.	Prince George's.	Queen Aone's.	Somerset	St. Mary'a.	Talbot	Washington.	Wicomico.	Worcester.	and Balto, City.	Percentage.
115	87	125	77	202	1. Typhold fever	122	21	9	17	2	5	5	5	10	19	7	4	6	5	8	3	9	5	6	4	10	17	7	18	324	1.52
					2. Typhus fever																::::										
4	10	7	7	14	3. Relapsing fever	i		1			1			2			1					2	1		5			'i :		15	0.07
					5. Smallpox,								2																		
6 29	9 20	10 46		14	6. Measles. 7. Scarlet fever	19 23	7	3	23	1	1		1		4	2	1	2		2	::::	1	1	1	- 4	1			1	33 72	0.15 0.34
36	44	46		80	8. Whooping cough	21	3	3	6	2	1		4			8	8	5		2	2	î	î	3	13	2	5	6	1	101	0.47
58	60	104		118	9. Diphtheria and cronp	64 97	11 14		31	2	7 3	8	3	3		7 9	5	4 2	. 1	1	2 11	11	i	2	. 1	5 4	10	5		182 225	0.85
62	66	94	34	128	10. Infineuza. 11. Miliary fever				7.4																		4		- 1	220	1.06
					12, Asiatie cholera			,																							
1 20	31	3 33		3	13. Cholera postras	12	1 2	····	7		2		2	····i	6	2		····i	1	2	3		2	···;	····i	1	1	6		63	0.01
20	91		18	51	14. Dysentery. 15. Plague,																								· · · · ·		0.00
					16. Yellow fever																							].			
15	11	23		26	17. Leprosy, 18. Erysipelas	16	4	3	7		····i			····i	1	2	ı i					2			1					42	0.20
1	2	1	2	3	19. Other epidemic diseases											1									2					3	0.01
27	7	28	6	34	20 Purulent infection and septichemia	29	3	1	9				. 2		4	2	2		1	1	2		1	1	1	1	1		2	63	0.30
					21. Glauders. 22. Anthrax.																						:::				
					23. Rabies,																	:									
15	8	12		23	24. Tetanus	11	1	2	3			1	2		2	1			1	::::		::::				1	1	2	2	34	0.16
5	6	5		11	25. Mycoses. 26. Pellagra,			2	5													i			1	····i		il.		16	0.003
			1113		27. Beribert			61		;:	25	****	28	31	66	106				33	46		33	1000							101111
737 27	675 22	876 21		1412 49	28. Tuberculosis of the lungs	821 50	44 6	1	500 19	15	25			31	2	3	12	31 3	16		20	57	33	55	42	32	50		45	2233	10.46 0.46
26	21	33	14	47	30. Tuberculosis meningitis	58	2				2				3		2		1		1		2				3		3	105	0.49
21 6	25 5	25 10		46	31. Abdomical tuberculosis	43	3		10		2	1	1	1		3	····i	1		2	4	1	1	1		1	1	4	1	89	0.42
1		10		11	32 Pott's disease																	::::		1	····i					19	0.09
5	7	9	3	12	34. Tuberculosis of other organs	9	1		2			1			3	1									1		1			21	0.10
8	3	3	3 9	6 15	35. Disseminated tuberculosis			···;	1 3			2		····;						2	1	1						``i ;		1·4 20	0.07
47	27	27		74	37. Syphilis.	102	6	8	37		1			ī		2					1					1	3	6	1	176	0.82
13	2	. 1	1	2	38. Gonococcus infection				1							• • • •		• • • •										1 .		2	0.009
10	9	17	5	22	39. Cancer and other malignant tumors of the buccal cavity	22		1	11			3			1						1.		1			1	2	1 .		44	0.21
109	119	198	30	228	40. Cancer and other malignant tumors of the			1												,			-			1	~	11			0.21
31	46	63	14	77	stomach, liver	226	26	7	60	1	4	12	10	3	7	15	6	8	2	3	10	14	6	8	2	7	16	4	4	454	2,13
0,	10	00			peritoneum, rectum, intestines	74	11	4	25	1		5	3	1		8	2	2	2	1	1	5		1		1	2	2		161	0.71
	84	72	12	84	42. Caucer and other malignant tumors of the										i																
1	66	58	9	67	female genital organs.  43. Cancer and other malignant tumors of the	88	5	2	22		1	6	4		3	12		3	• • • • •		41	6	3	2		1	7	3 .	•••	172	0.81
					breast	53	2	2	19		1	7	3			9		3	2	1	2	5		1		4	6	1 .		120	0.66
28	8	29	2	31	44. Cancer and other mallgoant tumors of the	10										2								0			-				
66	33	92	7	99	45. Cancer and other malignant tumors of	10	2		7		1	1	2		1	3								3		2	5	3	1	47	0.22
				3	other organs, and of organs not specified	121	18	3	20		1	5	6		3	6	2	7	5		3	4	3	2	1	3	6	1 .		220	1.03
	1	1		1	46. Other tumors (tumors of the female geol- tal organs excepted)	3					1																				0.02
16	11	18		27	47. Acute articular rheumatism	40	4	4	6			· · · i	1			i		2		1		4	1			1	1	:		67	0.02
15	11	18		26 4	48. Chronic rheumatism and gout	17	2	3	4			1	1	3		2	1	1	1	1										43	0.20
- 0	1		1 0		49. Seurty				1											_ 1						1		1 .		4	0.02



# TABLE C-Continued.


DEATHS IN MARYLAND, 1915, BY CAUSES, SEX. COLOR AND COUNTY.

														,	-															
Male.	Female.	White.	Colored.	Total, Rural Dis- tricts.	CAUSES OF DEATH.	Baltimore City.	Allegany.	Anne Arundel.	Baltimore.	Calvert.	Caroline.	Carroll.	Cecil.	Charles.	Dorcheater.	Frederick.	Garrett.	Harford.	Howard,	Kent.	Montgomery.	Prince George's.	Queen Anne's.	Somerset.	St. Mary's.	Talbot.	Washington,	Wicomico. Worcester.	Total: Maryland and Balto, City.	Percentage,
9	14	18	5	23	102. Ulcer of the stomach.	23	4	1	6						2		1		1		2	1			1	1	1	2	46	0.22
30			23	68	103. Other diseases of the stomach (cancer excepted).	69	3	8	15	2	2	1	2	2	2	1		4	2	1	3	5	1	2	2	1	5	4	137	0.64
361 74	315 93		212 31	167	104. Diarrhoa and enteritis (under 2 years) 105. Diarrhoa and enteritis (2 years and over).	479 83	42 8	5	106 45	2	26 7	12 5	4	24 3	48 9	29 9	10 3	28 7	3	3	19 7	44 9	3	37 9	18 2	23	41 7	36 16	1155 250	5.41 1.17
	· · · · I	····i	::::	·····i	106. Ankylostomiasis.				1							,													I	0.005
29 39	20 35		13 22	49 74	108. Appendicitis and typhlitis	70 85	11		6 16		1	1 2	1	1	3 7	8	1	1.					····i	2 2	2	2	5	6	119 159	0.56
5	4	8	1	9	110. Diseases of the intestines	5	1		2						1	1	1					1						2 .	14	0.07
2	1	2	1	3	111. Acute yellow atrophy of the liver														[								:::			
24 4	31		11	55 12		70 17	9	1 2	11		····i		2 2	1		3 2	3	····i				7	3	2	2	1	6	1	125 29	0.59 0.14
11	8	16	3	19	115. Other diseases of the liver	22	-4	1	2						2	1				1		1				1			41	0.19 0.009
1 13			9	27		11	3	4				3	2			4	ī					1	1				i	1	38	0.18
2	2	3	1	4	118. Other diseases of the digestive system (cancer and tuberculosis excepted)	/ 3	1	i l		l							1			1					1			1	7	0.03
80 511	70 419		53 234	150 930	119. Acute nephritis 120. Bright's disease	118 863	7 65	63	168	8	23	69	36	4 19	40	12 59	12 12	6 33	3 14	30				29	17	31		5 41 15		1.26 8.40
8	6	12	2	14	121. Chyluria	20			2						2	1									1	····i	3	1	34	0.16
14		1		19	123. Calculi of the urinary passages	12					1					i	····i			····i									3 31	0.01
					125. Diseases of the urethra, urinary abscess, etc.	3																							3	0.01
18 1			2 1	18 1	126. Diseases of the prostate 127. Nonvenereal diseases of the male genital organs.	29	2				,	1	2			3										1	1		47	0.22
					128. Uterine hemorrhage (nonpuerperal)																									
	4	3	1	4	129. Uterine tumor (noncaucerous)	11																							15 15	0.07
	8	3 5		8 10	131. Cysts and other tumors of the ovary	4			3			1				1									1	1			12	0.06
					male genital organs	17	2		1					1	1											2	2	1	27	0.13
	1	1		1	133. Nonpuerperal diseases of the hreast (can- cer excepted)							1																	1	0.005
	11		1 3	11 17	134. Accidents of pregonucy	12 9	2	2	····i						1 2	2										1		:::		0.11
	11	6	5	11	136. Other accidents of labor	6							1	î	ĩ	3					1			1	ĩ		1	1	17	0.08
	37 28	29 21	8 7	37 28	137. Puerperal septichæmia	23 23	10	2	3 3		1 1	1	3		2	3 2	2 2	1				1		3		1	3	2	60 51	0.28 0.24
	3	1	2	3		6			1					1															0	0.04
	4	2	2	4	140. Following childbirth (not otherwise de-	, i				1	1				i															
713	1		1	1	ficed)			1	1											1							:::	1	1	0.02 0.005
13 1	18	28	3	31	142. Gangrene. 143. Furuncle.	14		1	8			6	-1	• • • •										1	1	2		1 1	45	0.21
2			1	5	144. Acute abscess	6	1	٠					1		2														11	0.05
7	4	8	3	11		7			1							1													11	0.05
					cepted)	23	3	2	3			1	1															3	34	0.16
					rheumatism excepted)	2																							2	0.009
2	1	2		2	149. Other diseases of the organs of locomotion			: : : :																				1	2	0.009
							1																							

## TABLE C-Continued.

DEATHS IN MARYLAND, 1915, BY CAUSES, SEX, COLOR AND COUNTY.

																					_										
Male,	Female,	White.	Colored,	Total, Rural Dis- tricts.	CAUSES OF DEATH.	Raitimore City.	Allegany.	Anne Arundel.	Baltimore.	Calvert.	Caroline.	Carroll.	Cecil.	Charles.	Dorchester.	Frederick.	Garrett.	Harford.	Howard.	Kent.	Montgomery.	Prince George's.	Queen Anne's.	Somerset.	St. Mary's.	Talbot.	Washington.	Wlcomico.	ester.	Total: Maryland and Balto, City.	Percentage.
333 2 2 2 2 8 8 14 2 2 8 8 14 1 1 2 100 266 12 2 55 355 355 4 2 2 1 1 5 8 8 2 0 5 5 8 8 39 105 5 08 8 39 105	48 6 6 5 5 10 0 4 3 3 7 7 4 4 3 1 1 2 2 3 3 5 7 7 7 7 2 6 6 2 1 1 3 3 1 3 3 6 6 6 1 7 4 7 0 2 1 7 0 7 0	733 8 2 2 100 211 6 6 32 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3 3 3 15 14 147 147 147 148 188 240 4	81 88 13 24 46 46 46 17 19 19 19 19 19 19 19 19 19 19 19 19 19	50. Diabetes. 51. Exophthalmic goitte 52. Addisor's disease 53. Levelbernia. 54. Levelbernia. 55. Levelbernia. 56. According to the second sec	105 66 22 25 3 3 40 40 40 620 620 63 3 17 4 4 4 4 28 28 13 9 9 9 9 9 9 9	1 1 3 3 3 3 3 2 2 2 4 9 1 1 1 1 4 4 7 0 7 0	2 2 5 5 5 2 5 5 1 1 1 5 5 6 6 4 0 4 4	25 2 2 7 7 144 1 1 174 3 3 3 3 2 2 5 5 5 9 2 1 2 1 1 1	1 2 2 2 2 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	41 22 9 199 3 7	1 4 2 25 1 6 1	2 2 16 4 2 2 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1  3 3 1 1 13  8 3	4 1 1 3 5 5  4  2 4 4 5 8 2 0 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3 2 1 1 12 1 1	7 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 2 2 2 2 2 9 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 22 7 7		1 1 3 24 8 2 1 2 2		11	8	10 16 5	5 1  1 4  5  5 40 10 11 3 3 4  2 1 1  2 7 9	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1	188 14 5 28 40 8 71 4 11 17 97 23 63 67 1832 166 51 122 4 122 4 9 57 1 23 23 23 1855 103	0.87 0.07 0.02 0.12 0.19 0.04 0.83 0.05 0.08 0.45 0.07 1.26 0.50 0.24 0.31 0.02 0.24 0.31 0.02 0.24 0.31 0.02 0.04 0.31 0.02 0.04 0.04 0.05 0.05 0.06 0.07 0.07 0.07 0.09
13	22 4	24 4		35 4	rysm, etc. 82. Emholism and thrombosis. 83. Diseases of the veins (varices, hæmor-	145 20	2	13	55 8	1	1			2		10 3	1	5			3	1	6		5 1	1	12	1 2 .	- 1	320 55	1.50 0.26
1		1		1	rhoids, phlehitis, etc.)	2 3			.:			1								ľ							i i			6	0.03
3 1 58 29 283 304 9 21 3 14 3	3 4 53 37 223 267 8 15 	6 3 74 50 340 381 14 34 1 24 7	2 37 16 *166 190 3 2 2 5	2 6 5 111 66 506 571 17 36 3 29 8 22	55. Hiemorrhage; other diseases of the circulatory system anal fosses.  55. Diseases of the nami fosses.  56. Diseases of the threed hody.  59. Acroic bronchitis.  50. Caronic bronchitis.  50. Evenie bronchitis.  51. Bronchopacumonia.  52. Pienrisy.  54. Polinomary congestion, pulmonary apoplexy.  55. Gangrene of the lung.  56. Admina  56. Gangrene of the preparatory system.  59. Other diseases of the resultatory system.	4 3 72 54 504 605 18 30 8	2 10 8 31 26 2 3 3	1  8 1 39 40 1  2	2 2 15 10 *150 116 3 9 1	2 10 13	2 2 12 9 1	8 3 28 36 1 4	2 5 13 15 1 2	13 17  2	1 4 3 19 23 1 1 1	 1  6 5	2 3 12 8	1  1 7 2 12 17 1 1 1	1 9 16	4 1 16 11		3 16 35 2	13 15 1	3 4 12 14 1 I	7 1 10	6, 7, 19		1 2 13 15 2 1		6 	0.03 0.06 0.04 0.86 0.58 4.73 5.51 0.16 0.31 0.05 0.20 0.06
8 9 1	8	10 1		11 17 1	(tuberculosis excepted) 90. Diseases of the month and annexa 100. Diseases of the pharynx 101. Diseases of the œsophagus	9 5 9 1	1	2	,3 ,	1			1		2		3			3					ì	3	1		2	31 16 26 2	0.15 0.07 0.12 0.009



## TABLE C-Concluded.

DEATHS IN MARYLAND, 1915, BY CAUSES, SEX, COLOR AND COUNTY.

Male.	Female.	White.	Colored.	Total, Rural Dis- tricts.	CAUSES OF DEATH.	Baltimore City.	Allegany.	Anne Arundel.	Baltimore.	Calvert.	Caroline.	Carroll.	Cecil.	Charles.	Dorchester.	Frederick.	Garrett.	Harford.	Howard,	Kent.	Montgomery.	Prince George's.	Queen Anne's.	Somerset.	St. Mary's.	Talbot.	Washington.	Wicomico.	Total: Maryland and Balto, Oity.	Percentage.
523 3333 766 88 84 84 84 85 84 86 86 88 84 84 86 86 86 86 86 86 86 86 86 86 86 86 86	388 70 120 388 71 120 388 71 120 388 44 42 377 33 42 55	2 139 14 4 13 7 7 11 10 6 6 14 4 5 5 100 10 17 7 9 131 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 2 7 7 1 16 1 10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	636 114 15 204 14 14 14 18 13 23 33 31 11 17 7 7 60 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7	150. Congenital malformations (stillbirths not michodel).  151. Congenital debility, etterus and sciercum.  152. Congenital debility, etterus and sciercum.  153. Lack of care.  153. Lack of care.  154. Senility.  155. Suicide by poison.  155. Suicide by applyxia.  157. Suicide by applyxia.  157. Suicide by marging or strangulation.  158. Suicide by marging or strangulation.  159. Suicide by miching or piercing lostruments.  160. Suicide by miching or piercing lostruments.  161. Suicide by unting or piercing lostruments.  162. Suicide by unting or piercing lostruments.  163. Suicide by unting or piercing lostruments.  164. Poisoning by food.  165. Other acute poisonings.  166. Configaratioo.  167. Burus (configaration excepted).  168. Absorption of deleterious gases (configaration).  169. Accidental drowning.  170. Tranumatism by cutting or piercing instruments.  171. Tranumatism by flacture and the suicide of the suicide strains.  173. Tranumatism by machines.  174. Tranumatism by differents.  175. Tranumatism by other crushing (vebicles, railroad, landsides, etc.).  176. Injuries by animals.  177. Starvation.  178. Effects of bent.  189. Lighting.  181. Fracture cause not specified).  184. Fracture cause not specified.  185. Fracture cause not specified or ill-defined.  185. Cause of detain not specified or ill-defined.	3  4 9 5 83 10 58 7 2 91 10 109 3  1 12	111 111 5 5 2 1 1 1 1 1 1 1 1 1 1 1 1 1	30 14 9 9 2 3 3 1 1 1 1 1 3 3 2 2 2 2 5 	23 1099 177 1 1 2 2 8 4 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1	4	10 10 11 11 13 13 13 14 14	2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 2 6 6	2	13 13 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	8 1	1 8 8 5 1 1 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 		1 1 1 3 3 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3	2	1 2	2 10 7 7 7 6 6	51 9 11 14  2 1 1 1 6	1 2	5 299 229 192 193 16 16 16 16 16 16 20 3 255 11	2 5.24 7 1.25 1 0.10 3 1.40 0.20 3 1.40 0.20 3 1.40 0.20 3 1.40 0.20 3 1.40 0.20 3 1.40 0.20 3 1.40 0.30 3 0.40 3 0.30 3 0.30 3 0.30 3 0.40 3 0.40 3 0.40 3 0.40 3 0.40 3 0.40 4 0.40 5 0.40 6 0.
6386	5637	8618	3465	12,023	TOTAL	9327	822	740.	2700	153	284	566	373	267	502	780	228	390	199	274	436	523	273	400	270	314	751	424 3	54 21,350	

\*1 ('hinese, \*1 Indian. \*1 color unknown \*1 Chluese, †2 sec and 2 color unknown,



 TABLE D.

 DEATHS IN MARYLAND, 1915, BY COUNTIES, MONTHS, SEX, COLOR AND AGES.

		T					1	T	-T		ī				Ī	Ĩ					I						T		T	T	T	Т	П		
COUNTIES.	Japunery.	February.	March.	April.	May.	June.	July.	August. Sentember		October.	November.	December,	Male.	Female.	White.	Colored.	Total.	0 to 1.	1 to 5.	5 to 10.	15 to 20.	20 to 25.	25 to 30.	30 to 35.	2	2	45 to 50.	50 to 55.	55 to 60.	60 to 65.	65 to 70.	70 to 75.	75 to 80.	80 and over.	Unknown.
Allegany (*Y Indian). Anne Arundel. Anne Arundel. Baltimore (*Chinese). Calvart. Caroline. Ceel. Charles. Dorchester. Frederick. Harford. Howard (*1 Chinese). Keet. Howard (*1 Chinese). Keet. Sie Mary's. Taibot. Somerset. Si. Mary's. Taibot. Wiconico. Wiconico. Wiconico. Woorseter (*1 color unknown). Total 23 Counties. Total Baltimore City (*4 Chinese, I Japanese) Total Baltimore City (*4 Chinese, I Japanese)	233	9 449 2149 2149 2249 224 244 345 33 3 5 2 2 3 3 5 2 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	8 68 254 254 311 633 254 311 355 355 555 555 555 555 555 555 555	76 258 9 27 79 36 28 28 25 25 36 40 31 36 36 37 37 39 2 1,1722 899	52 54	47 588 190 111 10 39 222 16 38 63 31 12 30 16 42 22 31 44 48 48 33 19 63 31 19	764	63 269 1 9 9 30 50 50 50 50 50 50 50 50 50 50 50 50 50	20 25 35 35 31 22 35 31 22 31 31 31 31 31 31 31 31 31 31	7 23 33 31 25 40 68 15 20 12 26 30 43 22 45 27 43 31	73 48 182 8 18 27 23 49 49 28 29 24 42 21 31 13 26 51 28 29 860 755 624	86 66 252 12 25 44 31 31 58 82 22 20 39 39 42 23 36 78 83 44 28	453 399 1,520 899 132 305 200 131 261 373 109 208 117 134 224 286 146 148 133 176 *395 226 191 6,386 4,835	369 341 1,180 64 152 261 173 136 241 407 119 182 82 140 212 227 7 131 218 137 138 356 198 198 198 198 198 198 198		*49 375 *530 98 100 43 61 169 232 127 103 *73 145 165 226 130 166 139 156 139 136 3,405 *2,178		135 162 363 47 76 68 51 72 134 45 68 56 56 79 125 67 99 68 88 153 88 80		13 1 1 39 4 2 2 7 10 3 7 12 10 6 1 1 1 5 8 11 1 7 1 12	0 67 64 66 5 5 5 5 5 5 5 5 5 7 25 9 22 12 22 12 22 12 22 12 23 33 34 34 34 34 34 34 34 34 34 34 34 34	309	6 10 10 11 9 23 26 6 10 2 5 9 11 7 17 13 13 23 10 19			35 27 146 1 5 26 14 10 32 5 12 4 6 16 20 8 8 16 23 17 8 456 486	34 38 140 4 2 18 8 9 19 9 9 9 9 7 12 21 19 8 8 19 7 7 7 8 8 19 9 9 9 9 9 9 9 9 9 9 9 9 7 12 12 13 14 15 16 16 16 16 16 16 16 16 16 16	41 29 177 7 11 24 21 15 18 31 5 21 14 16 28 10 16 8 8 15 7 7 7 7 7 19 19 19 19 19 19 19 19 19 19 19 19 19	46 34 184 3 13 31 19 16 21 44 11 12 44 11 17 26 15 14 14 14 19 41 14 19 41 11 11 26 15 16 17 26 18 18 19 19 19 19 19 19 19 19 19 19 19 19 19	54 45 198 9 16 37 122 11 20 37 15 39 17 20 28 27 28 9 14 52 20 28 27	60 49 202 4 18 65 26 9 26 61 13 28 15 22 29 46 62 20 16 18 53 30 20 20 20 20 20 20 20 20 20 2	78 36 186 166 16 14 63 43 21 35 63 14 30 20 17 41 42 19 27 15 64 23 31 933 564 1,497	49 37 160 9 26 57 39 9 24 79 18 30 18 16 48 31 11 21 7 22 65 23 22 821 482 1,303	56 42 173 173 27 67 19 31 19 45 22 22 24 41 37 26 27 22 41 37 36 983 502 1,485	4 13 3 6 1 85 4 4 425 1 2 4 4 6 2 3 —81—81
																										-									

 TABLE, E.

 BIRTHS IN MARYLAND IN 1815, BY MONTHS, SEX, COLOR, NATIVITY, AND AGE OF PARENTS—RURAL DISTRICTS.

	10   10   10   10   10   10   10   10																																				-		,			-
	109   137   153   145   156   143   173   153   146   146   135																1		1	PARENT	s' NAT	IVITY.		. W									P	ABENTS	AOE,							
	180   137   103   145   156   143   153   153   146   145   156																		Ne	ative.		For	cign.	nkn				Age o	f Fath	er.							Ag	e of M	other.			_
Counties.	UNTIES.												Male,	Female.	White.	Colored.	Tota)		Both Maryland.	One Maryland.	Neither Maryland.	Both Foreign.	Father Foreign.	Both birthplaces u	15 to 20.	20 to 25.	25 to 30.	30 to 35.	35 to 40.	40 to 45.	45 to 50.	Unknown.	15 to 20.	20 to 25,	25 to 30.	30 to 35.	35 to 40.	40 to 45.	45 and over.	Unknown.	Mothers aged 14 years and under.	
Bultimore (*2 sex unknown); ?2 color unknown). Caroline Caroline Caroline Caroline Curroline Charles Dorriester (*1 sex unknown) Frederick. Garrett. Howard (*1 Chimese) Kest Montzomery (*1 sex unknown) Prince Georges (*1 Japanese) Sumerset (*1 sex unknown) Sumerset (*1 sex unknown) Sumerset (*1 sex unknown) Sumerset (*1 sex unknown) Talbot. Washington (*1 sex unknown)	99 257 297 47 02 49 62 122 31 36 27 79 33 70 42 33 131 66 40	110 220 33 60 37 44 80 120 45 28 22 66 74 31 56 38 37 115 66	107 218 21 46 72 32 38 84 100 40 40 77 47 37 120 60 65	228 15 49 58 44 42 60 47 26 31 71 73 20 62 49 128 65	156 93 197 24 38 63 33 82 111 48 44 23 16 53 73 34 64 44 35 121 25 54	143 94 257 236 36 61 41 43 73 118 47 35 20 30 60 80 36 52 32 37 124 48	153 113 251 19 51 63 45 45 61 104 41 50 20 27 63 63 80 36 56 56 56 56 56 56 56 56 56 56 56 56 56	119 267 249 49 777 511 39 96 84 53 60 32 34 50 105 35 67 45 38 8143	84 278 19 43 • 71 50 39 78 104 51 40 34 36 63 33 57 36 63 38 57	98 241 24 41 72 44 39 53 117 42 31 31 55 80 34 54 37 48	79 207 14 26 75 30 68 98 35 38 30 24 58 64 32 42 47 24 46	84 249 14 43 58 30 40 65 102 39 37 51 72 24 44 27 42 117 58	580 *1.478 *120 253 361 267 226 *446 626 248 192 149 *367 509 223 *359 242 232 *787	849 594 1,392 126 431 214 261 426 660 150 150 150 345 454 454 454 230 211 724 342 267	†2,64 9 36 74 43 20 51 1,13 55 41 23 20 48 61 25 25 41 48 33	51 413 225 55 155 155 255 255 25 25 25 25 25 25 25 25 25 25	13 1, 13 2, 13 13 14 15 15 15 15 15 15 15 15 15 15 15 15 15	550 498 342 334 712 963 403 701 472 443 511 697 546	795 777 1,868 239 314 674 429 762 278 375 270 265 474 545 552 423 360 863 542 376	507 119 431 6 126 77 108 40 68 192 161 79 46 50 112 170 42 81 170 42 81 112 170 42 81 113 81 82 81 82 81 82 81 82 81 82 81 82 81 81 81 81 81 81 81 81 81 81 81 81 81	305 53 154 1 40 27 30 8 18 74 28 17 12 104 193 8 20 4 4 18	99 171 235  13 8 24 7 6 4 20 25 5 6 21 10 25 1 1 20 3 1 3 1 3	51 2 33 1 123 5 9 9 5 1 9 9 1 9 1 9 1 1 2 7 1 1 2 7	5 1 9 2 2 7 2 2	15 15 24 5 11 16 13 10 28 17 7 5 9 8 25 8 25 8 8 25 8 8 30 9	313 2222 495 466 90 156 99 89 183 252 99 91 41 11 80 149 79 77 301 140 97	455 284 708 48 103 218 109 97 194 306 125 132 87 83 150 220 86 155 80 106 408 174 108	370 239 631 54 94 164 99 93 145 255 116 101 56 71 137 198 90 74 297 135 93	297 197 505 45 84 114 80 77 138 183 90 83 74 116 157 70 117 85 67 214 106	194 128 288 68 66 50 54 85 141 63 48 48 35 84 113 41 67 55 48 150 71 61	\$6 55 125 15 23 29 21 28 51 67 24 23 18 13 44 62 19 42 35 24 55 27 33	38 16 18 16 2 31 3	190 159 250 250 33 35 54 92 55 61 142 153 80 55 32 30 90 90 90 65 120 66 160 87 77	505 314 799 655 138 223 140 124 241 350 136 148 86 90 183 232 116 198 124 140 140 140 140 140 140 140 140 140 14	403 289 706 54 117 195 121 1123 194 296 127 119 72 75 145 250 84 150 103 166 118	189 569 51 87 144 79 77 142 223 98 96 84 53 142	289 151 363 32 69 87 71 113 175 75 52 248 47 89 136 51 91 81 53 190 86 78	51 151 5 28 41 12 24 25 71 28	11 6 18 5 7 3 4 2 12 1 1 1 1 2 5 10 3 3 1 2 10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	4 2 2 2 2 5 1 2 1 5 3 3 1 7 3 3 4 1 1 2 2	1 (14) 1 (13) 1 (14) 1 (13), 1 (14) 1 (13) 1 (14) 1 (14) 1 (14) 1 (14) 1 (14) 1 (14) 1 (14) 1 (13), 2 (14) 1 (13), 2 (14) 1 (13), 2 (14) 1 (14), 1 (14)	
Total Baltimore City (*2 cases of Chinese)  Total State of Maryland	1,123	1,129	1,221	1,018	1,054	1,143	1,269	1,181	1,167	1,138	1,103	1,162	7,028	0,680	11,46	1 *2,24	7 13,		,978 1	1,925	-	2,859 3,554							1,972 5,032	-	442 1	73 221	1,567 3,800	9,427	3,585 7,990		3,822	1,361	161	12	1(12), 2(13), 12(14)	

# DO NOT CIRCULATE





DO NOT CIRCULATE

